polymorphonuclear cells. No tubercle bacilli were to be found. On November 20th, there being still no improvement, 10 c.cm. antimeningococcal serum were injected hypodermically, the dose being repeated on November 23rd, 26th, and 27th, and on December 1st respectively. No definite improvement was apparent until after the last injection, when the temperature began to fall rapidly, consciousness returned, and all the serious symptoms abated. This satisfactory condition of things continued for five days, and then a relapse occurred, lasting for a further period of five days, during which the old symptoms were revived, though less acutely, with the temperature again shooting up to 103° F. It was then decided to inject 10 c.cm. antimeningococcal serum into the spinal canal, but an unexpected amelioration in the child's condition prevented this intention being carried into effect. From this time on recovery was uninterrupted, and here you see the boy apparently as well and healthy as he ever was. On January 5th another lumbar puncture for confirmatory purposes was done. The spinal fluid was then found to issue under a much diminished pressure; when centrifugized the deposit contained a few blood discs but no excess of leucocytes and no micro-organisms, and media inoculated with the deposit revealed no growth of the *Diplococcus intracellularis*.

I show you here the small flasks of antimeningococcal serum as they are sent out from the laboratory ready for use. Each flask contains 10 c.cm. We propose to inoculate the child we have examined this afternoon, in the first place hypodermically, and later on, if need be, by the intraspinal method.

It may be mentioned that in cases of tuberculous meningitis the fluid withdrawn by lumbar puncture, though turbid, is often sterile, but on centrifugalization tubercle bacilli can generally be demonstrated in the residue. The prevailing lymphocytes are of the small mononuclear type.

In suppurative cases the polymorphonuclear cells are in abundance, and the finding of the streptococcus, staphylococcus, pneumococcus, or other micro-organism in the centrifugalized residue affords confirmatory evidence of the type of meningitis which is under observation.

AN IMMUNIZING SUBINFECTION IN SCARLATINA.*

By A. D. EDWARDS, M.B., B.S.LOND., B.Sc., D.P.H.,
DEPUTY MEDICAL OFFICER OF HEALTH, BOURNEMOUTH.

THE continued investigation of scarlatina has given us proof abundant that the predominant factor in the spread of the infection is direct contact with persons suffering from the fever. It is true that some outbreaks of the disease are traceable to infection-bearing milk, and also that a certain number of cases may be due to infection from clothing—things as distinct from persons; but the predominant factor is usually recognized to be personal contact.

The more extensive supervision of contact cases and the routine examination of school classes in which scarlet fever has occurred have strengthened this belief, for they have resulted in the wider discovery in the affected classes of children who were passing through attacks of scarlatina so mild that they had been missed alike by parents and teachers, but who were acting as a continued source of infection to others.

Missed Cases.

These cases are "missed" because the parent or teacher did not see the rash on the child; the sore throat was so slight that the child was well in a day or so, and no medical man was called in; or a medical man may have been called in, and either because the rash was very faint or absent, or because the rash had faded, a correct diagnosis was not made. The attack is a mild one; the child returns to school, and, especially if there is any ear or nose discharge, acts as a means of spreading infection; in about ten days the child is found to be peeling. I would emphasize the importance in class outbreaks of a full inquiry and examination of all children who have been absent from school on account of sore throat. Whether or not peeling is a source of infection is doubtful, the evidence is apparently in the negative; but it at least gives information as to the source of infection in a fresh case which may have puzzled the investigator.

Mention has been made of the evanescent rash with very slight symptoms, and of scarlatina *sine eruptione*. The

existence of the latter has been regarded as doubtful, its reputed occurrence having been ascribed to the fact that the rash, being slight and evanescent, had been missed.

The Lower Grades of Infection.

But because scarlet fever infection varies so greatly in its characteristics as to produce all grades of rash, from the brilliant and vivid eruptions, hæmorrhagic and accompanied by miliary vesicles, to a slight erythema of low punctiform character which is hardly visible, it is not illogical to expect that the rash-producing properties of the infection may be traced to vanishing point.

In examining large numbers of children who were school class contacts of true cases of scarlet fever, I have found on not a few occasions that some of these children had no sign or symptom of scarlet fever except a condition of the tongue in which the papillae are well marked, with possibly a slight injection of the throat. In some of these children the surface of the tongue was furred, giving the "white strawberry" appearance; in others the "red strawberry" appearance was present. It has been my custom in dealing with school classes affected by scarlet fever to exclude any child who had a sore throat or any skin condition which might indicate the early stage of a scarlet fever rash, as well as any child who had ear or nose discharge. The children who showed this tongue condition were not excluded; they were watched, and in no case was I able to detect any other sign or symptom which would verify the diagnosis of an attack of scarlet fever. In tracing other cases of scarlet fever which subsequently occurred in such a class, I found that occasionally the tongue cases were possible, but rarely probable, sources of the infection of the fresh case.

The Scarlatina Tongue.

The strawberry tongue of scarlet fever is due to the following sequence of events: First, a thick white fur accumulates on the tongue, due to an excessive proliferation of epithelium; then this fur begins to exfoliate, the fungiform papillae showing through as prominent congested points, giving the appearance of the white strawberry tongue. By the fourth day the fur has exfoliated, the papillae are very prominent and congested, and the appearance of the red strawberry tongue so typical of scarlet fever is produced. The white strawberry tongue is by no means diagnostic of scarlet fever. It occurs also in conditions such as measles and diphtheria; but the red strawberry tongue is said to be pathognomonic.

An Immunizing Subinfection.

Many of the contact-children to whom I have referred had the red strawberry tongue, but showed no other sign of scarlet fever at the time; and in none of the cases which were kept under observation was there any subsequent sign to verify the diagnosis of scarlet fever in any of its stages. But all these children had been in contact with scarlet fever cases; and of the other children in the class who had also been in contact with these cases, but who did not show this tongue sign, some developed scarlet fever. It is not unreasonable, therefore, to suppose that these children who had the tongue sign may have had scarlet fever in so low a degree of infection that whilst it was too attenuated to produce any visible effect upon the more opaque and resistant epithelium of the skin, it was concentrated enough to produce an effect on the more delicate epithelium of the tongue; and, further, that the attenuated infection by which they had been attacked had immunized them against the more marked disease.

Another fact which appears to indicate the frequent existence of a scarlatinal infection of very low virulence is the occurrence in every outbreak of scarlet fever of sore throats among contacts who do not actually develop scarlet fever. The prevalence of these sore throats is, I believe, greater at the beginning of an outbreak of scarlet fever. My records are not complete enough for proof, but it is recognized that the autumnal outbreaks of sore throat are coincident with the onset, but not markedly continuous with the progress, of the autumnal rise in scarlatinal infection. This would support the belief that a high percentage of persons in contact with scarlet fever are infected, but many have so slight an infection that it causes no sign but sore throat and may immunize the individual from the greater infection. The infection of

* Read to the Bournemouth Medical Society.

persons, therefore, during the prevalence of scarlet fever may be classed as follows: (1) The well developed cases of scarlet fever; (2) the cases which are mild, but whose infection has resulted in sore throat and a slight, evanescent rash with subsequent peeling; (3) the cases in which there is a still milder grade of infection, not sufficiently concentrated to produce a skin rash, but which produces a sore throat and may produce "exfoliation" of the tongue. I have mentioned that in none of these latter cases which I have followed up has true scarlet fever developed, but the record of a very much larger number of cases is necessary to prove the hypothesis.

Tentatively, therefore, I hold it true that Class III, composed of persons exposed to scarlatina who have developed nothing but a sore throat and an unrecognizable degree of scarlatina, have received an immunizing subinfection which has protected them from the full attack. This subinfection has produced a response in the form of increased resistance, the host becomes better armed with antibodies, and a "positive phase" is produced in the same way as it is produced in vaccine therapy.

The possibility of this increased immunity being produced is strengthened by the fact that in persons exposed to infection, but who do not contract the disease, a definite leucocytosis may be demonstrated.

The Russian Vaccines.

In 1906 Gabritchewsky of Moscow introduced a vaccine made from the *Streptococcus scarlatinae*, which he used with a view to protecting children from attacks of scarlatina. As a result of the injection he obtained in a certain number of cases an erythema resembling that of scarlatina. Zlatogoroff and Langoway in Moscow obtained similar results, there being in some cases a slight rise of temperature and pharyngeal injection. These results are not conclusive, but at least they are in favour of the specific nature of the *Streptococcus scarlatinae*. The further research going on at the present time should give us more definite knowledge concerning the elusive streptococcus, but, in view of our present lack of knowledge concerning the causative micro-organism, it becomes necessary to seek an analogy in the case of specific bacilli concerning which we know a little.

An Argument from Analogy.

No practical work on the typhoid bacillus can be complete which does not take into consideration the group of closely related micro-organisms including the *B. coli* and the bacillus of Gaertner. There is evidence of a series of bacilli of gradually decreasing virulence, some of which can be surely recognized by definite culture reaction, and there is moreover a slight variation in the type of disease produced which has resulted in the description not only of typhoid fever, but also of paratyphoid fever. Again, a clear analogy is afforded by the variation of the virulence of the diphtheria bacillus, from the most toxic infection through all grades to the vanishing point at which the person harbouring the bacilli becomes merely a carrier.

The variation in the degree of infection produced depends to some extent on the natural resistance of the person attacked, but that it depends also on the absolute virulence of the micro-organism is evidenced by the varieties of the diphtheria bacillus, the pathogenic properties of which are traceable from the Klebs-Loeffler bacillus of high virulence, through the apparently inert Klebs-Loeffler, to the non-toxic or only to become toxic pseudo-diphtheria bacillus.

The Public Health Standpoint.

If the subinfections of scarlatina are immunizing, then they are of advantage to the individual. From the public health point of view it must be asked, "Are they sources of infection to others?" If so, they may be of disadvantage to the community, for the infection which will have only a protective and immunizing effect on one person may, when transferred to other persons of low resistance, result in graver pathological conditions. If these subcases of scarlet fever are infectious, it would appear that usually their infectivity is of low degree; the infection is not likely to produce otorrhoea or rhinorrhoea, but if these discharges are coincident, then the infection is likely to be spread. This would lead to the conclusion that every effort should be made to cure such discharges, especially in school children who come into close contact with each other, and

herein, as in so many other directions, there is hope of much benefit from the institution of routine school medical inspection.

ON THE RESULTS OF THYROID TREATMENT IN INFANTILE WASTING.

By J. W. SIMPSON, M.D., F.R.C.P.E.,

ASSISTANT PHYSICIAN, SICK CHILDREN'S HOSPITAL, EDINBURGH.

ON examination of thyroid glands taken from at least 100 children dying of various diseases, pathological changes were noticed in a large number. The fact that these changes were especially constant in children who had died in a "marasmic" condition led me to inquire if this state of the thyroid might not play some part in the all too common condition of infantile wasting. In a former paper¹ I described the pathological changes seen, and also the results of treatment of 20 marasmic cases with small doses of thyroid extract. These results were decidedly encouraging, but the number treated was too small to make them at all convincing. Since that time I have had the opportunity of treating a much larger number of wasting cases (80), and this second paper deals with the results obtained from these.

In cases of infantile malnutrition the history is usually one of improper feeding, and this being so, one frequently finds that when put on a more suitable diet the children at once begin to improve, without the addition of any drug. All my cases were therefore at first carefully dieted for at least two or three weeks, and thyroid was only given to those who showed no signs of improvement after this time.

In dieting, some modifications of milk, such as peptogenic milk, peptonized milk, milk and cream mixture, whey and milk mixture, etc., according to the conditions present, was first tried. I found, however, that the diet which on the whole gave the best results when thyroid was being administered was a diluted milk and cream mixture treated with sodium citrate (1 to 2 grains to the ounce of milk). Cream was usually added after the first two or three days, beginning with half a teaspoonful to the bottle. Why thyroid reacts better with this diet than with the others it is difficult to say, but it may be due to the alkaline character of this milk mixture, as in cases of wasting there is frequently some intestinal disturbance which leads to the presence of fatty acids, and this acid condition of the intestinal tract may prevent the absorption of the drug.

Results in Cases Treated with Thyroid.

Out of a total of 80 cases treated with thyroid, 72 were infants under nine months, and their history was usually one simply of wasting. In the 8 over this age the wasting had supervened after some acute disease. Out of this total (80), 12 died and 5 did not return to hospital after treatment had been carried on for two or three weeks. Of the 12 cases which ended fatally, 2 were extremely ill when first seen, and probably the stage of their illness was too advanced to allow of the administration of thyroid being beneficial. In both these children the weight was under 4 lb., and both showed evidence of failing circulation, oedema of legs, etc. One died two days, and the other four days, after being first seen. Two others were cases of marasmus due to congenital syphilis; of the other 8 deaths, one of which improved for some little time, but eventually died of tuberculous meningitis, 2 were country cases and were only seen a few times, whilst the remaining 5 did not react to treatment in the slightest degree. The remaining 63 cases, however, all did well. No doubt a number of those infants would eventually have improved on dieting alone, but the fact remains that they failed to put on weight for the first two or three weeks on such treatment, and that on the addition of thyroid (the same dieting being continued) a number of them at once increased rapidly in weight, while some who remained stationary for a week or two showed beneficial results after that time.

In the cases treated some interesting points were noted. One specially instructive was that of the children of a woman who was suffering from well-marked chronic phthisis. Five years ago she brought her infant (aged