INTRATHYROID HAEMORRHAGE FOLLOWED BY ACUTE DYSPNOEA AND DEATH.

BY

S. GURNEY CHAMPION, M.D., C.M.EDIN.,

AND

C. B. MOORING ALDRIDGE, M.A.CAMB., M.R.C.S.ENG.,

MEDICAL CASUALTY OFFICERS, LOWTHER ROAD BRANCH, ROYAL
VICTORIA AND WEST HANTS HOSPITAL, BOURNEMOUTH.

As the following case is of rather an exceptional character, its narration may perhaps prove interesting.

The patient was a domestic servant, aged 24, who had been employed for the previous three weeks at the Lowther Road Hospital, Bournemouth, and who jumped off a tram travelling about eight miles an hour at 8 p.m. on Thursday, April 20th. She fell on the left side of the face, receiving two small cuts, one on the outer side of the left orbit, with a marked surrounding ecchymosis, the other on the under side of the chin; there was comparatively little bleeding from either of these cuts.

When helped up she was perfectly conscious, and acknowledged that the accident was due to her own carelessness. She remounted the tramcar with the assistance of the conductor, and was taken to a doctor's house twenty yards away. The doctor being busy with other patients at the time, the girl's injuries being apparently trivial, the hospital where she was engaged being only a few hundred yards off, her appearance not bad enough to excite alarm, and she herself professing to be quite well, she was allowed to return to the hospital about five hundred yards distant.

Soon after she commenced walking more urgent symptoms appear to have arisen, for, a quarter of an hour later, she was found on the doorstep of the hospital in a collapsed condition, breathing with great difficulty and clutching at her throat.

Her noisy inspirations were heard at some distance and attracted the attention of the sister in charge, who, with the nurse, came to her help. It was during the next few minutes that she related the history of the accident.

Within five minutes one of us was present, and realizing the urgency of the symptoms, telephoned for the other's assistance.

State on Examination.

We found the girl lying in a supine position, with head slightly raised on a pillow; she was unconscious. The noisy inspiratory dyspnoea was then the most pronounced symptom; each inspiration was accompanied by a throwing forward of the neck.

The thyroid gland was markedly enlarged, and we understood that she had previously suffered from a simple goitre, which had given her very little trouble; there was no localized area of fluctuation over the gland.

Her face was very pallid; the pulse was regular, rate 100, and small tension. The pupils were equal, somewhat contracted, and did not react to light. She had had incontinence of urine, but not of faeces. The skin was moist.

There was no exophthalmos, no marked flaccidity or rigidity of the limbs; there were no convulsive movements of any kind.

Treatment and Result.

We did everything that we thought suitable for relieving the profound shock present and the urgent dyspnoeic symptoms. Oxygen was freely given, but the heart-beats became steadily more rapid and more feeble, reacting for a short time only to the various remedies. There was no vomiting, no haemorrhage from the nose, mouth, or ear, and no escape of cerebro-spinal fluid. The pupils continuing equal, became slowly dilated and insensitive to light.

Finally she died at 9.30 p.m., an hour and a half after the accident, respirations having continued after the heart had stopped.

Necropsy.

At the necropsy held two days later the following condition was found: The body was well nourished. On one of the legs was the scar of an operation for varicose veins. On removal of the skull cap the cortical vessels of the brain were found to be much engorged, but there was no fracture or intracranial haemorrhage; both middle meningeal arteries were intact.

The examination of the chest revealed a healthy heart, with competent valves; there was no effusion of blood present. The abdomen was normal; the various organs were found to be healthy.

On the examination of the neck the thyroid gland was found to be much enlarged, the main bulk of the gland being formed by a haemorrhagic cyst with very definite walls, chiefly occupying the middle line, but also encroaching on the lateral lobes. Its approximate measurements after escape of its contents are 7.5 cm. by 4. This thyroid haemorrhagic cyst was situated low

down, between the upper portion of the manubrium sterni in front and the lower cervical and upper dorsal vertebrae behind. There was no extravasation of blood into the surrounding tissues.

The thyroid gland itself was much diminished in size after the escape of blood. It weighed 5½ oz., against the normal 1 to 2 oz. There was no obliteration or displacement of the trachea.

One or a combination of the following theories is suggested as a cause of death:

(a) Death from double shock—first the shock of the accident due to the violent impact of the head upon the ground, and, secondly, the shock of the sudden onset of dyspnoea.

(b) From the direct pressure on the trachea during life, this cause not being apparent *post mortem*.

(c) From thyroid poisoning due to the sudden tearing of the glandular substance, thus liberating an excessive quantity of thyroid secretion into the general vascular system.

(d) From pressure on the vagus nerve producing an inhibitory effect upon the heart.

(e) From pressure on the middle and inferior cervical sympathetic ganglia with a reflex effect upon the heart via the cardiac nerves to deep cardiac plexus.

The following cases are interesting with reference to the above, but in none of them is it related that the dyspnoea was preceded by a severe accident.

CASE I.—In the BRITISH MEDICAL JOURNAL for May 1st, 1909, there is a case of a Chinese male, aged 25, who had a sudden enlargement of the thyroid with dyspnoea. This man recovered without operation in about three days' time.

In the *Lancet*, November 16th, 1907, Berry narrates the following cases:

CASE II.—That of a boy of 17, who had chronic severe dyspnoea due to a thyroid. Postponed operation till it was too late. Very little blood was lost; but just after the thyroid tumour was removed, he collapsed and died.

CASE III.—Another in a woman, aged 68, who, after the operation of removal of the thyroid gland for the relief of dyspnoea, died from cardiac failure without any sign of respiratory distress.

CASE IV.—Still another one is mentioned of a girl of 14 who had a bilateral parenchymatous goitre, noticed only three weeks. On Berry's arrival the girl was deeply cyanosed and unconscious. Fortunately he had everything ready for operation. Thyroidectomy was performed; the patient recovered, not regaining consciousness, however, for one hour after the termination of the operation.

CASE V.—He relates, too, a case of a young woman with thyroid enlargement attending as an out-patient at one of the London hospitals who died on the doorstep of the hospital from sudden dyspnoea. *Post mortem* there was found to be present an extensive haemorrhage into an encapsulated tumour of the thyroid.

CASE VI.—Osler, in his book on *The Principles and Practice of Medicine*, fifth edition, mentions a fatal case from haemorrhage into the thyroid gland and cellular tissue.

The cases mentioned above nearly all occurred in young people. Monnier says that the prognosis is very bad in urgent operations for dyspnoea resulting from sudden enlargement of a thyroid gland.

THE DISCRIMINATION OF COLOUR.*

BY

F. W. EDRIDGE-GREEN, M.D., F.R.C.S.,

BEIT MEDICAL RESEARCH FELLOW.

(From the Institute of Physiology, University College.)

In a paper on the relation of light perception to colour perception,¹ and in previous writings,² I have stated that if a portion of the spectrum be isolated, it will appear monochromatic, the length of the monochromatic region varying with the intensity and wave-length of the light and the colour perception of the observer. Most normal-sighted persons make about eighteen such divisions in a bright spectrum.

In a paper in the *Proceedings of the Royal Society*,³ Lord Rayleigh, whilst agreeing that the facts were as I stated in the conditions described by me, expressed the opinion that he could distinguish between the wave-lengths included in a monochromatic division to the extent of discriminating between the colours of the two D lines. Lord Rayleigh kindly lent me the colour box with which he had made the experiments, and on repeating them in the manner described by him I arrived at similar results. I hope, however, to be able to show that the results obtained by Lord Rayleigh were due to the

* Read before the Royal Society.