Whatever the nature of this remarkable case might have been, I wish again to draw attention to the persistent cervical stiffness and morbus cordis with arthritis.

The last case merits a complete account at some future date, but I append these few facts:

A. H., male, aged 6 years, came first under the care of the surgical side of the hospital, having suffered for fourteen days from pain in the back, and limitation of extension of the cervical spine. As a result of this his head was bent somewhat forward. Four weeks later both legs, but particularly the right, became weak, and both knee-jerks were difficult to obtain. Spinal caries with commencing paraplegia was suspected, and he was put in a Phelps's box. In September, 1908, while in the box, I saw him for what appeared to me an attack of rheumatic fever with multiple arthritis and heart disease, and after repeated examinations I felt doubtful of the presence of spinal caries. A year later he had another attack of what appeared to be rheumatic fever with pericarditis.

Once more I went over the case from the point of view of spinal caries, and came to the conclusion that it had probably not occurred. At this time there had been considerable pain in the shoulders with marked atrophy of the right fingers and hand, with a less degree of the same condition in the left. There was also some wasting of the right leg. Over the upper extremities tactile sensation was diminished. Both hands were blue.

It would seem, then, possible that we have here a smouldering rheumatic inflammation of the cervical fasciae and vertebral joints with a spread of the same process to nerve roots in the neighbourhood.

The outcome of my last observations comes to this: that if we study some of the stiff necks in childhood associated with rheumatic heart disease by the light we may get from studying rheumatic processes in some obvious situation such as the elbow or the knee joints, we may discover that the simulation of spinal caries by this disease may be a more frequent occurrence than we have perhaps imagined, and we may learn something of the clinical history of spondylitis rhizomelique.

This concludes my lecture but does not exhaust my subject, for there is much that is of interest in the study of renal disease and of cerebral disease and disorders of nervous function to be learnt from the study of rheumatism in childhood.

FOUR CASES ILLUSTRATIVE OF CERTAIN POINTS IN PSYCHO-ANALYSIS.

BY

HUGH WINGFIELD, M.D.CANTAB.,

CONSULTING PHYSICIAN, ROYAL HANTS COUNTY ASYLUM.

ALTHOUGH the theoretical side of Freud's work has attracted some attention in this country, the practical possibilities resulting from an application of his principles seem to have been almost entirely neglected. Possibly this may be due to the difficulty and the time so frequently involved in carrying out the process of psycho-analysis; but it does not appear to be realized that the practical use of his principles is often a comparatively easy matter, and of so immense a value, in a large number of "borderland" cases, that nothing else holds out any real hope of relief.

The great fact which he pointed out is that many of the symptoms of hysteria (such as pain or paralysis), and still more the mental symptoms of certain allied conditions (such as the strange phobias of neurasthenia or psychasthenia), are not merely haphazard occurrences, but are produced by emotional ideas, actually and vividly at work, though in some unknown region of the mind. These ideas are, in all cases, the result of some painful impression which, like all other impressions, could for a time be readily recalled to memory, and indeed could only with difficulty be dismissed from consciousness at all. Later, however, sometimes perhaps in the natural course of events, and certainly sometimes by the agency of a determined effort to forget, the memory of the painful event has become blurred, and has finally faded beyond recall.

In certain conditions, which I need not here describe, these forgotten painful impressions remain permanently fixed in that part of the mind which is often called the sub-consciousness, though there are, perhaps, serious objections to the use of the term. Yet these hidden ideas never manifest themselves as they really are in all their nakedness to the consciousness of the unfortunate sufferer. They may, for instance, give rise to vague but intense feelings of

terror, while the patient remains quite ignorant of the causes of his fear. The "sub-consciousness," however, is perfectly aware of what it feels, but merely infects the waking consciousness with a feeling of obscure terror, keeping entirely to itself the secret object of horror.

The first case given below illustrates this characteristic rather well. The patient was a prey to terrible foreboding, but never at all suspected that the cause was a fear that he would die the same appalling death as he had witnessed in his own father.

Again, we may get hysterical pain, which appears to be often a kind of defence-neurosis, the unendurable sub-conscious mental agony being converted into the more bearable physical suffering.

The particular part of the body attacked by hysterical pain is often decided by some accidental condition in the patient at the moment of the original shock of anguish. Thus, in Cases II and III, the patients were respectively suffering from slight left facial neuralgia and from aching in the left arm at the moment when the blow was struck. Both patients developed hysterical pain, one in the left side of the face, the other in the left arm.

To cure such cases it is, as Freud shows, absolutely essential to discover the submerged idea or ideas which cause the mischief. According to him, when the ideas are discovered, and discussed, the manifestations will cease, the idea dissipate and become harmless, restoration to the memory of the original mental trauma, and the discussion of it with another person, producing what he calls "abreaction," though the actual mechanism by which this process effects a cure is still a matter of dispute, many disagreeing with the theories advanced by Freud.

In Case II this result certainly occurred; but, as I had constantly assured the patient that the pain would vanish if she could tell me everything, I cannot say how much was due to suggestion and how much to "abreaction." At any rate, in Cases I and IV the mere discovery of the original cause of the condition did little or nothing to relieve the patient's condition; but suggestion during hypnosis directed against the ideas, when they were brought to light, produced instant amelioration.

I wish to emphasize particularly the point that (except in Case III, where the cause was known at the first sitting, and where, therefore, suggestion succeeded at once) though hypnotic suggestion had been tried over and over again before the original trauma was laid bare, such suggestions proved of no real effective value whatever, being directed merely to the removal of the symptoms.

Whilst in every one of these cases hypnotic suggestion was employed, it was used with a twofold object—first, to unearth the actual emotional idea which caused the disorder; and, secondly, to remove or counteract that idea when discovered.

The first case occurred many years ago, long before psycho-analysis was recognized, and, being then in general practice, I had no opportunity of making similar experiments.

The patient was an American gentleman, aged 36. He came to me complaining of intense depression and an overwhelming dread of some terrible impending catastrophe. He had suffered in this way for many years, but the depression and terror had increased rapidly for some months before he came to me, and he had strong impulses to suicide. His own medical man had died some six months previously, and perhaps the increased severity of his distress may have been due to the fact that he had not confided his state to any one else, but had kept it entirely secret. He was, I found, readily hypnotized to a slight degree, though not capable of deep hypnosis, and for nearly three weeks I tried giving him suggestions that his dread would vanish. The treatment, however, had no effect; he became, indeed, rather worse than before. He had no idea as to what he apprehended; and I was about to give up the case in despair when I thought that he might possibly tell me the cause of his terror by means of automatic writing. I found that he had on several previous occasions tried planchette writing, so I made him place his hand on a planchette, and asked what frightened him. The planchette wrote, "Father's death." Asked why this event frightened him, he wrote, "Will die same." It appeared that when a child he had witnessed his father's death from pulmonary embolism, and had been most painfully impressed.

Having thus discovered what he subconsciously dreaded, I immediately began making suggestions under hypnosis that it was impossible that he should die in this way, that his heart was absolutely sound, and other statements of the same nature. He began to improve at once, and in less than three weeks there was no trace of his former condition.

The remarkable point is, of course, that the earlier general suggestions of improvement had no effect, but that

suggestions directed against this particular fear were immediately successful. It is worth noting that even when the patient was told what his peculiar foreboding was he could not be brought to believe it, so utterly was it absent from his waking consciousness.

The second case was that of a young lady of 24. For five years she had suffered from pain in the left side of her head, which for eighteen months had been gradually increasing in intensity. When I saw her first, in October, 1909, she was obviously in a state of acute suffering, and was considerably wasted. The pain I concluded to be hysterical, and tried suggestion under hypnosis. She was a fairly good subject, but the result was practically nil, for, though during hypnosis I was able to diminish the torment, it returned as fiercely as ever within half an hour of waking. After thirteen trials with similar consequences, I determined to search farther for the actual cause. During hypnosis I told her she would dream of something connected with her illness. On the third subsequent trial she dreamt she saw a glass of water on a table. Her pain was increased when she was roused, so that I felt that probably the dream was intimately concerned with it. On following occasions she had other dreams, and at last she suddenly identified the scenes of these, but resolutely refused to give me any more information. Things remained thus for nearly three months, when, owing to my constant reiteration of the fact that I could do her no good unless she gave me her confidence, she at length told me the following history.

Rather more than five years previously she had been staying with friends in South Africa, and had at first sight fallen violently in love with a young man who was a frequent visitor to their house. Her hopes were suddenly dispelled only ten days after their first meeting by finding that he was already engaged. At the time when she made this discovery she was suffering from slight neuralgia of the left side of the face.

When she had finished her recital, she conversed about her experience, and asked when she should come to see me again, whereupon I told her to let me know how she was in a few days. Three days later she wrote saying that the pain had entirely vanished, and that she was quite well. I have not been able to follow up her case, for she left four months later for South Africa, and I have not heard of her since.

The case conforms completely with Freud's theory, the mere recalling of the fact to the conscious memory and the discussing it with someone else being sufficient to dissipate the emotional idea which formed the core of the mischief. The glass of water appeared as a crucial incident in the story. One day the young man called and asked for a glass of water. He drank it, filled himself a second glass, and drank half of that. After he had gone, the patient, when she was alone, had drunk the remainder. The glass had a key pattern engraved on it, and so had that which appeared in her dreams. The localization of the pain in the face was doubtless caused by the neuralgia, which was present when the blow fell. The continuance of this pain was a defence-neurosis, the mental being converted into physical torture.

The characteristic localization is reproduced exactly in the next case, which in many respects presents identical features.

Miss X., aged 23, was sent to me by Dr. Risien Russell, suffering from hysterical pain in the left arm, which was sufficiently acute to incapacitate her for work of any kind. For thirteen months she had been treated, first in a home, and latterly in a hospital. The pain, she said, had become less severe, but was still fairly acute, and occurred at frequent intervals, both day and night. It had commenced in 1908, and I at once inquired of her mother whether she had had any love disappointment at that time. It transpired that she had, the circumstances being very like those in the previous case. But she had always recollected it keenly, as was natural, the attachment having lasted some years before the man in question had become engaged to another woman. She herself told me a little about her trouble, and I then hypnotized her and suggested that the remembrance should no longer distress her, and that the pain should cease. It vanished immediately, and I saw her on three other occasions during the ensuing week. The pain remained in abeyance, and I fondly hoped that it would never return. But two months later it suddenly revived, and she again came to me. On inquiry I found that the young man had written to her, and that the pain had seized her the same evening. I now had an opportunity of discovering why the suffering chose the left arm as its seat. It appears that one morning she had been cleaning out a large aviary, which labour had made her left arm ache considerably, and that it was at this moment that she received the intelligence which so upset her. The pain was much more intractable than on the first occasion, and I was obliged to see her no fewer than seventeen times before it completely vanished. The interest of this case lies in the localization of the pain, and also in the fact that the prime cause of distress seemed to exist in the conscious memory, so that psycho-analysis by an elaborate method was unnecessary. But it is so curiously like the last case that the comparison of the two seems instructive.

The next case was also sent to me by Dr. Russell, and

though not yet completed, is, I think, sufficiently advanced to give as it stands.

Miss X., aged 21, had for some six years suffered from a feeling of dread while in church, and also if alone in the street. The terror in church was just as bad when she had a friend with her as when she was alone, but out of doors the presence of a companion prevented an attack. She had no idea as to why she was afraid, but felt that people were looking at her. Suggestions of improvement made under hypnosis produced no real effect. After trying these for three weeks, I suggested, under hypnosis, that she should remember the original incident which caused the fear. This she could not do, but she became aware that something, she knew not what, had occurred before her first attack, which happened during the singing of a hymn when she was at school. I then suggested that she would recollect it during the day, or during sleep that night. The same afternoon she recovered the incident whilst she was resting. It appears that when a child she had been reading a story book, in which she became completely engrossed. In the course of the story one of the characters suddenly became mad, and was taken off to an asylum. On reading this the child shut the book, in a crushing sense of horror, and, though she spoke of it to no one, she remained terribly distressed for several weeks. The incident had completely faded from her mind, and could be unearthed only by some psycho-analytic method. I found that she frequently dreamt that she had gone mad, and was being seized by attendants to be carried off to an asylum. Clearly, then, what she unconsciously dreaded was the onset of insanity, and her terror was due to the idea that people thought she was going mad, and were ready to hurry her off to imprisonment.

Under hypnosis I at once began to make suggestions to combat this idea, and the result is that for the first time for five years she can now go about alone. I have not yet allowed her to try attending church, but I hope to do so shortly. The improvement was marked even after the first counter suggestion.

Of course, in cases of this kind, there is often, in addition to the actual subconscious fear, a secondary fear lest an attack should come on, which is not subconscious, and which can frequently be eliminated only by actual experience of lasting immunity. That is why I have, so far, forbidden her to try church-going. The points of interest in the case are these: First, suggestions of improvement were of no value whilst directed only against the general feeling of fear. Secondly, the mere recalling of the original shock to her waking memory did not of itself alter her condition. Thirdly, suggestions specifically made against the actual emotional idea when it had been discovered produced immediate improvement.

Cases of the kind described above seem to show the paramount importance of a really accurate diagnosis in many such instances, if we hope to cure them at all. Sometimes, as in the second case, the actual diagnosis and discovery of the hidden cause is sufficient to effect a cure. For other patients suggestion is needed, but is often of use only if directed against the actual morbid subconscious idea. And, as a rule, that idea must be discovered by some process of psycho-analysis, different methods being applicable to different cases. But the time which this treatment takes is often a bar to its use. Thus Freud often consumes more than a year in completing an analysis. The instances given above took only a very short time, and may, I hope, encourage others to apply the same principle to similar cases.

THE fourteenth part of the Nyasaland Protectorate's Sleeping Sickness *Diary*, prepared by Dr. A. H. Barclay, Acting Principal Medical Officer of Nyasaland, states that further cases of the disease have been reported. Dr. Sanderson is now building a sleeping sickness camp at Ngani Hill, which is in the centre of the infected district, but is itself fly-free. All the cases are being sent to him, and are being treated with atoxyl. It is hoped to try salvarsan as soon as a suitable supply of the drug can be obtained. The most serious point to note at present, Dr. Barclay thinks, is the occurrence of 3 cases of sleeping sickness on the south-west arm of Lake Nyasa, a district which Dr. Barclay has for some time feared might be infected. He has also heard that the country between Kasankas and Dowa will also provide cases. As there is a continuous belt of fly from Kasankas down to Matope, and as fly appears to be encroaching steadily on new areas, it is feared that further cases may occur towards the south. Dr. Drummond finds that the removal of patients has caused the natives to become suspicious and to hide from him; another sleeping sickness camp is therefore to be built in South Nyasa in a district free from fly. Professor Newstead, of the Liverpool School of Tropical Medicine, has been appointed to the Protectorate for six months in order to study the different species of tsetse flies occurring in these areas.