1,261, and of these 662, more than half, were performed under spinal anaesthesia. The first cases were done in October, 1907, so that the method has ceased to be a novelty, and the increase has been a gradual one. Moreover, a large number of these operations were done by assistant surgeons or house-surgeons, who used the anaesthetic they preferred.

To sum up, it is clear that spinal anaesthesia, owing to its limited field, can never replace chloroform. If we are to be limited to one method that must, of course, be general anaesthesia. But, in the ground common to both, there is not a vast difference between the two methods in ease, safety, and effectiveness, if these are taken together over a large number of cases. What there is, is slightly in favour of general anaesthesia. Yet in a number of cases spinal anaesthesia offers advantages which cannot be got in any other way, and its apparatus and technique are so simple that it is worth any surgeon's while to use it whenever there is anything definite to be gained by it. In the large number of cases where there is nothing to choose between the methods, the use of stovaine becomes a matter of taste and convenience.

There seems to be a tendency in England to regard spinal anaesthesia as a dangerous method, and one that can only be employed successfully by a few experts who have devoted their attention to it. As far as my experience goes this is less true than it is of chloroform—the danger of spinal anaesthesia is about the same, and the skill required for its administration less.

In conclusion, I should like to express my especial obligation to Dr. Aly Bey Ibrahim, my assistant surgeon, and my thanks to the English and Egyptian residents, especially Dr. Thom and Dr. Maloney, to my successive house-surgeons, and to Dr. Shahin, recently appointed anaesthetist to the hospital, who have helped me by anaesthetizing cases, by keeping notes, and by making valuable suggestions.

REFERENCE.
1 *Lancet*, October 1st, 1910.

A CASE OF PAROXYSMAL TACHYCARDIA.

BY

R. O. MOON, M.D. OXON., F.R.C.P.,

PHYSICIAN TO THE WESTERN GENERAL DISPENSARY, MARYLEBONE,
AND TO THE HOSPITAL FOR DISEASES OF THE HEART, SOHO.

M. K., a tailoress, aged 35, came up to my out-patient department at the Heart Hospital in 1909 complaining of palpitation of the heart and pains in the chest for some years. On feeling the pulse I found it was beating at the rate of 180, but examination of the heart disclosed nothing abnormal beyond the rapidity of the pulse.

She was said to have been delicate from childhood, and began to be subject to these attacks of rapid heart beating and palpitation some fourteen years previously. There was no history of any other illness. The attacks used to come on about every five or six months, beginning quite suddenly and unexpectedly, very often soon after food, but also quite frequently in the night. She herself associated the attack with indigestion. The attack would last any time from four to twenty-four hours and then pass off gradually. As is generally noted in these cases of paroxysmal tachycardia, she was not very much distressed, and would go to her work as usual; she could not, however, lie down during an attack, and therefore when it occurred at night she was a good deal troubled by it. On only one occasion did she complain that she had severe pain in an attack.

I saw her from time to time, when her pulse was usually 76, but I never saw her with the tachycardia again as on the first occasion. She could never be persuaded to come into the hospital, but by treatment of the indigestion she became much better, the attacks very infrequent, and her condition generally improved. In August, 1910, however, she had two attacks lasting seven and ten hours respectively, after which there were no more. I saw her in January, April, and May, 1911; each time she complained of feeling generally weak, but there were no more attacks of tachycardia, and after May 15th I did not see her again. Apparently she had been fairly well till September 2nd, and was at work that day the same as usual, but about midnight she was seized with an attack. The friends with whom she lived were not alarmed, as it appeared to them no different from other attacks; but she soon complained of very severe pain, calling out with the pain, which she had never done before, and saying that she was dying. At 4 a.m. she became very quiet, and died at 4.30. As she had not been to the hospital for four months an inquest was held, and the post-

mortem examination, which was made by Dr. Freyberger, disclosed an unusually large patent foramen ovale, but an otherwise normal heart.

Clinically I had never found any evidence of an organic lesion of the heart, and I looked upon the case as one of simple paroxysmal tachycardia. Had we been able to take her into the hospital, some features pointing to the congenital lesion might have revealed themselves. I have not come across, in the literature on the subject, any other instance of paroxysmal tachycardia being associated with a congenital lesion, and consider that in this instance the association was a purely accidental one. Although the ultimate prognosis in paroxysmal tachycardia is unfavourable, yet I think anything in the nature of sudden death is distinctly rare; as a rule, the attacks become more frequent and last longer, the heart dilates, and we have all the signs and symptoms of cardiac insufficiency with backward pressure.

In my experience, most attacks of paroxysmal tachycardia terminate abruptly, the heart appearing to the patient to be jerked, so to speak, into the right place. However, in this case the patient was quite clear that the symptoms passed off gradually.

LEUCODERMIA UNDERGOING PIGMENTATION ON EXPOSURE TO SUNLIGHT.

BY

K. H. JONES, M.B., CH.B. VICT.,

STAFF SURGEON, R.N.

IN view of the interest recently exhibited in certain pathological conditions of the skin, concerned with the comparative absence or excess of pigment in its layers, notes of the following case may be worthy of record.

G. A. L., aged 18, entered the Royal Navy as a youth, in January, 1911, and came under my care for a scalp wound in March of the same year. A few days after his entry on the sick list I saw him stripped to the waist, and the fact that he was subject to leucoderma in a high degree was at once obvious. The condition affected the skin of the neck in front and behind, as well as that of both groins. The usual condition obtained in this case—there was a brownish pigmentation of the skin, and the patches devoid of pigment lay in the coloured portion. There was a yellowish ochreous coloration of the skin of the sides and the front of the neck above the clavicles, and above the vertebra prominens behind, and a much slighter pigmentation of that over both scapulae. On the left side of the neck were two large oval patches of leucoderma about 1½ in. long, and about ¾ in. wide, their long axes being approximately horizontal. In front of these patches and above them was a much smaller one, of about the same shape. On the right side of the neck were two small oval patches. Posteriorly a large butterfly-shaped patch had its head at the vertebra prominens, and extended over both scapulae. In the region of the groins pigmentation and leucodermic patches were less well marked than in the neck.

This youth was of a fair complexion, with blue eyes, and his skin was of the cream-coloured tint that tans a deep brown on exposure to the sun. He was sturdily built, enjoyed general excellent health, and had had the leucoderma ever since he could remember. There was not the slightest suspicion of syphilis in this case, but it is of interest that the distribution is that usually associated with syphilitic leucoderma.

In July I had occasion to look at this youth again, and at once noticed a change in all the leucodermic patches on the sides of the neck which were exposed to light. In all there appeared a reddish blush of inflammation, and in all, but especially in the two largest on the left side of the neck, there had developed numerous oval and rounded areas, 2 or 3 mm. in diameter, and of a brown colour; whilst a pigmentary invasion of the periphery of the patches was also seen to be taking place. At the back of the neck and in the groins, where no light reached the leucoderma, no change in them had taken place.

This youth had previously to his entry into the navy worn a high collar, and had been engaged in an indoor employment. Having to wear the characteristic open collar of a bluejacket, exposed to the light, and particularly the sunlight, portions of his skin which previously had been covered, and the pigmentary invasion of the leucodermic patches was probably the result.

It is stated by many authorities that the pigment increases in these cases during the summer, and particularly at the edges of the leucodermic patches. Stelwagon of Philadelphia, cites a case in which all pigment was

removed by the extension and coalescence of the leucodermic patches; and that thereafter the skin was unaffected by the sun. Crocker quotes Ehrmann as stating that in one case brown spots occurred in the leucodermic patches, as in the case above described, but does not ascribe any cause.

There is little doubt that with continued exposure to the sun the whole area in this case would become pigmented, the exact opposite of what usually obtains in such cases. From the reddish blush of the skin in this case there is little doubt that the increased pigmentation is inflammatory, probably due to the actinic rays of the sun, and very closely allied to ordinary sunburn.

Memoranda:

MEDICAL, SURGICAL, OBSTETRICAL.

ECTOPIC PREGNANCY TWICE IN THE SAME PATIENT IN SIX MONTHS, NECESSITATING TWO SEPARATE LAPAROTOMIES.

THE patient was sent to me by Dr. J. McLaren of Tottenham. She is 27 years of age, and has been married four years. Three years ago she was confined with a 7 months child, which lived three days. Six weeks after this confinement the menstrual discharge made its reappearance, and thereafter it recurred regularly every month until April, 1911. On May 18th, after an amenorrhoeic cycle of seven weeks, she was suddenly seized with severe pain in the lower abdomen, and at the same time a haemorrhagic discharge made its appearance vaginally.

From May 18th until June 12th, when I performed the first laparotomy, there had been continuously a haemorrhagic discharge from the vagina and there had been three separate attacks of pain. The physical signs then noted were the following: On palpating the abdomen there was felt in the hypogastrium a small swelling, but the abdomen generally was so tender that satisfactory palpation was impossible. The cervix, which was somewhat nearer the anterior vaginal wall than usual, was lower than normal. The body of the uterus—not enlarged—was felt in front bimanually. In Douglas's pouch was felt a boggy swelling, which extended into the left pelvis and appeared to be of about the size of a cocoanut.

At the operation the omentum with much small bowel was adherent to a swelling in the left and posterior segments of the pelvis. On separating these adhesions dark blood immediately escaped. The left tube, which contained a quantity of blood clot, was removed with the left ovary. An embryo of about six or seven weeks was found, and villi were discovered in the ampullar end of the tube.

Menstruation began as usual on August 30th, and lasted for the usual number of days (four). She expected to menstruate on September 27th, but she missed this period, and almost at the same time complained of feeling sick in the morning. On October 25th she complained of slight pain in the lower abdomen and almost at the same time she remarked that she had a haemorrhagic discharge. She continued to lose freely for five days, and altogether for ten days. During the first five days there was abdominal pain, but this was never severe. It was never at any time like the pain experienced on the previous occasion. She was sent to see me on November 15th, 1911, merely because she complained of tenderness of the abdomen, and although there had been no vaginal discharge since November 4th she had become apprehensive that something was wrong.

The physical signs noted on November 15th were the following: There was nothing detected on palpating the abdomen except slight tenderness in the right iliac region. The cervix uteri was centrally placed; it was not soft and the os was not patent. The body of the uterus was felt easily bimanually; it was not enlarged. Behind and to the right of the uterus was felt a small, tense, movable swelling, which was only very slightly tender. This swelling was of the size of a hen's egg.

At the second operation a large amount of small gut was adherent to the small swelling in the posterior and right segments of the pelvis. The haematoma burst immediately it was grasped. The right tube and ovary—the former with blood clot and the latter with a corpus luteum—were

ligatured and removed. The embryo on this occasion was about $\frac{3}{4}$ in. in length. The patient made a good convalescence, with no untoward symptom.

London, W.C.

JAMES OLIVER, M.D., F.R.S.Edin.

EPIDEMIC JAUNDICE.

THE cases mentioned in the BRITISH MEDICAL JOURNAL of November 18th, p. 1353, of catarrhal jaundice occurring in an epidemic form, are interesting to me, because it was only a few days ago I was called to see a boy 8 years old similarly affected. He complained of pain in the abdomen and slight vomiting, and there was also headache and malaise. His temperature was nearly 102°, his tongue was coated, and there were some very prominent bright red papillae visible, but his throat was all right. The conjunctivae were yellow and the urine discoloured. The skin was very hot, but not apparently tinged. There was pain and great tenderness in the epigastrium and hypochondria, chiefly the right. I do not think that the liver was enlarged, but he would scarcely allow me to examine him. There was not much tenderness below the umbilicus. He had had no rash, and his mother told me that the other three children (who were younger) had had their eyes jaundiced and also the urine, but that they had recovered.

He had been ill for some days, but was going on well until he went out, and then he rapidly got worse and the pain appeared in his abdomen, and that was the reason I was called to attend him. There was nothing that they had taken that could have caused it, and I said that it was a case of "epidemic jaundice." He was given a mixture containing strong doses of sodium bicarbonate, a little dilute hydrocyanic acid, and chloroform water, which was to be taken every four hours. He was to be kept on a fluid diet and in bed, and if the pain increased linseed meal poultices were to be applied, and I was to be informed if he got worse. I first saw him on Wednesday, November 8th, and I next visited him on Friday, November 10th.

I was told that a poultice had been applied on the afternoon of November 8th over the painful region, and that it had relieved him. He had now got up, the conjunctivae were clearing and also the urine, his temperature was normal and he felt well and he had practically recovered. He has continued to improve, and there has been no necessity to visit him.

The whole of the children in the house were affected. It is situated on the countryside and in a bleak position, and the weather was cold and wet. I am not aware of any other cases in the district, and I think it appropriate to communicate this because of the apparently comparative rarity of these epidemics.

Glandowey, Cardiganshire.

D. OWEN WILLIAMS, M.B.

PERIODICAL HAEMATEMESIS AND MENSTRUATION.

ON account of its rarity the following case may be of interest to readers of the BRITISH MEDICAL JOURNAL.

M. J., aged 29 years, has been under my care for over twelve months. Menstruation is quite regular and seems normal, but at each menstrual period she expectorates considerable quantities of blood. For twelve months both oral and uterine discharge has occurred regularly every twenty-eight days, and neither has occurred without the other during that time. The woman has always been allowed to go to bed in the Infirmary for a few days at each menstrual period, and the stages of the process have been repeatedly and carefully observed.

She becomes uncomfortable with a feeling of sickness, and this sensation is relieved when a mouthful of dark-coloured foul-smelling blood clot is brought up, by an effort akin to vomiting, without any straining. This occurs about ten to twelve times in the twenty-four hours, and continues about four days.

The woman seems healthy and expresses herself as quite well between the periods. Her history is not much to be relied on, but she has had one miscarriage.

One hears and reads of vicarious menstruation, but as the case I refer to does not agree with any one I have ever seen or heard of I should be glad to know if any similar case has been recorded.

DAVID FLECK, M.B., B.Ch.,

Medical Superintendent, Eastern Counties Reformatory,
East Harling, Norfolk.