a tendency to accumulate in the system. In my own experience veronal, of the newer drugs, while sometimes producing giddiness, unsteadiness of the gait, headache, and sickness, has been the most useful. In some cases of melancholia the blood pressure is high, and sleep will not be obtained till this symptom has been treated.

The third symptom needing treatment is the tendency to suicide, and this is of course done by means of continuous supervision exercised by skilled mental nurses over the actions of the patient. It is not necessary to add anything to what I have already said on this important subject. The nurses are also expected to occupy and interest the patient, and if by their attention and sympathy they win his confidence, they become therapeutic agencies of great value.

Three Special Methods.

There are three other modes of treatment to which I desire to refer briefly. The first is directed to a lowering of the blood pressure, which has often been found to be abnormally high. A warm bath lowers the blood pressure, and this is often followed by temporary relief of the symptoms. It also enables the patient to sleep if taken at bedtime. The most convenient way of reducing the blood pressure is by the administration of tetranitrate of erythrol. Half-grain tablets may be given twice daily, slowly increasing the dose till two or three grains daily are given. It is thought by some that the high blood pressure has an important bearing on the presence of feelings of depression, for in the converse state of maniacal exaltation the blood pressure is usually low. This form of treatment sometimes acts well, and is worth trying when other measures have failed. In one case of mine it was very successful where high blood pressure and depression were found to vary together in a remarkable way.

Another form of treatment also directed to remove the mental pain is the administration of opium. Some believe this feeling of pain is due to a sense of resistance and difficulty in the environment, the result of muscular sensations and impairment of movement. It is more probable that it has an organic basis, and is due to the general malnutrition of the body, the result of which is that the important organs transmit nervous impulses to the centres of a disagreeable kind. These impulses we are not conscious of, but it is believed that the massive effect of the whole forms the background of our consciousness, and produces our feelings of well-being or malaise as the case may be. The rational way of removing mental pain due to such a cause would be to remove the malnutrition by tonic treatment. This is attempted, but in addition it is believed that this feeling of depression can be alleviated by opium. In France it has been largely used for this purpose, and I have found it of marked service, especially in some cases of anxious melancholia. It is best given in the form of tinctura opii, starting with 5 minims three times a day, and rapidly increasing the dose till definite results are obtained. This form of treatment has many disadvantages, among which are two: It tends to increase the difficulties caused by constipation and sluggish action of the liver, and it tends to create a habit to which the patient may fall a victim. If administered, it should be done without the knowledge of the patient.

Lastly, melancholia tends sometimes to become very chronic, and in these cases, if the state of the nutrition permits it, and there is no organic disease to contraindicate its use, fairly large doses of thyroid may be given for one week. Three tabloids of 5 grains each should be given the first day, then 6, 9, 12, 9, 6, 3 daily in succession—an increase of three each day, till a maximum dose of 60 grains is reached on the fourth day, followed by a decrease of three each day. The patient must be kept strictly confined to bed and the pulse carefully watched. This drug undoubtedly acts as an excitant of the nervous system, but it is believed that its beneficial effects in melancholia are largely due to its influence on the general metabolism. There is a stirring up and a quickening of the trophic activities, and on the basis of this change a fresh start towards recovery may be made.

PROPHYLAXIS.

In conclusion I desire to say one word on prophylaxis. In all other departments of medicine the prevention of

disease is considered the highest aim, and we have now reached a stage in psychiatry in which our attention is being given to the prevention of mental disease. I say nothing at the present time on the interesting subject of eugenics, but I would urge on the family physician the necessity of dealing promptly and seriously with the earliest indications of disordered mental function. This is particularly necessary in the cases of those who have suffered from previous attacks, and in the members of families known to them to have a neuropathic history, when they are undergoing any severe strain. For example, a neurotic daughter will nurse an aged mother, with morbid conscientiousness and anxiety, losing her natural rest and depriving herself of relaxation and fresh air and exercise. When the shock of the inevitable end falls upon her, or even before then if she be a victim to influenza, she suffers a complete mental breakdown. It is the duty of the family physician to foresee the possibility of this, and to prevent it if that be possible. The early stages of mental disease are functional, and this is more applicable to melancholia than to any other form. Complete rest, a change of scene, agreeable companionship, and simple attentions to the health may then in a few days or a week or two be the means of averting a threatened attack. The mental as well as the physical health of these individuals and families should be in the mind of the family physician.

THE ETIOLOGY OF IRITIS.

BY

T. HARRISON BUTLER, M.D. OXON.,

HONORARY OPHTHALMIC SURGEON TO THE COVENTRY AND WARWICKSHIRE AND TO THE WARNEFORD, LEAMINGTON, AND SOUTH WARWICKSHIRE GENERAL HOSPITALS.

ON July 16th, 1910, an article appeared in the JOURNAL upon the etiology of iritis, which led to a very interesting correspondence. Mr. Kenneth Campbell stated that in his experience, taking all cases of iritis into consideration, 70 per cent. were due to syphilis, 20 per cent. to gonorrhoea, 9 per cent. to oro-alimentary, uterine, and tubal infection, and 1 per cent. to tuberculosis. He denies the existence of "rheumatic iritis." Römer, in his new textbook, embodying his lectures at the University of Greifswald, states that 25 per cent. of his cases are due to syphilis, a disease which is at least as common in Germany as in England. He devotes several pages to a consideration of rheumatic iritis, and carefully distinguishes it from gonococcal iritis. Worth, in a letter which appeared in the JOURNAL for August 13th, 1910, finds that only 30 per cent. of his private cases are due to syphilis, and ascribes more than one half to oral sepsis and kindred conditions. Beaumont, of Bath, believes that the rheumatic iritis is uncommon, and that many cases ascribed to rheumatism are really gonococcal. C. W. Jennings (*Ophthalmology*, 1909) found that among 500 cases of iritis 25.4 per cent. were rheumatic, and only 5.2 per cent. were gonococcal. Kenneth Campbell states that "of double iritis the cause is almost certainly syphilitic."

In view of this wide divergence of opinion it has seemed to me worth while to go through my casebooks, both private and hospital, and extract all the cases of iritis with a view to determining what proportion must be allowed to the different sources of infection, and whether any guide to diagnosis can be ascribed to the fact that one eye or both is affected.

Among 5,710 patients I found 100 suffering from iritis, a percentage of 1.75 only. I have included all cases of primary irido-cyclitis, but have excluded examples of interstitial keratitis complicated with iritis, and all traumatic cases, which are very common in a manufacturing town like Coventry. A septic foreign body in the cornea often induces a fairly severe iritis.

I shall first deal with the cases as a whole, and then compare my private with my hospital patients. I am fully aware that it is impossible to formulate any absolute law upon 100 cases, but it is possible to gather some valuable information, even from such a small number.

Exact diagnosis is exceedingly difficult, often quite impossible. In most cases one cannot determine the etiology

from the local signs, although occasionally the appearance of the iris is quite pathognomonic. I have examined the cases in the following way. The local examination has always been made in the dark room with the aid of a Zeiss loup, and with the ophthalmoscope. The patient has then been examined for rashes, enlarged glands, sore throat, and has been questioned closely to exclude syphilis. As iritis is chiefly a secondary symptom, when no other signs of infection can be detected, and no history can be obtained, it seems fair to regard the syphilitic etiology as improbable unless the local appearances are characteristic of that disease. Römer carefully distinguishes between an early syphilitic roseola of the iris, corresponding to the early skin roseola, an early papular form, a late papular form, and a gummatous iritis. The syphilitic changes are situated deep in the iris, and the adhesions to the anterior capsule are broad and are not confined to the pigment layer of the iris. But what one often sees in early secondary iritis is a ciliary blush, slight muddiness of the aqueous, little or no change in the appearance of the iris, with tiny thread-like adhesions between the capsule and the iris pigment layer. In addition, there are usually precipitates upon the anterior capsule, and less frequently in these very early cases upon the back of the cornea. When the pupil has been dilated with atropine—and it often dilates fully—a ring of pigment is seen on the anterior capsule. A ring in contradistinction to isolated islets of pigment is at least suggestive of syphilis.

If syphilis can be excluded and the patient is a man, one can inquire about recent or past gonorrhoea. It is quite unnecessary that the gonorrhoea be recent. I have seen gonococcal iritis in a man who had had the disease twenty years before, and had been free from it since that date. In gonococcal iritis there is a tendency for a gelatinous-looking exudate to accumulate upon the iris, and occasionally in the anterior chamber, but this condition is often absent.

Having excluded syphilis and gonorrhoea as far as possible, I then inquire for joint pains, and evidence of osteo-arthritis, the so-called "rheumatism." This condition is almost certainly an autotoxaemia, and I would prefer to class these cases as autotoxaemic rather than as rheumatic.

The teeth and nose should now be examined, for iritis, and especially an insidious cyclitis, are frequently due to toxins absorbed from pyorrhoea alveolaris, from sinus disease, and from other septic foci in the nose. Probably a septic otorrhoea may be responsible for some obscure forms of irido-cyclitis. Suppuration elsewhere in the body should not be neglected.

It is now well to go into the question of tuberculo-toxaemia thoroughly. When other causes have been excluded, and the patient has enlarged glands, or a history of such, a tuberculous family history, or other signs of this infection, the iritis may be due to the presence of tuberculo-toxins in the blood. Such examples should be tested by the subcutaneous injection of tuberculin, and treated by von Hippel's method. I have dealt fully with this subject in a paper in the *Ophthalmoscope* for December, 1910, and it need only be mentioned. Tuberculous iritis proper is rare. I have only seen one case among over 5,000 patients. If all these probable causes have been excluded, the probability of intestinal sepsis must be investigated, but it is hardly scientific to ascribe iritis to this cause unless indican, phenol, and an excess of sulphates are present in the urine. I am certain, however, that intestinal conditions can and do originate chronic cyclitis. I saw a case a year ago in which chronic cyclitis was associated with mucous colitis. When the colitis was actively treated the evidences of cyclitis, precipitates upon the back of the cornea, entirely disappeared.

The results of my investigation appear in the accompanying table. As there are 100 cases all the figures are percentages.

Cases of iritis formed 1.84 per cent. of 1,630 cases seen in private, 1.16 per cent. of 600 cases seen at the Warneford Hospital, and 1.77 per cent. of cases seen at the Coventry Hospital. The combined hospital percentage is 1.69, as against 1.84, the percentage among my private patients. Twenty-nine per cent. of my private patients had an iritis of unknown etiology. In half of these 8 cases only one eye was affected; in half, both. Five were females, three males. It is only to be expected that in hospital practice

Table showing the Etiology of Iritis in 100 Cases, both Private and Hospital.

Etiology.	Number.	Male.	Female.	Single.	Double.
Doubtful	41	16	25	27	14
Syphilis	22	9	13	10	12
Oral sepsis, nasal, etc.	12	7	5	7	5
Gonococcal	6	6	—	4	2
Rheumatic	6	5	1	5	1
Tuberculo-toxaemia.	5	2	3	3	2
Intestinal stasis ...	4	1	3	1	3
Albuminuric	2	—	2	2	—
Diabetic	1	1	—	1	—
Tuberculous	1	1	—	1	—
Gout	0	—	—	—	—
Total	100	48	52	61	39

Table comparing the Etiology of 31 Private Cases of Iritis with 69 Hospital Cases expressed in Percentages.

Etiology.	Hospital.	Private.
Doubtful	47.8	29
Syphilis	18.8	25.8
Oral, nasal, and general septic conditions	5.8	25.8
Gonococcal	7.2	3.2
Rheumatic... ..	5.8	6.4
Tuberculous toxaemia	5.8	3.2
Intestinal sepsis... ..	2.9	6.4
Albuminuric	2.9	—
Diabetic	1.45	—
Gout	—	—
Percentage total	99.45	99.8
Total	69 cases	31 cases.

the proportion of doubtful cases is higher. But in most of my cases the advice to see the dentist was written upon the prescription paper, and not in the notes, which until recently were kept in books. A comparison of the percentages of private and hospital cases due to oral sepsis shows that over four times as many private cases suffered from oral sepsis as hospital patients, which is contrary to reason. It is only recently that I have realized the immense importance of oral sepsis in the genesis of iritis, although I have long known its etiological value in cyclitis. I am of opinion, then, that a large proportion of the doubtful cases are really due to oral sepsis. Few of them can have had secondary syphilis, for I have invariably gone into this question very carefully.

The net result of the inquiry is that in at least 30 per cent. of all cases it is almost impossible to determine the real origin of the disease. Twenty-five per cent. are due to syphilis, and only about 5 or 6 per cent. can truly be termed rheumatic. An examination of the figures shows that in 22 syphilitic cases 10 were monolateral and 12 bilateral, so that this question cannot be considered in coming to a diagnosis. Cases due to oral sepsis and to intestinal stasis are more frequently bilateral than syphilitic cases. One patient put down as rheumatic iritis had had an acute attack of rheumatic fever only five weeks before the iritis came on. Two cases were associated with albuminuria. In one there was albuminuria, iritis, and a single retinal haemorrhage, but no retinitis. The other was a case of albuminuria and general anasarca. No case due to gout has ever been seen by me. The investigation lends no colour to Mr. Kenneth Campbell's view that 70 per cent. of all cases are due to syphilis. Even allowing that all my doubtful cases were syphilitic, which

is absurd, even then the figure only reaches 63 per cent. I believe that Römer's estimate, 25 per cent., is the correct one. It is interesting to note that in Palestine iritis is almost unknown, although pyorrhoea is very prevalent. I only remember seeing two or three cases among several thousand seen at the British Ophthalmic Hospital.

A FEW POINTS ON SOME COMMON AFFECTIONS OF THE EYES.*

By H. PERCY BENNETT, M.B., C.M.EDIN.,
HONORARY SURGEON, NEWCASTLE-ON-TYNE EYE INFIRMARY.

I PROPOSE to touch upon some affections of the eyes which every busy general practitioner is liable to meet fairly frequently, and will endeavour to give a few practical points that may be of some help.

We all know the difference between the injection of an eye in iritis and conjunctivitis. In iritis the injection is immediately around the cornea and is of a pinkish hue, whereas in conjunctivitis the hyperaemia is more marked towards the periphery of the bulbar conjunctiva. But it is not very uncommon to come across cases of conjunctivitis where there is, from the appearance of the eye, some doubt as to iritis being also present, and I am sure you will all have seen cases like the following:—

A patient with one or both eyes markedly red, with no diminution of the injection towards the margin of the cornea. We often find that the eye has been treated at home by keeping it tied up, or by poulticing it—perhaps with tea leaves. Such treatment, besides increasing the vascularity of the conjunctiva, may cause partial maceration of the corneal epithelium, and this, if one is not on one's guard, may be mistaken for turbidity of the aqueous, which is another sign of iritis. To proceed with the diagnosis, we next test the reaction of the pupil to light, and, if this is not satisfactory, inquire as to pain; stabbing pain in the eye, waking the patient about 2 or 3 a.m., is very characteristic of most cases of iritis. If still in doubt whether there is or is not some iritis present one next puts in a drop of $\frac{1}{2}$ per cent. atropin (provided glaucoma can be excluded), and if the pupil dilates well in half an hour, with no iris pigment or exudate on the anterior capsule of the lens, we may then conclude that there is no iritis. Giving such a patient atropin to use regularly at home, in addition to the correct treatment for the conjunctivitis, may in many retard the cure, and is only permissible when, after a thorough examination, there is still some doubt as to threatened iritis; in such a case it is better to order weak atropin drops ($\frac{1}{2}$ to 1 grain in the ounce) and see the patient in a day or two, when the eye may be found free from any suspicion of iritis, and the drops can then be discontinued.

The mistake usually made in treating conjunctivitis is in using strong astringents or strong antiseptics at the commencement; the condition is thus usually aggravated.

Boric acid lotion, to wash out the conjunctival sac thoroughly, or even only sterilized water with some sodium chloride in it, will in many cases bring about the cure. In other cases astringents may later on be necessary, and as there are many of these to choose from, it is better to avoid the *lead lotions*, because, if there be an ulcer on the cornea, a white and disfiguring incrustation of lead forms at the site of the ulcer. This could only be removed afterwards with great difficulty. I have seen this occur on more than one occasion after a patient has had a lead and opium compress on the eye for a short time to relieve pain. These ulcers proved most obstinate to get to heal, required scraping several times, and were very painful. Attention to the patient's health and surroundings and getting him into the open air are extremely important, and may effect a cure without any applications to the eyes.

Unless advised not to do so, patients think they should stay indoors when they have what they call "cold in the eyes," and that they should even keep in a darkened room. There is no advantage in keeping a patient suffering from conjunctivitis in a darkened room. Dr. Berry points out that though the actinic rays

are known to cause irritation of the conjunctiva, which may indeed, as in snow blindness and electric light ophthalmia, be so great as to lead to violent inflammation, on the other hand they have a powerful bactericidal effect. The light of an ordinary room when the sun's rays are not directly passing into it has no had effect on conjunctivitis.

There is one form of inflammation of the lids, *angular conjunctivitis* (*Morax diplobacillus*), in which zinc salts are the most effective treatment; here the inflammation is of a subacute type, with the redness most marked at the angles of the lids.

Poulticing, or what comes to the same thing, tying an eye up (as the tears and discharge from the eye keep the bandage moist), is usually harmful; it increases the vascularity, and favours the growth of germs which are always present in the conjunctival sac. Nature by the act of blinking does much in the way of cleaning the sac.

If one eye alone be affected by conjunctivitis, be on your guard against the *tear sac* being the source of the trouble. In chronic blenorrhoea of the sac without marked distension the condition may easily be overlooked. Here it is the contents of the tear sac regurgitating through the puncta that keep up the irritation of the conjunctiva. Firm pressure over the sac with one finger will force some purulent matter through the canaliculi, and show the cause of the conjunctivitis.

Many early cases of inflammation of the tear sac, especially in young infants, can be cured by instructing the patients or the parents to empty the sac by firm pressure applied to it by the finger as frequently, it may be, as every half-hour. If the secretions are allowed to remain in the sac they act as an irritant to the mucous membrane, which becomes still more swollen, and so any stricture at the junction of the sac with the nasal duct is increased. Inflammation of the tear sac besides causing troublesome epiphora is a constant source of danger owing to the risk of infecting any corneal wound; a slight scratch of the cornea may in this way lead to a hypopyon ulcer, and perhaps to the ultimate loss of the eye. Similar danger to the cornea must be guarded against in ophthalmia neonatorum and in gonorrhoeal ophthalmia of adults; rough handling in examining such an eye may easily cause an abrasion of the cornea, and so allow its inoculation by the conjunctival discharge, which may result in a perforating ulcer and total loss of the eye. It is advisable, therefore, before examining such eyes to first instil a drop of cocaine inside the lids.

Crédé's method is an excellent prophylactic. I find the number of cases of ophthalmia neonatorum seen at the Eye Infirmary within the last few years has markedly decreased.

You may be called to see a patient complaining of pain in one eye, and on looking at it you find no conjunctivitis to speak of, but distinct circumcorneal injection. Assuming that the tension of the eye is not raised, and that glaucoma can therefore be excluded, you have to deal with either an affection of the iris (or deeper part of the eye) or of the cornea. It is not necessarily the iris that is affected; it may be a corneal ulcer, only to be discovered on minute examination. Many, I am afraid, at once jump to the conclusion that it must be iritis, and the patient thereby may be placed at a great disadvantage. A small corneal ulcer without any infiltrated edge may have to be searched for very carefully to be detected; the same remark applies to herpes of the cornea, or to the presence of a minute foreign body on the cornea, or to a slight abrasion of the cornea.

Here I wish to draw attention to the great advantage of using fluorescein (resorcin-phthalein anhydride) as a means of diagnosis in corneal affections. It may be used as drops (fluorescein 2 and sodium bicarbonate 3 in 100); one placed in the conjunctival sac, and then the excess washed out with, say, boric lotion or sterilized water, will leave a brilliant green stain on any part of the cornea that has been denuded of epithelium, thus mapping out beautifully any corneal ulcer. A minute foreign body on the cornea would be more readily detected after using fluorescein, owing to the green ring which would appear round it, and it would therefore be much more easily removed. Fluorescein can be obtained in small tablets to place inside the lids, a more convenient form for carrying about.

* Read before the Newcastle-on-Tyne Division of the British Medical Association.