procuring me the "hypress" blower, and for many valuable hints; to Mr. Nesbitt, of the Thompson Yates Laboratories, for making my instrument; and to my colleagues, Messrs. Thelwall Thomas, Bickersteth, Litler Jones, F. Jeans, and H. Woolfenden, for allowing me to anaesthetize some of their patients.

ACUTE EPIPHYSITIS.

BY

CHARLES M. KENNEDY, F.R.C.S.ENG.,

SURGICAL REGISTRAR, LONDON HOSPITAL.

THE cases of which I propose to speak are not strictly cases of acute epiphysitis, but cases of acute juxta-epiphyseal diaphysitis—that is to say, cases of acute osteomyelitis due to pyogenic organisms, and commencing in the soft vascular line where new bone is replacing cartilage, a line which is anatomically part of the diaphysis.

I have chosen this title because, in the first place, this group of cases forms a definite clinical entity quite distinct from the other groups of osteomyelitis, with which I do not intend to deal, and secondly, because acute juxta-epiphyseal diaphysitis, the only accurate name I know for the disease, is too much of a mouthful for everyday use. The term "osteomyelitis" is too vague, for, though it includes this group, it also includes many other conditions.

Acute pyogenic inflammation of the epiphysis itself is so rare that it may be neglected. In the ten years 1901 to 1910 (inclusive) there is, judging from the notes, only one case that might have started in the epiphysis (and this case is doubtful) out of 309 cases of acute epiphysitis. The term, then, though unscientific, is convenient, and in using it I am following the late Mr. Barnard,¹ who in 1903 read a paper under the title "The Relation of Trauma to Acute Epiphysitis."

CAUSATION.

The exciting cause is a pyogenic organism. The organism is usually *Staphylococcus aureus*. Trendel,² when dealing with osteomyelitis in general, found staphylococcus, streptococcus, and pneumococcus. In this series we find corresponding results, as follows:

Bacteriological Results in 49 Recent Cases.

	Cases.
Staphylococcus was found in	38
Type not stated " " " " " "	5
<i>Staphylococcus aureus</i> " " " " " "	28
<i>Staphylococcus citreus</i> " " " " " "	2
<i>Staphylococcus albus</i> " " " " " "	3
Streptococcus " " " " " "	7
Mixed streptococcus and staphylococcus was found in " " " " " "	2
Pneumococcus was found in " " " " " "	2

That is, 77 per cent. staphylococcal.

The organisms are carried to the bone by the blood stream. The route by which they reach the blood stream is somewhat debated. Thompson and Miles³ state that they may get in through septic sores, but that probably they are more often due to sore throats and other infections of the mucous membranes. Dolbey⁴ regards sore throats as the commonest source of entry of the organism.

I found a note of a possible source of infection in only 44 cases, but these 44 amounted to 79 per cent. of the cases where any note as to the presence or absence of a source of infection was made. In these cases a septic condition of the skin was by far the most common; there were 36 cases of septic foci in the skin, most of them septic sores, 1 a septic burn; 1 case, in an infant of 2 weeks old, was apparently due to umbilical sepsis; 1 was a boil on the neck, and another multiple superficial abscesses. The other foci found were 2 cases of whitlow, 3 of otorrhoea, and 6 of tonsillitis; but 3 of these latter had also skin sepsis.

It would appear, therefore, that the organisms usually reach the blood by the skin—a conclusion borne out by the predominance of staphylococci as the causative organisms, for if the tonsils were the common site of entrance streptococcal cases should predominate, for they are the organisms usually found in cases of septic tonsillitis. Again, if the intestinal mucosa be the site of

entry, why is the *B. coli communis* so rarely the causative organism? Moreover, acute epiphysitis is very much more common in the poorer classes than in the well-to-do. This is usually explained by saying that the better classes are better nourished; but the disease usually attacks healthy-looking, well-nourished children. The children of the better classes play, and are probably nearly as much exposed to injury as the children of the poor; so that the trauma will not, I think, explain this greater frequency in the poor. Septic sores are, however, very much rarer in the well-to-do classes, and, moreover, if children of this class have a more serious septic skin lesion they are so nursed and coddled that they run little risk of trauma, the great predisposing cause. This greater frequency of acute epiphysitis in the poor, then, may be regarded as confirmatory evidence that the source of infection is a septic skin lesion in the great majority of cases.

The first theory that occurs to explain why the organisms in the blood should settle in the juxta-epiphyseal region of a bone and set up acute inflammation there, is that the disease is due to a specific organism which has a predilection for newly formed bone. This view Dolbey⁴ adopted in 1906. It is, however, negated by the fact that several varieties of organism are found, varying presumably with the source of infection. It may be objected that in the cases where organisms other than *Staphylococcus aureus* are found the disease is atypical, and does not truly belong to this group. This is, however, not the case; for in all my streptococcal and pneumococcal cases the notes would lead one to suppose that the cases were quite typical and genuine. There remains one other possibility—that the staphylococci and streptococci are secondary infections, the real causative organism having so far eluded us.

If we examine the known etiological facts, we find as follows:

Sex.

The disease is more common in males than in females. Thompson and Miles³ give the proportion as 3 to 1. In the series under consideration there were 217 males and 92 females, or 7 to 3.

Age.

Trendel² and Moore⁵ state that more than half the cases occur between 13 and 17 years. In the cases here considered, 70 per cent. occur between 7 and 17. The age incidence is quite different in the two sexes, a point I shall have to speak of again in relation to trauma. It is often stated that in young infants the disease occurs chiefly in congenital syphilitics. I have, however, found little evidence that this is so, and take the view that congenital syphilis is a negligible factor in the etiology of this condition. Albee⁶ states that the scrofulous and rheumatic dispositions predispose. I have already stated that this is not true of the scrofulous, as the disease usually appears in healthy subjects, and here I am in agreement with Dolbey.⁴ With regard to the rheumatic disposition, I find that in only 3 cases out of 309 is a previous history of rheumatism given.

Trauma.

Trauma has long been recognized as the chief predisposing cause in this condition. I find that although only 30 per cent. of cases gave a history of trauma, if only those cases were taken in which a definite note was made as to trauma, 61 per cent. gave a history of trauma. Trauma of some slight degree is so common in children that the number of cases giving a history of trauma is largely a matter of chance. Nevertheless, that trauma is a very important predisposing cause is shown by the sex and age incidence of the disease.

If trauma be the cause of the greater frequency of the disease in males, as is generally stated, the difference in sex incidence should be nil up to 3 years old, up to which age both sexes are equally liable to trauma, and should become more marked as the cases grow older. This is so; for in ten years at this hospital there were 53 cases under 3 years of age, and of these 27 were males and 26 females. Again, boys are most liable to trauma between 7 and 17, and 70 per cent. of cases in them occurred between these ages. Whereas in girls, 69 per cent. of cases occurred before 10 years of age. So that in both sexes the disease is most common when trauma is most common.

Again, if we take the incidence of the disease in limbs we get presumptive evidence of trauma; for in the first few years of life, when the child is practically a quadruped, crawling and falling about on arms and legs equally, the upper limb is affected twenty-five and the lower twenty-eight times, whereas in the later years the lower limbs provide the great majority of cases. The tibia and femur alone giving 77 per cent. of the cases between twelve and seventeen.

Admitting, then, that trauma is an important etiological factor, how does it act?

Ollier⁷ pointed out that the injuries to the epiphysis may be divided into para-epiphyseal strains, where the epiphysis is partially separated, and para-epiphyseal sprains, where the epiphysis is completely separated, but not displaced, and the well-known condition of separated epiphysis, where there is also displacement. As he pointed out, any trauma (for example, a twist) is most likely to affect chiefly the weakest part of the limb—namely, the soft, newly-formed bone. The epiphysis being injured there is a little effusion of blood, a little inflammatory reaction with stasis of the blood stream, and any organisms circulating in the blood and reaching such a spot will have time to grow and multiply and set up an acute inflammation.

The way in which trauma acts was very lucidly explained by Mr. Barnard¹ in his paper in 1903. He pointed out the gradations of the condition, from the simple para-epiphyseal sprain or strain, with its painful spot, but no deformity, and its aching for a few days, but no infection and no pyrexia, to the similar condition with mild infection (or high resistance on the patient's part) with its symptoms slightly more marked, with pyrexia for a few days, and lastly the typical acute epiphysitis.

The first group of cases we all see, but as a rule they are not recognized unless there is some slight displacement, as in Schaller's disease. The second group we also often see. The cases which come in as "epiphysitis," which are watched, and which recover without operative interference, and are then diagnosed as non-articular rheumatism, nearly all, I believe, belong to this group. That this is so is strongly suggested by cases such as the following:

A boy, aged 8, knocked his ankle in July, and had slight pain in it, which made him limp. He was seen at two other hospitals, and the ankle was painted with iodine. In December the ankle began to be painful and swollen, and he came to the London Hospital, and was admitted under Mr. Milne, who opened the lower end of his tibia, and found a localized abscess. The pus from this abscess yielded staphylococci on cultivation.

Nor is this case an isolated example. I have in the course of looking up the notes of acute cases accidentally come across the notes of two cases almost identical with that just quoted.

Trauma, then, will explain why acute epiphysitis always attacks the juxta-epiphyseal regions of bones without any necessity for a specific organism.

It would seem, however, that all cases of acute epiphysitis cannot be put down to trauma. If trauma be such an important causal agent, why do we never see a case of acute epiphysitis following on a separated epiphysis? There were no cases in my 309, so far as I could tell from the notes. Mr. Barnard met this objection by pointing out that acute epiphysitis does, though very rarely, occur in separated epiphyses. Now we see all the cases of separated epiphysis, but only a few of the cases of sprains and strains. In other words, acute epiphysitis is a complication of separated epiphysis, as of any other trauma, but separated epiphysis is so much less common than minor degrees of trauma, that, as one would expect, acute epiphysitis is much less common in the former group.

That this explanation is entirely satisfactory is very doubtful; for if trauma were the all-important cause, then the common sites of acute epiphysitis and of separated epiphysis should coincide. They do not, for it is found that acute epiphysitis is most common in the upper end of the tibia and lower end of the femur; whereas separated epiphysis is most common in the upper end of the femur. Acute epiphysitis is more common in the upper end of the humerus, whereas the lower is that most often separated. In the radius the lower epiphysis is both most often separated and most often inflamed; but separated epiphysis is rare in the radius as compared with the dislocated

elbow, which for anatomical reasons takes the place of the separated epiphysis of radius and ulna. Moreover, if cases of over 16 be taken, an age when one would expect all the patients to remember any trauma severe enough to injure an epiphysis, the percentage of history of injury rises, but only up to 65 per cent.

Both Ollier and Barnard admitted that all cases are not due to trauma. Ollier suggested "chill," and Mr. Barnard lowered resistance in the patient, as the other causal factor. Albee⁶ states that:

Spontaneous inflammations of the bones occupy seats of election, corresponding to the points of greatest activity of physiological development. The morbid disposition is, all other things being equal, in direct ratio to the activity of proliferation of anatomical elements.

If we take the lower limb we find the disease most common at the upper epiphysis of the tibia, and the lower of the femur, the most active epiphyses. If we take the upper limb we find the disease most common in the upper epiphysis of the humerus, again the most active epiphysis.

Thompson and Miles attribute the incidence of acute inflammation at the ends of bones, in part at least, to the arrangement of the vessels. The arteries run in the long axis of the bone and end in capillary loops just under the epiphyseal cartilage. It is, however, hard to see why micro-organisms should stick in these loops, and, moreover, this does not account for the disease being limited to the growing age, for the arteries and capillaries retain very much the same arrangement in adult life.

In growing bone vessels spring from these loops and grow vertically upward for a space, then join to form fresh loops. These vessels must at some period form blind alleys into which organisms may easily be driven and entrapped. This is, I believe, the reason why acute inflammation always attacks the juxta-epiphyseal line. The bigger the epiphysis and the more active the growth the more of these bacterial traps there will be and the greater the liability to acute epiphysitis, which explains largely the bone incidence. Now organisms would in this way be entrapped only for a short time and when the loop is completed be driven on again. Unless the organisms be numerous or virulent the patient escapes epiphysitis; but if trauma occurs, with perhaps some effusion of blood and stasis of the blood stream, even a single organism, perhaps, placed at such an advantage may grow and multiply and cause a fulminating disease. It is noticeable that in the bones more rarely affected there is almost always a history of trauma. The disease, then, may be due to any pyogenic organism (generally the staphylococcus, because the site of entry is usually the skin) which settles in the juxta-epiphyseal line for purely mechanical reasons dependent on the anatomy of growing bone. Trauma is an important predisposing cause, but it is by no means essential.

DIAGNOSIS.

There is no need to go fully into the clinical aspect of these cases, every one is so familiar with them. The sudden onset of pain near one joint, the bright eyes and flushed face, the rigors (or, often in babies, fits), the delirium at night, and occasional vomiting, are all familiar features.

The difficulties in diagnosis depend upon the condition of the patient when seen. If there is pus under the periosteum in any large amount, there will be a red, tender, fluctuating swelling. Here the condition has to be distinguished from cellulitis and abscess. If the patient is very ill, with a history of delirium and rigors, or if there is fluid (not necessarily pus) in one of the neighbouring joints, the diagnosis of acute epiphysitis is fairly safe. Very often, however, it is impossible to make a certain diagnosis until the abscess is opened, when the introduction of the finger and the feeling of bare bone immediately settles the diagnosis.

Whether this superficial swelling, caused by pus under the periosteum, will appear early or late depends on whether the initial lesion is peripheral or central. Mr. Barnard aptly compared the site of the initial lesion to the positions on a target. If the initial focus be in the position known as "an outer," the periosteum will be early and chiefly affected; if a "bull's-eye," the early spread will be down the Howship's vertical lacunae, and the medulla will be early and chiefly affected; if a

"magpie," the spread will be about equally distributed between medulla and periosteum. The compact bone is of course affected in all these positions, but owing to its greater density spread does not occur so rapidly through it.

Even when there is no pus under the periosteum, only increased vascularity and thickening, there may be oedema and redness over the affected part. The fluid in the neighbouring joints is often clear, and proves sterile on cultivation.

Even when the joint is full of pus, as it often is, this is usually a secondary infection of a clear effusion of this kind. Sometimes it is due to direct spread, but this is rare, because the thick, comparatively avascular epiphysis forms a formidable barrier to even the most virulent organism; spread under the periosteum, and so under the ligaments into the joint, though more common, is rare, because the periosteum of the epiphysis is very firmly bound down. Direct spread through the epiphysis is only noted twice (apart from the hip) in this series. In the hip-joint, however, the epiphyseal line being within the joint, the pus usually bursts early into the joint. This epiphysis, too, is one often separated by the morbid changes in acute epiphysitis.

If pus be found in the joint of a child, unless there is some obvious mode of infection, it is nearly always due to acute epiphysitis.

We now come to the more difficult cases to diagnose, those in which there is no obvious inflammatory swelling. They are usually mistaken for acute rheumatism. As a rule, the history will put one on the right track. The pain usually started in one joint, and on close inquiry not in but near one joint, and has remained there since. There is often a history of delirium at night, and perhaps of rigors. Where there is a history of delirium and rigors be very loth to diagnose acute rheumatism. Where the pain is always in one joint, and one joint only, be very careful in the search for evidence of acute epiphysitis. Search for a tender spot in the bone near the epiphysis. Sometimes a little discoloured patch, like a bruise, or occasionally like a patch of erythema nodosum, will point to the tender spot.

An early case of acute epiphysitis, especially if in a deep-seated bone, may be very difficult to diagnose; but this difficulty becomes tenfold when (as happened in several cases of this series) the disease has started with pains in two or three or even many joints before finally settling in one.

To meet these cases of great difficulty what assistance have we? A leucocyte count may help us, for in acute epiphysitis it is usually high (20,000 to 30,000 Albee);⁶ but in acute rheumatism there is also leucocytosis (the average being 16,000, Cabot).⁸ A blood count will often fail us, for in the very acutest cases leucocytosis may be absent, and moreover in acute rheumatism, especially if sodium salicylate has been given, it may be high enough to be confused with that found in the typical acute epiphysitis. The leucocytosis, then, is not to be trusted.

Sodium salicylate usually brings the temperature down, if only for a few hours, in acute epiphysitis, and so may lead to a false sense of security, and the waste of an invaluable day.

Radiography in most cases does not help. In this series where a note of a radiograph was made nothing abnormal could be seen. Ironside Bruce⁹ states that "in early stages of osteomyelitis no abnormality of outline or structure can be made out." In mild cases where the focus is early localized, a radiograph may show this focus. Of course, in chronic cases, and in cases which have been progressing for some time, radiography is invaluable.

Rheumatism is not the only thing for which acute epiphysitis may be mistaken. When house-surgeon I insisted on the full duty house-physician taking in a case as "scurvy rickets" which I afterwards had the mortification of taking over, when it was transferred to Mr. Openshaw's care, with several ounces of pus in the medulla and under the periosteum of its radius. When a dresser I well remember a case of Mr. Openshaw of acute epiphysitis of the iliac crest which closely simulated an appendix abscess. He was a boy of 13, taken suddenly ill with pain in the right iliac fossa and admitted with a lump there and a distended abdomen; but he had a history of a

fall. He was delirious and he had not been sick (though these cases often are), and he was correctly diagnosed.

Amongst the notes I find a case admitted moribund, which for three weeks had been treated as "enteric fever." It was to this type of case with its high fever and low muttering delirium, that Chaissagnac gave the name of "typhus of the bone."

Occasionally the diagnosis of the position of the lesion is very difficult; this is especially the case when the acetabulum is attacked. A careful history and close examination are the only trustworthy guides to diagnosis.

PROGNOSIS.

The mortality is high—higher than that of enteric fever or pneumonia; much higher than that of acute appendicitis. It was in the ten years 1901 to 1910, 34.6 per cent.

The majority of the cases die from pyaemia, with such lesions as purulent pericarditis, empyema, abscesses in lungs, myocardium, and kidneys, and not infrequently acute endocarditis. Abscesses may occur anywhere, but especially tend to attack other bones. This, as Mr. Barnard pointed out, is probably due to the organisms developing a taste for the "soil" on which they have been growing, or perhaps to the mechanical conditions found in growing bone, to which attention has already been called. Comparatively few cases die of pure toxæmia without *post-mortem* evidence of pyaemic spread.

Secondary hæmorrhage accounts for a few deaths. Two out of five cases of severe secondary hæmorrhage met with in this series died, in spite of prompt amputation. In most cases the secondary hæmorrhage is from the popliteal or a large branch in acute epiphysitis of the lower end of the femur.

Even if the patient escape with his life the outlook is not very bright. The average stay in the hospital is twelve weeks, and few cases are completely healed when discharged. They generally have a discharging sinus for months—at least 49 of the cases of this series were readmitted for sequestrotomy in periods varying from a few months to six years. A case was admitted under Mr. Openshaw last summer with chronic osteomyelitis of the femur, which dated back to an acute epiphysitis thirty-one years ago, and which had been giving trouble on and off since, in spite of many operations in various hospitals. Of nine cases which I have been able to see or hear of, at periods varying from one and a half to nine years after, only two were free from symptoms; but this is misleading, as in several cases the symptoms were the cause of my seeing them. The case of longest duration I heard of accidentally from Westminster Hospital nine years after, and he still had a sinus. This long-continued suppuration, with its risk of lardaceous disease, is not all, for many cases after a long interval of apparent cure suddenly develop signs of Brodie's abscess. I have come across the notes of cases fifteen, twenty-three, and twenty-six years after the primary onset.

In giving a prognosis, the extent and duration of the disease and the general condition of the patient are the chief guides. The age also helps:

The mortality before 1 year is 70 per cent.
The mortality from 1 year to 6 years is 41 per cent.
The mortality from 6 years to 11 years is 30 per cent.
The mortality from 11 years to 20 years is 29 per cent.

The mortality varies also with the organism, being only 14 per cent. in streptococcal cases.

The presence of albuminuria—a condition met with in 13 of the 55 cases (23 per cent.) in which I noted the condition of the urine—is of grave significance. More than half the cases in which it was found died; only one-fifth of these cases in which it was not found died. This albuminuria is apparently due to a toxic nephritis rather than to pyaemic abscesses in the kidney.

TREATMENT.

"Many lives and limbs have been lost because the surgeon was content with a diagnosis of periostitis and an incision through the periosteum," so says J. E. Moore, an American surgeon, and it is very true.

In every case where the pus is found under the periosteum in a child, unless the cause of that periostitis is very definitely known, the compact bone should be chiselled away and the medulla explored. If the medulla is not obviously diseased, the surgeon may be content

TABLES SHOWING AGE, SEX, AND BONE INCIDENCE IN 298 CASES OF ACUTE EPIPHYSITIS.

Females.

Age:	0-1.	1-2.	2-3.	3-4.	4-5.	5-6.	6-7.	7-8.	8-9.	9-10.	10-11.	11-12.	12-14.	14-16.	16-18.	18-20.	Total.	Died.
Tibiae: Total	2	—	4	4	—	—	2	3	5	1	2	4	1	1	2	—	31	6
" Upper end	1	—	1	3	—	—	—	3	2	—	—	2	—	—	2	—	14	4
" Lower end	1	—	2	1	—	—	2	—	1	1	1	1	1	1	—	—	12	1
Fibula: Upper end	—	—	—	—	—	1	—	—	—	—	—	—	—	—	—	—	1	0
" Lower end	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Femur: Total	3	3	1	3	2	1	—	—	2	—	—	2	3	3	1	—	24	10
" Upper end	—	—	—	—	1	1	—	—	—	—	—	—	—	1	—	—	3	3
" Head	—	—	—	—	1	—	—	—	—	—	—	—	—	—	—	—	1	1
" Great trochanter	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
" Lower end	2	3	1	3	1	—	—	—	1	—	—	2	2	2	1	—	18	7
Ilium	—	—	—	1	—	—	1	—	—	—	—	—	—	—	1	—	3	1
Os calcis	1	—	—	—	—	—	—	—	—	—	1	—	1	—	—	—	3	1
Astragalus	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Metatarsal	—	—	—	—	—	1	—	—	—	—	—	—	—	—	—	—	1	0
Phalanx	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Clavicle	1	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	1	1
Scapula	—	—	—	—	—	—	—	—	—	—	—	—	1	—	—	—	1	0
Humerus: Upper end	3	—	2	—	—	—	—	—	1	—	—	—	3	1	—	—	10	4
" Lower end	1	1	1	—	—	—	—	1	—	—	—	—	—	—	—	—	4	1
Radius: Upper end	1	—	—	—	—	—	—	1	—	—	—	—	—	—	1	—	3	2
" Lower end	1	1	—	—	—	—	—	1	—	—	—	—	—	—	—	1	4	1
Ulna: Upper end	—	—	—	1	—	—	—	—	—	—	—	—	—	—	—	—	1	0
" Lower end	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Total	13	5	8	9	2	3	3	6	8	1	3	6	8	6	5	1	87	27
Deaths	8	2	1	5	1	1	0	2	2	0	0	0	2	2	1	0		

Males.

Age:	0-1.	1-2.	2-3.	3-4.	4-5.	5-6.	6-7.	7-8.	8-9.	9-10.	10-11.	11-12.	12-14.	14-16.	16-18.	18-20.	Total.	Died.
Tibiae: Total	3	5	3	2	1	8	3	3	2	7	6	3	20	21	10	1	98	43
" Upper end	1	1	2	1	—	5	2	2	2	4	3	2	11	15	7	1	59	29
" Lower end	1	3	1	—	1	4	1	1	—	2	2	—	8	3	3	—	29	11
Fibula: Upper end	—	—	—	—	—	—	—	—	1	—	—	—	—	—	1	—	2	0
" Lower end	—	—	—	1	—	—	—	—	—	—	—	1	—	—	2	—	4	0
Femur: Total	1	1	—	4	1	3	1	5	5	4	1	3	8	9	6	3	53	12
" Upper end	1	—	—	1	—	—	—	3	3	1	—	1	1	4	3	1	19	5
" Head	1	—	—	—	—	—	—	—	1	—	—	—	—	—	—	—	2	2
" Great trochanter	—	—	—	—	—	—	—	—	—	—	—	—	1	2	2	—	5	0
" Lower end	—	1	—	2	1	1	1	2	2	3	1	2	6	5	3	2	32	6
Ilium	—	—	—	—	—	—	—	1	—	1	1	2	1	2	3	0	11	5
Os calcis	—	—	1	—	—	—	—	—	—	1	—	1	1	1	—	—	5	4
Astragalus	—	—	—	—	—	—	—	—	—	—	1	1	—	—	—	—	2	1
Metatarsal	—	—	—	—	—	—	—	—	—	—	—	—	1	1	—	—	2	1
Phalanx	—	—	—	—	—	—	—	—	—	—	—	—	1	—	—	—	1	1
Clavicle	—	1	—	—	—	—	—	1	2	—	—	1	1	—	—	—	6	3
Scapula	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Humerus: Upper end	—	2	3	—	—	—	—	—	—	—	—	—	—	3	1	2	11	6
" Lower end	—	1	1	1	—	—	—	—	—	—	—	—	2	—	—	—	5	1
Radius: Upper end	—	1	—	—	—	—	—	—	—	—	—	—	1	—	—	—	2	0
" Lower end	—	1	1	1	—	—	—	—	—	—	—	—	1	1	1	—	6	0
Ulna: Upper end	1	—	—	—	—	—	—	—	1	—	—	—	—	—	—	—	2	1
" Lower end	1	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	1	1
Total	6	12	9	9	2	11	4	10	11	12	9	12	37	38	24	6	211	79
Deaths	5	4	5	3	0	7	2	3	5	3	5	1	13	12	8	3		

N.B.—The figures in the bottom line appear at first sight to be incorrect, because several cases come under a general as well as a particular heading; care has been taken to count these figures only once.

with removing a large window in the compact bone, and this window should be as wide as the bone in question, but the medulla should not be gouged out. If the medulla is scraped out the endosteal membrane is removed, and the compact bone, already stripped of its periosteal blood supply, is more likely to die and form a sequestrum (Nichols). If the medulla is diseased, then the compact bone must be removed as far along the shaft as the disease has spread and a little further, and the diseased medulla gouged out. Where the medulla is gouged no overhanging ledges of compact bone must be left: they impede dressing, and usually form sequestra. In the lower end of the femur the periosteum is nearly always stripped off the posterior wall, and this posterior wall should be removed if sequestra are to be avoided.

Where the shaft is found completely, or almost completely, stripped of periosteum, a primary diaphysectomy may be done, or the whole medulla must be widely exposed and gouged out. If the patient's general condition is bad, a diaphysectomy is preferable, as it can be more rapidly done. If, on the other hand, the patient's condition will stand it, the chiselling away of one aspect of the entire shaft with gouging out of the medulla will leave enough compact bone to act as a splint. When the periosteum has formed enough bone to retain a good shape, a secondary diaphysectomy should be done. If the patient's condition is very critical, probably a primary amputation is the most rapid efficient operation, and should be done. It should, however, seldom be necessary.

In this series primary amputation was performed three times, with one death. In the case which died the toe was amputated for osteomyelitis, and at the *post-mortem* examination osteomyelitis of the ileum was found, which was probably primary, osteomyelitis of the phalanges being very rarely primary, though often secondary. This case also had general pyaemia with purulent pericarditis.

If, after efficient initial treatment, the case continues to go downhill, secondary diaphysectomy is worth trying, and if this fails, secondary amputation should be done.

The existence of pyaemic abscesses in either bones or soft tissues is no contraindication to amputation, as in these cases the patient often begins to get the better of his pyaemic foci as soon as the primary focus is removed by amputation. Secondary diaphysectomy sometimes gives very good results, as in a case reported by Scudder,¹⁰ and in several cases here. It is well worth a trial before proceeding to amputation, and may save a useful, though often unsightly, limb. After diaphysectomy, the periosteum should be stitched so as to draw it loosely into position over gauze packing. If, as is usually recommended, it be sewn to the skin, the newly formed bone when it grows is apt to be four or five times its normal width, as a result of the spreading out of the periosteum when the skin retracts.

The chief hope of reducing the mortality centres in early and radical operation. Possibly in the future vaccines may be instrumental in reducing the mortality. In this series only a few cases were treated with vaccines, but the results in these few cases have been anything but encouraging:

Mortality.	
General mortality of the series	... 34.6 per cent.
With vaccines, 7 deaths in 16 cases, or a mortality of	... 43.7 "

If the cases in which the bone was thoroughly opened up from the first be considered separately, one finds:

Group I.	
Without vaccines, 39 deaths in 123 cases, or 31.7 per cent. mortality.	
With vaccines, 2 deaths in 6 cases, or 33 per cent. mortality.	

If the cases in which at first only the periosteum was opened be taken, one finds:

Group II.	
Without vaccines, 50 deaths in 139 cases, or 35.9 per cent. mortality.	
With vaccines, 4 deaths in 7 cases, or 57 per cent. mortality.	

That the mortality is not higher in the whole of Group II. is due to the fact that in many cases the bone was opened up at a second operation. Possibly the very high mortality of cases treated with vaccines in this group may be due to vaccine treatment having been allowed to replace, instead of supplementing, the more important surgical measures

In conclusion, I must express my thanks to the surgical staff of the London Hospital for their permission to use the notes from which these figures have been compiled. To Mr. Walton and Mr. Benians I owe my thanks for many suggestions.

The tables on the previous page show the bone, age, and sex incidence in 298 cases.

REFERENCES.

- ¹ H. L. Barnard, *London Hospital Gazette*, February, 1903.
- ² Trendelenburg, quoted by Albee, *N. Y. Med. Journ.*, lxxxvii, 1077, 1909.
- ³ Thompson and Miles, *Manual of Surgery*.
- ⁴ Dolbey, *London Hospital Gazette*, Clinical Supplement, 1906.
- ⁵ J. E. Moore, *N. Y. Med. Journ.*, xci, 431, 1910.
- ⁶ Albee, *N. Y. Med. Journ.*, lxxxvii, 1077, 1909.
- ⁷ Ollier, *International Encyclopaedia of Surgery*.
- ⁸ Cabot, *Clinical Examination of the Blood*, 1897.
- ⁹ Ironside Bruce, *System of Surgery*, edited by Choyce, vol. i.
- ¹⁰ Scudder, *Surg., Gynec., and Obstet.*, Chicago, vi, 169, 1908.

APPENDICITIS IN PRIVATE AND PUBLIC
HOSPITALS FOR THE INSANE.*

BY

JOHN F. BRISCOE, M.R.C.S.,

ALTON, HANTS.

My excuse for a communication on this physical affection is an attempt to prove, from its rarity in private and public asylums for England, Wales, Scotland, and Ireland, that it is preventable. In the ten years from 1902 to 1911 there are recorded 75 deaths from appendicitis, typhlitis, or perityphlitis, ascertained in the majority of cases by *post-mortem* examination, among the insane detained in all the asylums of England and Wales. It would be instructive to know how many of the 75 were affected before admission into the respective asylums, and how many could have been saved by operation. For the statement that appendicitis is a rare malady in private and public hospitals I have the support of several well-known mental specialists; not only do these give me their assistance, but moreover I have the help of the pathologist to the London County Asylum at Claybury, who states, "I do not believe there has been a single case of appendicitis on the *post-mortem* table at Claybury since I have been pathologist. Consequently, in more than 2,000 *post-mortem* examinations there has been no case." The comparison of the Claybury hospital for the insane with the teaching hospitals of St. Bartholomew's and Guy's for the sane is reliable. For instance, at St. Bartholomew's Hospital 1,645 *post-mortem* examinations were made between 1909 and 1911, and 69 were designated cases of appendicitis. At Guy's Hospital in 1900, of the first 500 *post-mortem* examinations 12 were recorded as appendix cases. I find that at the same hospital there were 8,588 admissions in 1910, and of these admissions 306 were subjects of appendicitis, 187 being under the care of surgeons, while 119 were under the care of physicians. In connexion with the statistical tables from the Lunacy Boards of Scotland, it is believed that formerly death from appendicitis was usually returned as death from peritonitis, but these diseases had no separate heading in the boards' tables of causes of death, and are not yet separately notified. If a case of death from appendicitis were returned it would be placed in the table under diseases of the digestive system; but it has been ascertained that no death in a Scottish asylum has been returned during the past five years as due to appendicitis. In the report of the inspectors of lunatics for Ireland, 1911, typhlitis is grouped among the causes of death; but this table contains no return of the complaint for that year, and there were only 11 cases of peritonitis. The Secretary of the Office of Lunatic Asylums, Dublin Castle, sends me the returns of deaths from typhlitis during the period 1902 to 1910 as follows: 1902, 0; 1903, 1 (male); 1904, 0; 1905, 1 (female); 1906, 0; 1907, 1 (male); 1908, 2 (male and female); 1909, 0; 1910, 0—namely, 5 cases in nine years. The cause was ascertained by *post-mortem* examination, except in one case. The death-rate from appendicitis, as chronicled in the Registrar-General's Report for 1909, shows a slight gradual increase from 1901 to 1909.

The caecum, with the bud of the appendix, as stated by Quain, grow out together from a straight uniform piece of intestine at the sixth week of intrauterine life. The appendix is an abdominal and persistent anatomical arrangement in the plan of the permanent anchorage of the

* Read at a meeting of the Winchester Division.