THE FUNCTIONAL AND HISTOLOGICAL EFFECTS OF INTRANEURAL AND INTRAGANGLIONIC INJECTIONS OF ALCOHOL.

BY
OTTO MAY, M.D.CANTAB., M.R.C.P.LOND.,

BEIT RESEARCH FELLOW.
(From the Research Department, University College Hospital Medical School.)

SINCE its introduction by Schlösser¹ in 1903, the injection of alcohol into nerves for the treatment of neuralgia and other painful conditions has become a well-established therapeutic measure. In his earlier paper² he recommended its use, not only for sensory, but also for motor and mixed nerves, especially the facial in "tic convulsif," and the sciatic in "sciatica." In the case of motor and mixed nerves, however, he aimed, not at actual penetration, but only at injection into the neighbourhood of the nerve, a procedure which produced a paralysis lasting only half an hour to one hour. In a later paper³ he states that 80 per cent. alcohol applied locally to a nerve-trunk produced degeneration and absorption of all parts except the neurilemma. He gives no details of the histological

Fig. 1.—Cat 2. Part of section of Gasserian ganglion (polychrome blue), showing chromatolysis of some of the cells. (Drawn under $\frac{1}{3}$ in. objective.)

examination, nor any statement as to how deeply the penetration had occurred. In this same paper he gives statistics of 202 cases, including 38 of sciatica and 9 of tic convulsif (all the latter recurred in three to seven months), and adds a warning as to the danger of producing motor paralysis.

The only other experimental work on the subject which has come under my notice is that of Finkelnburg,⁴ who injected 0.5 to 1.5 c.cm. of 60 to 80 per cent. alcohol into the sciatics of dogs. A complete paralysis of the corresponding muscles was produced, remaining marked for months. Microscopically, an almost complete degeneration of the nerve bundles was found. If the injections were made only into the neighbourhood of the exposed nerve and the wound closed, well-marked motor weakness resulted, lasting several days. Examined by the Marchi method after fourteen to twenty-one days, the outermost bundles of the nerves showed a high degree of degeneration; in the central parts there was considerable black "stippling"; in addition, further off, were some slight haemorrhages under the sheath. The axones do not appear to have been examined.

The present paper deals with the functional and histological results obtained by alcohol injections into various nerves and ganglia. The experiments may be divided into three groups, as follows:

1. Injections into a pure sensory nerve—the infra-orbital.
2. Injections into the Gasserian ganglion.
3. Injections into mixed nerves—the sciatic and the anterior crural.

The animals employed were, with one exception, cats. In all cases they were fully anaesthetized during the operations, which were carried out with strict antiseptic precautions. The nerve or ganglion was exposed by careful dissection, and the alcohol injected obliquely from a syringe fitted with a fine needle. The excess of fluid was lightly mopped out, and the wound closed.

Histological Stains Employed.

1. For a general survey, extent of fibrosis, etc.—(a) Delafield haematoxylin and eosin, (b) Weigert haematein and Van Gieson.
2. For nerve cells—Unna's polychrome blue method.
3. For the medullary sheaths—Marchi's method, counterstained, in some cases, by Mann's eosin and methyl blue mixture.
4. For the axones—Cajal's and Bielschowsky's silver impregnation methods.

GROUP I.—INJECTIONS INTO THE INFRAORBITAL NERVE.

Four experiments of this nature were performed, as follows:

EXPERIMENT 1.—Cat. Floor of right orbit exposed, and 0.5 c.cm. of 80 per cent. alcohol injected into (?) the infraorbital nerve.

Fig. 2.—Cat 2. Section of part of infraorbital nerve from the floor of the orbit (haematein and Van Gieson) (photograph $\times 40$), showing the bundles of degenerated nerve fibres embedded in newly formed fibrous tissue.

The animal was again anaesthetized seventeen days later, and both infraorbital nerves stimulated by the faradic current at their points of exit from the foramina. No difference could be detected in the result on the two sides. The animal was then killed and the nerves, etc., examined as below:

Macroscopically, a tract of haemorrhage was seen on one of the branches of the nerve on the floor of the orbit.

Microscopically, no obvious changes were found either in the Gasserian ganglion, or in the parts of the infraorbital nerve central and peripheral to the point of injection. Sections at the site of haemorrhage showed a considerable amount of blood lying round the bundles of the nerve, but no obvious change in the bundles themselves.

The intracerebral roots of the nerve showed no trace of degeneration.

EXPERIMENT 2.—Cat. Experiment as in 1.

Twenty-one days later the animal was again anaesthetized and both infraorbital nerves stimulated. The left produced the usual reflex quickening of respiration, the right practically no effect.

The *post-mortem* examination showed that the nerve in the floor of the orbit was embedded in dense fibrous tissue firmly adherent to the bony floor.

Microscopically, the results were as follows:

(a) *Gasserian Ganglion*.—Some groups of cells showed definite chromatolysis, with eccentricity or disappearance of nucleus,

* Except in Experiments 3 and 4, in which the alcohol was injected without preliminary dissection.

breaking up of the chromatin granules, etc. (Fig. 1). These cells were chiefly in the part of the ganglion adjacent to the middle division, but some were scattered among other groups of cells.

(b) *The part of the infraorbital nerve in the scar* showed great fibrosis with implication of the bundles to varying degrees and corresponding destruction of nerve fibres (Fig. 2).

(c) *Proximal* to this the nerve was apparently normal (both axones and sheath).

(d) *Distal* to the scar the nerve was completely degenerated; no normal fibres could be seen, and the axones had practically entirely disappeared. There was great proliferation of the nuclei of the sheath.

(e) *The intracerebral roots* showed no degeneration.

EXPERIMENT 3.—Cat; 0.5 c.cm. of absolute alcohol injected through the skin into the left infraorbital foramen (under ether anaesthesia).

On the following day there was some protraction (paralysis) of the corresponding nictitating membrane, which passed off in the course of a day or two. No other symptoms were noticed.

Nineteen days later the animal was anaesthetized with ether and both infraorbital nerves exposed distally. On faradic

stimulation the effect on respiration seemed less marked on the left side than on the right.

Macroscopically nothing abnormal could be seen.

Microscopically the results were as follows:

(a) *Gasserian Ganglion.*—Some cells showed eccentricity of nuclei and disappearance of chromatin; none, however, had the shrunken, deeply stained appearance characteristic of complete destruction.

(b) *Infra-orbital Nerve.*—In the anterior part of the floor of the orbit it showed, by Marchi's method, degeneration of the majority of its fibres; some had, however, escaped, and showed the normal appearance. In some of the former group the degeneration appeared incomplete—stretches of normal medulla with scattered globules

Fig. 3.—Cat 4. Section of part of infra-orbital nerve from the floor of the orbit (haematein and Van Gieson (photograph $\times 40$), showing destruction of the nerve bundles without much fibrosis; one bundle (a) has, however, almost entirely escaped.

of black-staining fat. The proximal part of the nerve at the sphenoidal fissure showed no degeneration.

(c) *The brain-stem* showed no degeneration.

EXPERIMENT 4.—Cat. Procedure as in (3); symptoms identical.

Twenty days later the animal was anaesthetized with ether, and both infraorbital nerves stimulated as in (3). No definite difference could be observed—that is, respiration appeared to be quickened in both cases.

Macroscopically, no obvious change.

Microscopically, the results were as follows:

(a) *Gasserian Ganglion.*—Changes practically identical with those in Animal 3.

(b) *Infraorbital Nerve.*

1. Branches emerging from the foramen. By Marchi's method these showed a very considerable amount of degeneration with some normal fibres left. These nerves were very difficult to tease, due apparently to an increase of fibrous tissue.

2. In anterior part of floor of orbit. Sections stained by haematein and Van Gieson showed destruction of the nerve bundles with some increase of fibrous tissue and considerable round-celled infiltration. One bundle had, however, partly escaped (Fig. 3).

3. The nerve just central to 2. Examined by Marchi's method, many of the fibres showed the junction of the pathological and the normal parts (Fig. 4)—that is, the distal part showed numerous black globules, getting gradually less as the fibre was traced centrally, until a normal medullary sheath was reached. There were more completely normal fibres here than in the more distal part—that is, the alcohol effect had spread back unevenly, having extended further along some fibres than others.

(c) *The brain-stem* showed no degeneration.

In considering the results obtained in these four experiments one is struck by the fact that in No. 1 the nervous structures completely escaped injury. Apparently the needle entered the nerve, producing the haemorrhage described, but then must have slipped out of it before any alcohol had entered. This is an accident which presumably occurs not infrequently in clinical work, in those cases in which the alcohol fails to relieve the pain, or relieves it only for a very short time. In the three other experiments a considerable effect was produced in each

case—a more or less complete degeneration of the peripheral part of the infraorbital nerve. In No. 2, in which the alcohol was injected by the "open method"—that is, after exposure of the nerve by operation—the degeneration was complete, and the whole nerve was found imbedded in dense new fibrous tissue. In 3 and 4, in which the injection was made by the method employed clinically, the degeneration was not complete, though quite considerable in extent. In none, however, had the degeneration "ascended"; the nerve proximal to the part affected by direct contact of the alcohol was in each case normal. The Gasserian ganglion showed a similar change in all three cases—chromatolysis in some groups of cells, but no sign of actual cell death; in other words, no cells exhibited the shrunken, deformed, deeply-staining appearance suggestive of complete destruction. In harmony with this, there was total absence of change in the proximal roots of the trigeminal nerve, both extra- and intracerebral.

The conditions resulting from such injections may be summarized as a local destruction of fibres by the alcohol, with some consequent chromatolytic change in their cells, but without any ascending degeneration or nerve-cell necrosis. Such conditions are comparatively favourable to regeneration, as will be shown below, and this chemical section of the nerve is probably followed more quickly by regeneration than is mechanical section, or, better, resection of part of the nerve.

GROUP II.—INJECTIONS INTO THE GASSERIAN GANGLION.

The four following experiments were performed:

EXPERIMENT 5.—Cat. Left Gasserian ganglion exposed by the temporal route, and injected with about 0.75 c.cm. of 80 per cent. alcohol.

Seven days later the animal was killed, and the brain perfused *in situ* with formol-Müller.

Microscopically, the results were as follows:

(a) *Left Gasserian Ganglion.*—Sections were stained both by polychrome blue and by haematein-Van-Gieson. The most marked change was in the part free from nerve cells, that is,

These operations were kindly performed for me by Sir Victor Horsley, to whom my best thanks are due.

Fig. 4.—Cat 4. Some teased fibres of the infra-orbital nerve (Marchi) (photograph $\times 80$), showing the junction of the normal part (a) with the degenerated part (b).

the commencement of the division into the three main branches. Here were small haemorrhages, and patches of round-celled infiltration, with obvious destruction of nerve fibres. This was most marked at the base of the middle division, the superior and inferior being barely affected. The nerve cells, by comparison with the normal side, showed little change. Most seemed quite unaffected, but a few showed marked shrinking and dark uniform staining, suggestive of destruction.

(b) *Trunk of Middle Division.*—Marchi staining showed a considerable amount of early sheath degeneration, the medulla being broken up into large dark-brown blocks, many of which contained fine black granules. There were also a few normal fibres left. Cajal staining showed a marked degree of "axolysis" in the majority of fibres.

(c) *Trunk of Inferior Division.*—Most fibres were normal, only a few showing a slight grade of degeneration.

The superior division was not examined.

(d) *The brain-stem* showed a considerable amount of degeneration in the spinal root of the trigeminal; the mesencephalic root appeared normal.

EXPERIMENT 6.—Cat. Operation as in (5). No symptoms resulted.

Eighteen days later the animal was killed, and the brain perfused with formol-Müller.

Microscopically, the results were as follows:

(a) *Left Gasserian Ganglion.*—Here again the cells were surprisingly little affected, most of them looking quite normal, or showing moderate chromatolysis (Fig. 5). Only a few appeared to have undergone actual necrosis. As in (5), there was con-

Fig. 5.—Cat 6. Part of section of Gasserian ganglion (polychrome blue), showing many normal cells. (Drawn under $\frac{1}{2}$ in. objective.)

siderable destruction of fibres, with patches of round-celled infiltration and fibrosis.

(b) *Trunk of Middle Division.*—Many bundles had undergone complete degeneration, but a few had escaped.

(c) *Trunk of Inferior Division.*—This showed almost complete degeneration (Marchi and Bielschowsky), only a few fibres having escaped.

The superior division was not examined.

(d) *The Brain-stem.*—The spinal root of the trigeminal showed practically complete degeneration (Fig. 6). A few degenerated fibres were present also in the mesencephalic root of the nerve.

EXPERIMENT 7.—Cat. Operation as in (5), except that absolute alcohol was used instead of 80 per cent. No symptoms were observed.

Sixteen days later the animal was killed and the brain perfused with formol-Müller.

Microscopically, the results were as follows:

(a) *Left Gasserian Ganglion.*—Here again the most obvious change was in the commencement of the main divisions, a large part of the middle and a smaller part of the inferior trunk showing disorganization. The cell changes were similar to those in Cat 6—that is, the majority apparently normal, but some showing the shrinkage, distortion, and deep staining suggestive of destruction. The ganglion as a whole showed only slight fibrosis but considerable round-celled infiltration with one or two patches of necrosis.

(b) *Trunk of Middle Division.*—Most of the fibres showed marked degeneration, a few normal ones being scattered between these.

(c) *Trunk of Inferior Division.*—Most of the fibres were normal, only a few showing degeneration.

(d) *The Brain-stem.*—The spinal root of the trigeminal showed marked degeneration, but not so complete as in (6). The mesencephalic root showed only slight degeneration.

Recently Wilfred Harris⁵ has published a paper giving an account of several cases of tic douloureux treated by injections of alcohol into the Gasserian ganglion through the foramen ovale. He states that there is clinical evidence of practically complete destruction of the ganglion as a result of a single injection. The experiments recorded above hardly support this contention. In not one of these was there anything approaching this result; in each case a very large number of cells remained histologically normal. Sections of the ganglion showed definite changes, but the effect on the fibres seemed relatively more marked than on the cells. Thus, in Experiment 5 the middle division was largely degenerated, in Experiment 6 the middle and inferior divisions, and in Experiment 7 chiefly the middle division.

A very striking feature in each of these was the considerable amount of degeneration found in the spinal root of the trigeminal. This was quite out of proportion to the extent of cell destruction in the ganglion. In Experiment 6 the root was practically completely degenerated, while the majority of cells appeared to have escaped. The explanation of this fact is not at first sight obvious. Experiments 1-4 in Section I show that damage to the axones peripheral to the cells does not produce retrograde degeneration in the central processes constituting

Fig. 6.—Cat 6. Section of upper part of medulla (Marchi) (photograph $\times 5$), showing the complete degeneration of the spinal root of the trigeminal (a).

the spinal root. Hence, in the absence of complete cell destruction, we are driven to the conclusion that the alcohol must have passed centrally and reached the proximal root of the nerve without actually infiltrating the whole ganglion. In other words, it must tend to creep along the superficial parts of the ganglion under its fibrous sheath, apparently following the line of least resistance. In order to investigate this manner of spread more clearly, it was decided to repeat the experiment on a larger animal in which the structures approximated more closely in size to those of man. For this purpose the following experiment was performed on a young goat, nearly full-grown:

EXPERIMENT 8.—Goat. Right Gasserian ganglion exposed by the temporal route, and injected with about 0.75 c.cm. of absolute alcohol.

Eighteen days later the right eye showed very pronounced "neurotrophic" changes—a marked keratitis and conjunctivitis. The animal was killed, and the brain perfused *in situ* with formol-Müller.

Macroscopically, the right ganglion was found apparently much larger than the left, this increase being due to a large amount of new fibrous tissue surrounding it, and rendering it adherent to adjoining structures.

Microscopically, the results were as follows:

(a) *Right Gasserian Ganglion.*—Sections showed a fair number of profoundly altered cells, but here, again, the large majority appeared normal. In the neighbourhood of the destroyed cells there were patches of fibrosis and round-celled infiltration.

(b) *Trunk of the Superior Division.*—Stained by the Marchi and Cajal methods, this showed a considerable amount of degeneration, amounting to perhaps half its fibres.

(c) *Trunk of the Middle Division.*—This showed only a few degenerated fibres.

(d) *Trunk of the Inferior Division.*—This was practically free from degeneration.

(e) *The Root of the Trigeminal* (proximal to the ganglion).—This showed very considerable degeneration, amounting to at least half its fibres.

(f) *The Brain-Stem.*—Here the appearances confirmed (e). The spinal root showed wide degeneration, amounting in the pons to approximately half its fibres. Further back, in the medulla, this proportion diminished, so that at the level of the pyramidal decussation it was about a quarter of the whole.

This experiment confirms the conclusion arrived at from the results of Nos. 5-7, that the alcohol when injected into the ganglion tends to spread centrally to the proximal root without destroying the larger number of cells *en route*. The amount of "central" degeneration was undoubtedly far in excess of the sum total in the three peripheral trunks, and disproportionate to the cell destruction. This raises the interesting question: Are these fibres of the spinal root capable of regeneration? It is usually held that such is not the case, that fibres in the central nervous system, even those of exogenous origin, do not undergo complete regeneration. If this is true, then from the therapeutic point of view a complete section, mechanical or chemical, of the root of the trigeminal nerve proximal to its ganglion, should be equivalent to removal of the Gasserian ganglion itself. I am at present engaged in trying to elucidate how far this is true—to what extent regeneration can occur in the cord after section of posterior roots, etc. Clinical experience is against the view that simple section of the trigeminal nerve proximal to the ganglion is as effective as removal of the latter; the possibility must not be lost sight of, that pain impulses might be carried into the brain by fibres only imperfectly regenerated—for example, fibres which consist of an axone without a morphologically perfect medullary sheath.

GROUP III.—INJECTIONS INTO MIXED MOTOR AND SENSORY NERVES.

Eight such experiments were performed, of which the following is a brief summary:

EXPERIMENT 9.—June 28th, 1910. Cat. 0.5 c.cm. 80 per cent. alcohol injected into left sciatic in lower part of thigh; 0.5 c.cm. normal saline similarly into right sciatic.

Immediately after operation both hind legs were weak, so that the cat was plantigrade, with both heels to the ground and a tendency to turn the foot over in walking.

July 1st. Right leg practically well; left leg *in statu quo*.

Fig. 7.—Cat 9. Photograph taken a month after injection of alcohol into the left sciatic nerve, showing the weakness of the left hind-leg. Two months later recovery was complete.

in the nerve. By the Marchi method, a few fibres of the left posterior tibial nerve showed some fine black granules, but most were normal. The rest of the tissues were not examined.

Fig. 8.—Cat 13. Section of left posterior tibial nerve (Cajal stain); on the left of the section are seen the normal axones, and on the right the fine, newly regenerated ones. (Drawn under $\frac{1}{2}$ in. objective.)

EXPERIMENT 10.—June 30th, 1910. Cat. 1 c.cm. 90 per cent. alcohol injected into left sciatic nerve in mid-thigh.

July 1st. Cat walks with left heel to ground. (Cf. Experiment 9.)

August 3rd. Still in same condition.

August 12th. Leg showing definite improvement; heel beginning to be raised in walking.

August 26th. Walking is now almost normal.

September 26th. (Eighty-eight days after operation.) No difference between the two legs. Cat killed; weights of the two gastrocnemii equal (10.5 grams). Macroscopically, the left sciatic nerve looked normal; no inflammatory adhesions were present.

By Marchi method, the left position tibial nerve showed some bundles quite healthy, while in others there were slight remains of degeneration in the presence of fine black granules, lying for the most part in "phagocytic" cells between the fibres. The sciatic trunk showed numerous patches of round-celled infiltration, chiefly at the periphery of the bundles; the fibres themselves looked normal.

EXPERIMENT 11.—July 18th, 1910. Cat. 0.75 c.cm. 50 per cent. alcohol injected into left sciatic in mid-thigh. No weakness of the leg was produced, the animal walking quite well on the following day. It was kept eighteen days without developing any symptoms, and was then killed.

EXPERIMENT 12.—October 26th, 1910. Cat. 0.5 c.cm. 80 per cent. alcohol injected into left sciatic in mid-thigh.

October 28th. Cat limps slightly, with left heel lower than right.

November 7th. Weakness less marked.

November 14th (nineteen days after operation). No difference observable in the two hind legs. Killed. No macroscopic change in the nerve.

Microscopically, there was scattered degeneration in both the peroneal and posterior tibial nerves, involving only a minority of the fibres. The axonal staining showed most axones to be normal, but a few were broken up into irregular fragments.

EXPERIMENT 13.—November 3rd, 1910. Cat. 1 c.cm. 90 per cent. alcohol injected into and around left sciatic nerve in mid-thigh.

November 7th. Marked ataxia of left hind-leg; left foot is dragged, and the animal tends to walk with the foot turned over, so that

the dorsum is in contact with the ground.

December 8th. The calf muscles are greatly wasted.

December 18th. The walking has improved, though the wasting of the calf muscles is still very marked.

January 6th, 1911. The cat walks well, without any obvious ataxia.

Sixty-four days after operation, the left gastrocnemius, tested by Leduc's current (6), showed partial "reaction of degeneration." Killed. Right gastrocnemius 30.4 grams; left gastrocnemius 19.4 grams.

Microscopically, the results were as follows:

(a) *Left sciatic* (above injection, Cajal stain), quite normal.

(b) *Left posterior tibial*: Cajal: A patch at periphery of normal large axones; the larger part of the nerve showed disappearance of these, with a few fragments of debris and numerous fine new regenerating axones (Fig. 8). Marchi: Marked advanced degeneration in the greater part of the nerve; a few normal fibres were, however, present.

EXPERIMENT 14.—June 1st, 1911. Cat. 1 c.cm. 90 per cent. alcohol injected into and around left sciatic nerve in mid-thigh. Very little appeared to have penetrated the nerve.

June 6th. Wound healing well, but cat shows no symptoms at all from the injection.

June 7th (six days after operation). Killed. No macroscopic changes found in the nerve or muscles. No microscopic examination made.

EXPERIMENT 15.—June 1st, 1911. Cat. 1 c.cm. 90 per cent. alcohol injected into and around left anterior crural nerve just above Poupart's ligament.

June 4th. Marked weakness of the quadriceps extensor of the left thigh.

August 1st (sixty-one days after operation). Very marked wasting and weakness of these muscles.

The cat was anaesthetized and the left rectus femoris examined with Leduc's current. It showed the typical "reaction of degeneration" (complete).⁶ Faradic stimulation

of the trunk of the anterior crural nerve produced no contraction of the muscles. Killed. Weight of right quadriceps, 48 grams; weight of left quadriceps, 20 grams.

Microscopically, the results were as follows:

(a) *The Anterior Crural Trunk*.—At the site of injection the nerve showed profound changes—a massive new formation of fibrous tissue, with patches of round-celled infiltration, involving and destroying the greater proportion of the nerve fibres (Fig. 9). A few fibres had escaped, at any rate in these particular sections, but the destruction was very extensive—a veritable “nerve cirrhosis.” Above this point, Cajal and Marchi preparations showed a practically normal nerve; the process is a local disorganization, not leading to any ascending change.

(b) *The distal branches* (by Marchi, Cajal, and Bielschowsky methods) showed active preparation for regeneration—the long nucleated protoplasmic columns (the “apoptrophic cells” of Marinesco), without, however, any new axones. It looked as though the marked fibrosis at the site of injection had delayed regeneration, so that the process was slower than, for example, that after successful primary nerve suture.

Fig. 9.—Cat 15. Section of the anterior crural nerve at the site of injection (haematein and Van Gieson), (photograph $\times 40$), showing profound disorganization of the nerve, with great fibrosis. Two bundles of fibres which have escaped complete destruction are marked (a).

EXPERIMENT 16.—June 6th, 1911. Cat. 0.5 c.cm. absolute alcohol injected into and around the left sciatic nerve in mid-thigh.

July 8th. Marked plantigrade gait of left hind foot, with tendency to walk with the dorsum of foot to the ground.

July 26th. Weakness and wasting of the left calf muscles still very marked (fifty days after operation). The cat was anaesthetized, and the left gastrocnemius examined with the Leduc current. It showed complete “reaction of degeneration.” There was no muscular response to faradic stimulation of the sciatic trunk either above or below the site of injection. The animal was then killed. Weight of right gastrocnemius, 23.2 grams; weight of left gastrocnemius, 8.5 grams.

Microscopically, the results were as follows:

(a) *The sciatic trunk* at the site of injection; the changes were practically identical with those found in the preceding experiment—that is, great fibrosis both between and in the nerve bundles. There were also the remains of numerous haemorrhages in both these situations.

Immediately above this point Cajal preparations showed greater regeneration activity, axones splitting up into complex bundles of fine fibrils, etc. (Fig. 10). Further up the appearances were entirely those of a normal nerve—that is, there was no trace of any ascending degeneration.

(b) *The peripheral branches* showed no new axones—that is, here again the fibrosis has proved a severe obstacle to the down-growth of new axones from the proximal part. They showed the usual appearances of late Wallerian degeneration—the

presence of numerous comparatively fine fat granules, some lying free, others contained in the protoplasm of the invading phagocytes.

A striking feature brought out by a consideration of

these experiments is, as in Group I, the variability of the results. Thus, in Nos. 11 and 14 the injection produced no obvious effect, in No. 12 only a slight transitory weakness, while in 9, 10, 13, 15, 16 the functional results were severe and more lasting. The technique was, as far as possible, identical in all these experiments, but the strength of alcohol used was not always the same. In No. 11 it was only 50 per cent., which may partially account for the lack of disturbance produced. On the other hand, in both 9 and 14 it was 90 per cent., yet the effects in these two cases afford a striking contrast. There can be not the slightest doubt that the explanation lies very largely in the difficulty of ensuring adequate penetration of the nerve

by the alcohol. The texture of the nerve in question has an obvious bearing on this point. The cat's sciatic is composed of several bundles held lightly together by loose connective tissue; such a nerve is more

difficult to infiltrate successfully than one of a more compact structure—for example, the posterior tibial. Further, if this difficulty is so prominent when the nerve is exposed by dissection, it becomes still more so when the injection is made through a skin puncture, as is the usual clinical method. On the other hand, when the injection is made into a bony canal (as in the case of the three divisions of the trigeminal), or into a space bounded by dense fibrous structures (as in the case of the Gasserian ganglion), it may well be that the alcohol, by remaining longer in contact with the nerve or ganglion, may produce comparatively more effect, even though little had penetrated during the actual injection.

The muscles supplied by the nerves operated upon in Experiments 9, 10, 13, 15, 16, showed the usual atrophic changes, identical with those following

Fig. 10.—Cat 16. Section of the sciatic nerve just above the point of injection (Cajal stain); this shows the axones splitting up into complex bundles of fibrils—a phase of active regeneration. (Drawn under $\frac{1}{2}$ in. objective.)

mechanical section of the nerve. It is interesting to note, however, the fairly rapid recovery which can ensue. Thus, in Experiment 10 this was complete when the animal was killed after

eighty-eight days. The gastrocnemius muscle weighed the same as that on the control side, and reacted equally well. Moreover, the animal had lost all trace of the limp which was so prominent for some weeks after the operation. Unfortunately, the histological examination was incomplete, as the branches to the gastrocnemius were not examined to see how far they had suffered and how far recovery was due to a true nerve regeneration. In Experiments 13, 15, and 16, however, the investigations were more complete; in 13 (sixty-four days) regeneration was shown histologically to be well advanced, and the muscles had so far recovered that, examined by the Leduc current, they reacted in a manner quite different from the typical reaction of degeneration. In 15 (sixty-one days) and 16 (fifty days) regeneration was much less advanced. In neither of these could any new axones be found in the peripheral branches, and the muscles gave the complete R.D. It is probable that this backwardness, contrasted with 13, is dependent on the greater degree of fibrosis produced by the alcohol in the former, the fibrous tissue forming, as I have found in other experiments, an effective barrier against the downgrowth of the regenerating axones. There is every reason to believe, however, that in due time complete recovery of power would occur in all muscles which had been paralysed by such injections. In other words, there does not seem to be any real danger of permanent paralysis in these cases, though in man the time taken for complete recovery might be a matter of a year or more. This is a matter of some interest, in view of the cases of peroneal paralysis recorded by Schlösser and others (v.a.) following the treatment of sciatica by these means, though such a procedure is, I fancy, no longer likely to be advocated at the present time.

CONCLUSIONS.

1. Alcohol, injected into the trunk of a peripheral nerve, produces a more or less complete local necrosis of the nerve at the point of injection.

2. The change is not an "ascending" one, the nerve above the point of injection remaining normal; the cells of origin of the fibres may show some degree of chromatolysis, but do not exhibit signs of permanent injury.

3. The conditions produced by such injection are more favourable to regeneration than those resulting from simple section without suture. The anatomical continuity of the nerve trunk favours rapid regeneration, though this is to some extent retarded by the fibrosis which occurs to a greater or less extent in every case of alcohol injection.

4. It is apparently impossible by a single injection of alcohol to produce complete necrosis of the Gasserian ganglion, its dense texture preventing complete infiltration. The alcohol tends to find its way under the sheath of the ganglion towards the proximal root, which is affected to a greater degree than the actual ganglionic cells.

REFERENCES.

- ¹ Schlösser, *Klin. Monatsblatt f. Augenheilkunde*, 1903, Bd. 41 (2), s. 244.
² *Ibid.*, *Versammlung d. Deutscher Ophthal. Gesellsch.*, Heidelberg, September, 1903. ³ *Ibid.*, *Verhandlung des Congresses f. Innere Medizin*, Wiesbaden, 1907, s. 49. ⁴ Finkelnburg, *ibid.* ⁵ W. Harris, *Lancet*, 1912, vol. i, p. 218. ⁶ O. May, *Brain*, 1911, vol. xxxiv, p. 272.

GREY HAIR ASSOCIATED WITH NERVE LESIONS.

By G. LENTHAL CHEATLE.

In previous papers¹ I have pointed out that when the sites of the points of incidence and areas of occupation are taken into consideration, then there is reason to believe a nervous influence exists which can induce greyiness of hair, at least in some cases. The particular cases to which I drew attention were of two classes.

Class 1. Persons having had grey patches of hair ever since they could remember.

Class 2. Persons who gradually become grey with advancing age. This class of case must again be subdivided into:

(a) The initial occurrence of patches of grey hair, which gradually fuse with the general greyiness of the scalp and beard.

(b) The gradual appearance of scattered grey hairs beginning in the temporal regions. In this subdivision there is an absence of patches.

It is those cases in Class 1, and (a) of Class 2, which show by the points of incidence a nervous genetic influence. In Class 1 the grey patches are often hereditary as well.

In the present article I wish to indicate a large class of cases which acquire greyiness of hair in association with other nerve lesions, and therefore probably connected with nervous changes.

Fig. 1 shows the nerve lesions on the right side of a case of Sir Victor Horsley's, to whom I owe my thanks for allowing me to publish it. The clear area on the forehead and frontal region of this scalp was anaesthetic, the skin was bleached and the hair was grey. The dotted area posterior to it was hyperaesthetic and the hair was not grey. In both areas there was slight overlapping of the left side, which otherwise was normal with regard to sensation and the colour of the hair. The case suffered from leucoderma.

Fig. 1.

Fig. 2.

Fig. 2 shows the state of things in a case of my own, an out-patient of King's College Hospital, aged 36 years. A subcutaneous neurofibroma growing on each of the sites marked by a cross on the cheek and forehead of the right side, and which were noticed about the same time ten years ago, when the hair also began to grow grey.

The state of the greyiness of hair was as follows: On the mid-orbital maximum point of Dr. Henry Head, there was a patch of normal brown colour, which appeared as an island in the completely grey V-shaped area seen in Fig. 2, behind which, and limited behind by the posterior boundary of the fifth nerve, was an area of scattered grey hairs. There was no disturbance of sensation. Tactile, heat, cold, and pain sensations were normal.

It would be of immense theoretical and practical interest to discuss the state of greyiness of hair, together with the protopathic nerve influence described by Dr. Head. The discussion would be very discursive, owing to the fact that our knowledge is not sufficient to enable us to state what makes hair appear grey; and although one may have opinions upon the subject they are not sufficiently matured for dogmatic pronouncement.

All I want to do now is to show that here is an epithelial change in state, which takes place *pari passu* with a change in the nervous system, and, therefore, probably due to it.

Although greyiness of hair is of itself an unimportant epithelial change, it appears at a time of life common to another and important epithelial change—namely, cancer. The genesis and etiology of an unimportant change in a tissue may illuminate the genesis and etiology of an important change in the same tissue.

I am, at any rate, assuming a correct attitude in saying that because a nerve influence can be shown at work in the genesis of one epithelial change, it ought, therefore, to be considered in the genesis and etiology of another epithelial change—that is, cancer.

Greyiness of hair may be said to be a degenerative change and cancer a proliferative change. The proliferative change in cancer can hardly be considered an object lesson in regeneration; its products are nearer degeneration products, so far as size, shape, and function of the cells are concerned.

REFERENCE.

- ¹ The Incidence of the Hair's Greyiness, *BRITISH MEDICAL JOURNAL*, July 22nd, 1905. The Mental Nerve Area and its Relation to Greyiness of Hair, *BRITISH MEDICAL JOURNAL*, July 4th, 1908.