forgotten that there is a tendency for the type of scarlet fever to become milder.

Dr. Foord Caiger, in his article on scarlet fever in *Allbutt's System of Medicine*, states that in London malignant scarlet fever is very rare. On the other hand, Dr. Claude Buchanan Ker points out that toxic cases formed approximately 0.6 per cent. of the total cases of scarlet fever admitted recently into the wards of the Edinburgh City Hospital. As, however, he includes in these statistics cases which die as late as the fifth day after admission, it seems probable that the majority are not of the same degree of malignancy as the cases already described. So far as the proportion of malignant to the total number of cases of scarlet fever is concerned the statistics of hospitals for infectious diseases are of little value, for of the most toxic cases many must succumb before it is possible to make the diagnosis and take steps to have them removed, while many others must be too ill to permit of their removal.

Dr. Ker* describes three varieties of toxic scarlet fever:

1. The fulminant type is very rare. Patients die with extreme rapidity, usually within twenty-four hours from the onset; cases are said to have occurred in which death resulted after a few minutes' illness. The attack commences very acutely and suddenly with intense headache, persistent vomiting, and profuse diarrhoea. The temperature is often high at first, perhaps sinking below normal in the later stages. The rash is absent, having scarcely time to make its appearance.

2. The ataxic form is not quite so severe as the fulminant, but in other respects is similar. Vomiting, diarrhoea, and delirium are severe, and the temperature keeps high. The rash appears late, or is atypical in character and distribution. The patient succumbs in from two to four days.

3. In the adynamic type the heart and circulatory symptoms are more prominent than the nervous. In this form also both vomiting and diarrhoea may be severe. The pyrexia is moderate in degree, and the rash may be well marked and possibly petechial. Death usually takes place on the fourth or fifth day.

In the forms just described the case is malignant from the beginning, but in other cases malignancy supervenes on an apparently simple attack.

Both of my cases which ended fatally belong to the first rather than the second group, while the case which recovered—and recovery is quite the exception—presented features more characteristic of the ataxic than the adynamic form.

It is undoubtedly true that in scarlet fever family susceptibility plays a part which is not evident in the other acute exanthemata. This fact is well illustrated in the series of cases narrated. Henoch, in the lectures previously referred to, remarks that "striking to me appeared the fact, noted undoubtedly by many other physicians, that if scarlet fever breaks out in a family and one of the children dies of the malignant form, very frequently also a second and a third child are taken off under similar conditions, and in this way whole families die out." It is worthy of note that in my cases the three patients so severely affected were all boys, who were nearest of age to one another, and were attacked with most startling suddenness on successive days. The child first affected, who evidently spread the infection, suffered from a mild form of the disease. It seems most probable, therefore, that the infection was not of great virulence but rather that the older boys possessed very slight powers of resistance, and were at once overwhelmed.

After puberty the susceptibility to infection diminishes. This fact also is well demonstrated in this series of cases. All the children who came into contact with the disease suffered, and most of them in a severe degree, while the mother and father and the nurse all escaped, in spite of the fact that the whole house was infected, and that they must have been constantly in the closest possible contact with the infection for many weeks.

Among other points of interest may be noted the late appearance of the rash in Frank's case. His illness commenced on September 4th, and it was not till September 8th that the rash appeared. Till then the nature and cause of the illnesses had not been clear. The diagnosis of scarlatina had been excluded when Fred's illness com-

menced, and ten days elapsed before the nature of the mysterious sequence of cases was made manifest, both deaths having occurred several days before. Most authorities agree that a definite diagnosis of scarlet fever is almost impossible before the rash appears. When the first toxic case was seen, ptomaine poisoning was at first suspected, and the help of the medical officer of health was sought. The absence of any source of such poisoning, and the occurrence of the other two cases on successive days sufficed to negative this condition.

It is only fair to say that I had had then only a small experience of scarlet fever. On looking back I feel that the first case should at least have been regarded with suspicion, and promptly isolated.

I desire to tender my thanks to Dr. G. D. Todd, of Selby, whose assistant I was at the time, for permission to publish the cases.

RECURRENT MONSTRIPARITY.

BY

E. NAPIER BURNETT, M.D., M.R.C.P.ED., F.R.C.S.ED.,

NEWCASTLE-ON-TYNE.

MRS. G., aged 35, a strong, healthy woman, gave birth to a full-time female child having a large cephalocele presenting from the vertex. The first pregnancy went to term, when a normal male child was born. The second pregnancy ended in abortion at the end of the third month; the fetus was anencephalic. The third pregnancy ended in abortion at the fifth month; the fetus (male) was anencephalic. The fourth pregnancy concluded at term, and is the subject of the present note.

On May 20th, 1910, when in her thirty-eighth week of pregnancy, the patient had a slight uterine bleeding which lasted a few hours. On vaginal examination no placenta could be felt, and the haemorrhage ceased with rest in bed.

The first stage of labour was somewhat protracted, the presenting part being high up and difficult to reach. The membranes could be felt as if being pushed to one side of the os by the presenting part, and on their being ruptured the presenting part progressed with each pain.

The second stage was delayed, no further progress being noticed, although the pains were vigorous.

I was asked to see the case in consultation. After the patient was anaesthetized and the bladder emptied I found it impossible to make out the nature of the presentation—it was a large soft rounded mass, the size of a fetal head.

At first I suspected the case to be one of ovarian dermoid obstructing labour, but the abdominal examination did not support this view. With the whole hand in the vagina I was able, on getting behind the mass, to recognize a vertex presentation with the tumour attached to it. Labour being still delayed, I incised the presenting mass and emptied out a large quantity of brain tissue, which showed both cerebral and cerebellar structure within the sac; the walls were $\frac{3}{4}$ in. in thickness. My fingers could now reach the cranial vault, and through an opening in the vertex I could insert three fingers into what appeared a normal cranial cavity containing brain tissue. There was no evidence of hydrocephalus either in the supernumerary cerebral structure or within the cranium. The labour terminated as a difficult forceps case.

The child was of female sex, with very large shoulders and short neck. The eyes were protruding and the nose flattened. There was very little liquor amnii, otherwise there was no associated malformation. The child lived about half an hour after delivery.

The third stage was completed normally, the placenta and membranes showed nothing unusual and there was no evidence of any adherent amnion.

The cephalocele was situate in the usual position of such tumours—namely in the mid-line in the region of the posterior fontanelle; the parietal bones were widely separated as if the two fontanelles were one.

Cephaloceles are comparatively rare, much more so than the anencephalic fetus. Ballantyne quotes Trélat as having had three cases in 12,900 labours, and in his own series of 325 specimens of antenatal deformity there were only eight cases of cephalocele, in contrast with forty-six of anencephaly.

There was nothing in the mental or physical condition of the patient throughout the whole of pregnancy to lend any support to the theories of external or internal maternal traumatism.

* In his textbook on *Infectious Diseases*.

Morgagni's theory, which seeks to account for these cephaloceles as being due to increased intracranial tension, especially in the form of hydrocephalus, seems every now and again to take on a fresh wave of popularity with the profession. In this case there was no evidence of such causation.

Examination of the specimen shows that it was of the nature of an accessory brain formation, lying outside, and distinct from, the intracranial cerebral tissue. Ballantyne¹ observes that

we must not think of intracranial parts being pushed out at cranial apertures and so becoming extracranial; we must imagine parts of the anlage of the brain (that is, the cerebral vesicles and the undifferentiated membranes) lying originally outside what is to be the cranium and developing in their ectopic position: if it be asked what is the cause of the accessory part, we have no satisfactory answer, but the idea of a purely germinal (pre-embryonic) factor suggests itself.

A further interesting feature to be noted is the deficiency of brain tissue (anencephaly) in the two previous fetuses, and in the present case Nature seems to have gone to the other extreme in providing cerebral tissue much in excess of the normal.

REFERENCE.

- ¹ Ballantyne, *Antenatal Pathology and Hygiene*, vol. ii, 1904, p. 359.

TURPENTINE ENEMATA IN THE TREATMENT OF ENTERIC FEVER.

By W. J. J. ARNOLD, M.B., B.Ch. R.U.I.,
CIVIL SURGEON, ST. HELENA.

It is probably not incorrect to say that none of the various methods of treatment have hitherto been successful in definitely controlling the course or limiting the duration of enteric fever.

The supreme importance of early recognition of the disease, the value of skilled nursing, regulation of diet, and attention to oral hygiene, must be accepted as first principles. In the further treatment, a measure employed at the Civil Hospital, St. Helena, during the past four or five years has resulted in a distinct shortening of the average duration of illness, and in a remarkable amelioration of all the symptoms. This particular measure consists in the administration, by the bowel, of turpentine and olive oil from the first day of treatment, and its regular use at

The dose may be repeated more than once if necessary, alternating with the enemata. The average intestinal tract contains an enormous amount of toxic matter, and after this has been got rid of it should not be permitted to reaccumulate. The turpentine and olive oil effectually prevent any accumulation. Under the influence of turpentine abdominal fullness is perceptibly reduced, and very soon the abdomen presents a hollow surface, especially in the region of the iliac fossae and hepatic and splenic flexures. It should be our aim to maintain this condition. With such physical signs we may be sure that bacillary activity and toxin manufacture is at its minimum, and this opinion is confirmed by the entire absence of unfavourable symptoms. Headache is never complained of after the second day; the tongue cleans early and remains moist, and patients, as a rule, express themselves as feeling quite well, their most serious trouble being an early desire for "something to eat."

It is very rare to meet with diarrhoea in the type of enteric occurring here. In an epidemic amongst troops stationed in St. Helena in 1902 constipation was the rule. At that time a simple enema of soap and water was given every other day; quinine, gr. ij, in solution was given by the mouth every three hours. Delirium was frequently present, and in over a hundred cases the mortality was about 10 per cent. I was then doing duty as a civil surgeon at the military hospital. Since I have used turpentine in the way described I have never once seen delirium in enteric fever, though formerly it was rather the rule than the exception amongst the island inhabitants. I have for many years invariably given quinine gr. ij or gr. iij, dissolved in hydrochloric acid, by the mouth, and believe firmly in its value both as a heart tonic and a bactericide. Amongst the last 30 cases treated as I have outlined there was one death.

The patient was a boy aged 17, one of four cases which came from the same house. He had a sharp haemorrhage the day before his admission to hospital, when I first saw him, and had been on ordinary diet up till that time. His abdomen was much distended, and though he improved markedly in the interval, he died some days after from perforation.

I regret that calomel—though in small doses—was given in this case, and I would not advocate its use when the disease is thought to have gone beyond the first week.

The two charts are (1) of a brother and (2) a sister of this boy. The girl stated that she had only felt ill for two days before admission. She was very drowsy, and had severe headache, but next day this had disappeared. This chart (2) shows clearly the beneficial influence of turpen-

Chart 1.—G. T., aged 20. Ill some days before admission. A, On admission.

Chart 2.—S. T., aged 22. Ill two days before admission.

stated intervals until the temperature has been normal for at least ten days. When the case first comes under observation, an enema of turpentine 3j and olive oil Oj is given by a funnel and tube; and, the foot of the bed being well raised, the emulsion is allowed to find its way slowly up the bowel. On the next day, or the day after, the same quantities are repeated. In most instances the amount of turpentine may then be reduced to 3ss with olive oil 3xv; this is administered on alternate days throughout the whole course of the disease, and continued until convalescence is fairly established.

A preliminary thorough cleansing of the bowel by a dose, or preferably small divided doses, of calomel, followed by castor oil, prepares the ground for the action of turpentine.

time in the course of the disease. The case promised from the initial condition to be a severe one; yet, though no sponging or antipyretics were used, the temperature came steadily down. Quinine was given in all cases, but in small doses is not an antipyretic.

The longest duration of fever was seventeen days; many cases touched normal in a week from the commencement of treatment, with no subsequent rise, and the average was about twelve days. There were no relapses, and alcohol was not indicated in any instance.

Vaccines may some day supersede other treatment. Until then, I recommend a well-known drug, used on a new plan, the adoption of which enables one to treat the disease with a confidence other methods fail to impart.