causes constitutional disturbance or death, but when septic infection has gained an entrance to it both frequently follow.

The uncomplicated primary "sore" of syphilis is not a sore at all, but a hard, dry, raised, painless lump.

Most of the pain, the odour, and the discharge of cancer are due to superadded sepsis. One of the conditions in which they are all pronounced is cancer of the uterine cervix. I have had opportunities for observing elderly spinsters with this disease. In all, the only early symptom was haemorrhage. Before the growth had caused serious disturbance it had in each instance invaded the bladder, was infiltrating the surrounding structures, and had become inoperable.

CHRONIC LYMPHATIC OEDEMA.

Amongst other causes of chronic lymphatic oedema of the extremities tubercle, syphilis, and cancer have to be remembered.

COMBINATIONS.

That of syphilis and tubercle is a deadly combination. I have seen cases in which tubercle attacked the lungs of a victim of secondary syphilis. In each instance the tubercle rapidly advanced to a fatal issue.

The most disastrous form of "scrofula" is a combination of congenital syphilis and tubercle.

Cancer and syphilis are very firm allies, and syphilis often provides a suitable site for the lodgement of cancer. If a person over 60 years of age contracts syphilis his death from cancer may be anticipated.

Tubercle and cancer favour the same sort of soil. Persons who have recovered from tubercle in their youth are exceptionally liable to cancer in their later years. It is more than a coincidence that both occur in families with a history of some members having attained to an extraordinarily long life.

THE ROENTGEN RAY APPEARANCES OF THORACIC ANEURYSM.*

By A. C. JORDAN, B.A., M.D.CANTAB.,

MEDICAL RADIOGRAPHER, GUY'S HOSPITAL, AND ROYAL HOSPITAL FOR DISEASES OF THE CHEST.

THORACIC aneurysm is a condition in which a Roentgen-ray examination is of the greatest value, for there are few cases in which such an examination does not clear up the diagnosis beyond possibility of doubt. In the normal thorax, viewed from the front or back, the normal aortic arch is usually entirely concealed by the shadow of the vertebral column. In a fair proportion of normal cases,

Fig. 1.—Diagrams of the aortic arch in the right anterior oblique view. 1, Normal arch. 2, Generally dilated arch. 3, Fusiform arch. c, Front of vertebral column. d, Diaphragm. Immediately in front of c is the posterior mediastinal clear space.

however, it forms a slight bulge to the left of the vertebral column below the clavicle. Thus, we must not fall into the error of taking a mere bulge to the left of the spine above the cardiac shadow as evidence of the presence of an aneurysm. It is necessary to supplement the view obtained in this plane by a view in the plane of the aortic arch itself. For this purpose the patient is placed vertically on a revolving seat in such a manner that he can be revolved upon his vertical axis between the x-ray tube and the fluorescent screen.

* A paper read in the Section of Medicine at the annual meeting of the British Medical Association, 1910.

It will be remembered that the direction of the course of the aortic arch from its origin is backwards and to the left. Hence we are able to obtain a view in the actual plane of the aortic arch by revolving the patient so as to bring the right shoulder forward. The exact angle to which he should be revolved in individual cases is determined by the appearance on the fluorescent screen. In this, the right anterior oblique view, the appearance of the normal aortic arch and of the aneurysmal arch is illustrated by the diagram (Fig. 1), which shows the appearance of the aortic arch in relation to the posterior mediastinum and to the vertebral column.

In both the normal and the generally dilated arch the sides of the aortic shadow are parallel, a line A drawn horizontally across the upper part of the shadow being equal in length to a horizontal line B across the base, where the aortic shadow and the shadow of the auricle meet. In the generally dilated arch, however, both A and B are greater in length than in the normal arch, and it is not uncommon to find them double the length of the

Fig. 2.—Large saccular aneurysm of the ascending portion of the aortic arch. The transverse portion of the arch is displaced to the left.

normal arch. In the fusiform arch the upper line A is longer than the line B at the base of the arch. A saccular aneurysm is merely an exaggeration of the fusiform arch.

The only form of aneurysm which is not revealed with certainty by either the anterior or the oblique examination is a small aneurysm confined to the concavity of the arch. I have no personal experience of this form except in association with general dilatation of the arch. If the symptoms, for example, those of pressure upon the recurrent laryngeal nerve, are too great in any particular case to be explained by moderate general dilatation of the arch, the fact should be borne in mind that there may be a localized saccular aneurysm in the concavity of the arch. In the case of a large saccular aneurysm of the aortic arch there can rarely be any doubt from the anterior view alone as to the diagnosis. When the saccular aneurysm affects the ascending portion of the arch, the transverse arch is usually displaced to the left by the pressure of the aneurysm, and is seen to the left of the vertebral column (Fig. 2).

In an aneurysm which involves the whole of the aortic arch it is usual to find the long axis of the heart far more oblique than usual. In explanation of this it should be remembered that a generally dilated arch is also elongated

Since the descending thoracic aorta is fixed in position against the vertebral column, it is the aortic arch, lying free in the thorax, whose position will alter. Consequently,

while a thin stream found its way through the oesophagus around the aneurysm.

In complicated cases the diagnosis may be far less easy.

Fig. 3.—Aneurysm of the innominate artery above a dilated aortic arch.

as the arch elongates, the commencement of the arch will take up a lower position in the chest, so much lower that in some cases the heart appears to be actually lying upon its right side with its long axis practically horizontal.

In less extreme cases, where the aortic arch shadow as seen from the front is decidedly greater than normal, the oblique view will show the exact nature and degree of the dilatation, and aortic arches which look very much alike from the front will be seen to be due in some cases to moderate general dilatation, in others to great general dilatation, and in others, again, to fusiform dilatation of the arch.

Aneurysm of the innominate artery usually forms a very striking and characteristic picture. In most cases the arch itself is dilated, sometimes considerably so. From the upper part of the shadow of the arch, just to the right of the spine, a considerable shadow ascends and is lost in the dense tissues of the neck (Fig. 3).

An unusual case is that of an aneurysm of the descending thoracic aorta, displacing the heart and aortic arch forward (Fig. 4). In the right anterior oblique view the outline of the heart and aortic arch are shown to be normal, but their distance from the vertebral column is increased. The posterior mediastinal clear space is occupied in its middle part by an opaque shadow, which was thought during life to be due to a mass of enlarged glands, but was shown at the *post-mortem* examination to be due to an aneurysm of the descending thoracic aorta. This aneurysm pressed upon the oesophagus and gave rise to difficulty in swallowing. A bismuth emulsion was found, when swallowed, to be obstructed at the level of the upper part of the aneurysm, the emulsion collected above it,

Fig. 5.—Aneurysmal dilatation of the descending thoracic aorta in its entire length.

Thus, in the presence of a large pleuritic effusion, where the lung above the effusion is compressed and the other lung is considerably congested, the outlines of the cardiac and aortic shadow are so far concealed that a diagnosis may be almost impossible, although, after the removal or the absorption of the pleuritic fluid, the diagnosis becomes obvious.

It is somewhat surprising to find how frequently an error in diagnosis may occur in cases in which the Roentgen-ray examination reveals the diagnosis at a glance. Thus phthisis is not infrequently diagnosed as aortic aneurysm, while an aortic aneurysm is still more often mistaken for pulmonary phthisis. New growth in the chest may give rise to areas of percussion dullness and to other signs and symptoms suggestive of aneurysm, while various swellings may, by receiving transmitted pulsation from the heart or aorta, simulate an aortic aneurysm very closely.

In other cases an examination of the aortic arch throws light on the nature of a case where the actual complaint is not connected with the thoracic aorta. Thus in a case of pulsating abdominal tumour examined recently, in which the diagnosis of aneurysm of the abdominal aorta had been made, although there was some possibility that the tumour was connected with the liver, it was shown that the aortic arch was dilated, while the descending thoracic aorta was uniformly dilated in its

Fig. 4.—Right anterior oblique view. A, Saccular aneurysm of descending thoracic aorta. B, Bismuth emulsion in the oesophagus above the aneurysm; a thin stream of the emulsion is seen passing thence down to the diaphragm, whence it enters the stomach. C, Aortic arch, normal in size, displaced forward by the pressure of the aneurysm. D, Diaphragm.

entire length. (Fig. 5.) Thus it may be said that the diagnosis of abdominal aneurysm was placed beyond a doubt, while, further, the utility of surgical treatment for it, in view of the condition of the thoracic aorta, was made clear.

Dilatation of the aortic arch usually follows atheromatous disease of the endothelium, and the question arises whether it be possible, by a Roentgen-ray examination, to determine the presence of atheroma while there is as yet no dilatation of the aortic arch. One is bound to admit that atheromatous patches as such are not recognizable either on the fluorescent screen or on a photographic plate, although calcareous patches have occasionally been shown when they happen to have been caught edgewise, but even this is of extremely rare occurrence.

There is one fact, however, which has considerable bearing on this question of atheroma, and that concerns the form of the cardiac and aortic shadows. We have shown that a generally dilated arch is elongated, and throws the heart into a more oblique position than is seen normally. When aortic regurgitation is present the form of the heart is characteristic, the apex being displaced to

Fig. 6.—Author's diagram showing the normal cardiac area (shaded) and the form of cardiac and aortic shadow commonly associated with aortic regurgitation due to atheroma of the aorta (curved black outline).

the left, while the left ventricle and the left auricle are both comparatively empty. In this way an excavated appearance of the left cardiac border obtains when viewed from the front, and this is accentuated by the prominence of the aortic arch to the left of the spine. Even before the aortic arch has begun to dilate this particular form of cardiac shadow is found, so that in any case in which the presence of an atheromatous aorta is suspected the discovery of this particular form of cardiac shadow affords valuable circumstantial evidence confirming the suspicion of atheroma (Fig. 6).

THE late Dr. Jameson John Macan, brother of the Master of University College, Oxford, left estate of the gross value of £7,628, with net personalty of £7,541.

ACCORDING to the *New York State Journal of Medicine*, the average annual income of the medical profession in the United States is 1,250 dollars (£250).

IT has been arranged that a special congress on the standard of purity in cocoa and chocolate will take place in Paris in January, 1911, when there will be present delegates from Germany, England, Austria Hungary, Belgium, Brazil, Spain, the United States, France, Holland, Italy, Mexico, and Switzerland. The congress is being arranged under the auspices of the White Cross Society of Geneva, whose delegate in the United Kingdom is Mr. Loudon M. Douglas, F.R.S.E., of 3, Lauder Road, Edinburgh.

MINER'S NYSTAGMUS.*

By H. S. ELWORTHY, F.R.C.S.,

INSPECTOR OF INJURED WORKMEN TO THE EBBW VALE COMPANY.

IN compliance with the request of one of your secretaries, who has done me the honour of asking me to read a paper before you, I propose to bring to your notice some observations on miner's nystagmus.

While engaged in examining injured workmen for the Ebbw Vale Company during the past three years a certain number of cases have been examined; but I must ask your indulgence if the observations are not very elaborate, owing to the limited time in which they have to be made, and also if the conclusions I have drawn from them may seem presumptuous or premature, considering the small number of cases and the short period of time in which I have been engaged in the work.

To begin with, miner's nystagmus is a disease peculiar to coal miners. I have made inquiries as to its occurrence in the Cornish tin mines, but have not heard of any cases. Nystagmus is not mentioned in the Blue Book on the Health of Cornish Miners, 1904. Inquiries as to its prevalence in the lead mines of the Isle of Man were also negative. For this information I am indebted to Drs. T. A. Woods of Douglas, and H. Gell of Peel.

ETIOLOGY.

On considering the question, "Why is it that nystagmus only occurs in coal mines?"—I come to the conclusion that it is the absence of colour that makes all the difference.

I take it that the cause is fatigue of the eye, resulting from working by artificial light with a black background and nearly black surroundings. There is the blackish "rubbish" or "slack" above and below the coal, and the jet-black coal with some sparkling facets which reflect the light. The monotony is not relieved by other colours. The miner may come to work with clean face and clothes, but in a few hours all is more or less black from the dust. Then he has to fix with his eye the point at which to strike with his mandril, or to lift out lumps of coal against a black background. When he turns his eyes away from the coal there is no colour relief. To make matters worse, the majority of miners prefer to work in the day. The consequence is that for the greater part of the year it is dark when they get up and go to work; they work in the dark, and in winter it is dark soon after they come out. They therefore get the minimum of daylight, and the opportunity of seeing a variety of colours by daylight, which means physiological relief to the eye.

Another cause of fatigue to the eye is when miners have to walk a mile or more underground to reach the "face" where they work. They proceed more or less in single file, each carrying a lamp which gives out light in every direction. The result is that the miner walking behind another gets the glare of that lamp in his eyes all the time. Consider what that means in a two or four mile walk.

Then, after working his eight hours and his eyes being more or less exhausted, he has to make a similar journey back. On inquiry I have generally found that men feel the effects of nystagmus most when they come from work with the abominable glare of the lamp just in front of them. Many have to sit down and wait until it has gone out of the way.

As to the theory that nystagmus is produced by working while lying down I can give no opinion, as the great majority of the Ebbw Vale colliers, I am told, work in the upright position; and I have not yet seen a case of oblique nystagmus. And, besides, other underground workers get nystagmus, such as hauliers, timbermen, and repairers in coal mines. The only class of workers in coal mines that seem immune is the ostlers, and that is because the stables are whitewashed. That it is not due to imperfect fusion is shown by its occurrence in one-eyed men.

The one essential factor seems to be exhaustion of the eyes from working in the general blackness without any colour relief.

* Read before the Monmouthshire Division of the British Medical Association.