THE ROUTINE TREATMENT OF MALARIA IN UGANDA.

BY

J. A. TAYLOR, M.B.EDIN., D.T.M.LIVERPOOL,
MEDICAL OFFICER, UGANDA PROTECTORATE.

AFTER more than ten years' experience of the treatment of many thousands of cases of malaria in Uganda, the results I have obtained with small doses of quinine have led me to conclusions so different from those expressed recently in your columns that I hope they may be worth recording.

Most of the cases of malaria met with in Uganda are simple subtertian without complications, and the routine treatment I adopted for these was as follows:

1. Calomel, in an average dose for an adult of not less than 5 grains, followed in some hours by a saline. This usually meant that the calomel was given last thing at night, and the salts at 6 a.m., or when the patient awoke the next morning; but if there were any signs of the calomel acting during the night the salts were given at once.

2. Quinine, 5 grains of the hydrochloride by mouth one hour after the saline. This was usually about 7 a.m., and between this time and noon three more doses of 5 grains each were given at more or less regular intervals—for example, 8.30 a.m., 10 a.m., and noon. On subsequent days, until the temperature had remained normal for twenty-four hours, 20 grains were given daily in the same manner, the first dose at 6 a.m., or as soon as the patient was awake, and the last dose at noon. After the temperature had been normal for twenty-four hours the quinine was reduced to 15 grains daily—5 grains before the morning, mid-day, and evening meals. This was continued for one week and then reduced to 10 grains daily—5 grains morning and evening. Ten grains daily were continued for a fortnight, and then only 5 grains each evening for two months.

3. Phenacetin in 5-grain doses up to 15 grains daily for headache alone, or aspirin in the same doses if complaint were also made of pains in the limbs and body.

Under this treatment the temperature usually remained normal after the third day, but occasionally, especially if a first attack, after four or five days; no complications occurred in any of my cases from the effects of the quinine or malaria, there were no recurrences of the fever during the three months' treatment, and attacks only occurred later when some definite fresh infection had taken place.

In the most typical cases all the quinine was given before noon while the temperature was little if at all above normal, and there was thus better absorption and little likelihood of vomiting. In the less typical but not uncommon cases in which the temperature had not fallen in the morning, my practice was to give only one or two of the 5-grain doses of quinine at three or four hours' interval in the morning and the remainder of the 20 grains during the afternoon and evening. If the quinine was vomited or the temperature rose above 104° F. before at least 15 grains of the quinine had been taken, an intramuscular injection was usually given. Intramuscular injections, however, were required in less than 1 per cent. of my cases, and usually on not more than one occasion. If the patient made no objection I gave the quinine in solution, but I cannot say I found this more satisfactory than in the compressed form. A very great objection on the part of the patient to taking quinine in solution—and the majority of one's patients have a very great objection—in my experience reacts on the stomach, the absorption is retarded instead of hastened, and vomiting sometimes ensues, whereas tablets would be retained and absorbed.

My remarks apply to subtertian malaria. I have met with no cases of pure quartan. Pure tertian was also uncommon in Uganda before the war, but during the war there was a considerable increase in the number of cases, no doubt brought about chiefly by the troops from India. In tertian I have carried out the same treatment as described for subtertian, but have waited until the temperature has remained normal for two clear days before reducing the dose to 15 grains.

With larger doses of quinine daily I have been unable to obtain results as satisfactory. Nor have I found the method of giving 10 grains of quinine two or three times

a day satisfactory. By the time the evening, and sometimes the mid-day, dose is given, the temperature may have already risen and the quinine may be vomited or only partly absorbed. Also, after an evening dose of 10 grains, the headache and general discomfort are increased, with a restless and sleepless night as a result. Even an intra-muscular injection when the temperature is high does not produce the same effect as a smaller dose by mouth when the temperature is low.

I do not think too much emphasis can be laid on the importance of giving the calomel is not less than a 5-grain dose, both in malaria and blackwater fever. The one or two grains sometimes given usually causes severe pain, griping and numerous visits to stool with slow and little result, whereas 5 grains causes little discomfort and acts only once or twice with very good and quick result.

As compared with other purgatives, a dose of calomel (5 grains) at the commencement of an attack of malaria appears to have an almost specific action in preparing the way for the quinine, and to it I attribute to a very large extent the absence of complications and quick response to quinine.

With blackwater fever my experience is that if the 5 grains of calomel can be given immediately an attack begins no serious complications arise, and, with careful general treatment, blackwater fever is removed from the category of the most dangerous diseases. I have had only two deaths among my patients from blackwater fever. In each case I was able to see the patient for the first time only on the third day of the attack; one patient had taken 2 grains of calomel, and had been on and off the stool all night with great pain and discomfort, but with so little result that several soap and water enemata were needed before a saline injection could be made into the rectum. In the second fatal case no purgative at all had been given. Blackwater fever, however, is, in my opinion, a complication of malaria which should rarely if ever occur. It appears to be the result of a final and drastic effort on the part of nature to eliminate the parasites from the body, usually after several untreated or insufficiently treated attacks of malaria. I have seen no case of blackwater fever occur during the treatment of malaria if both calomel and quinine had been given, but I have seen cases, usually slight, after quinine alone. In these, to my mind, the question has not been whether the quinine brought on the attack, but how much more serious the attack would have been if no quinine had been given.

With regard to large doses of quinine it may be of interest to record that during the war several cases of malaria were sent up to Uganda from Mombasa to try the effects of a change of air. Some of these whom I saw at Entebbe informed me that they had been taking 40 to 60 grains of quinine daily for several weeks without the temperature remaining normal for twenty-four hours. With most of these complete stoppage of the quinine brought the temperature to normal within three days. In one case the temperature came to normal and remained normal until the eighth day, when it again went up to 103° F. Subtertian parasites were then found in the blood, and treatment with calomel and quinine as described above was commenced. In three days the temperature was again normal, and there had been no recurrence of the fever when I last heard of the patient three weeks later.

I have had no experience of the treatment of malaria elsewhere than in the Uganda Protectorate. It is possible, although I have not seen this point discussed, that the effect of the altitude of Uganda—more than 3,000 feet above the sea level—upon the blood may help the action of quinine. In Uganda, also, I usually had to deal with only single infections, and earlier treatment was perhaps more often possible. Whatever may be the explanation there would appear to be a great difference between the doses and manner of administration required in other localities as compared with Uganda.

I cannot help thinking that too much quinine, especially too much in a single dose, can be given as well as too little. One grain of calomel given to many Uganda natives will cause intense salivation and other alarming symptoms of mercurial poisoning, while a larger dose given to the same person may have only a purgative action. Some analogous action may occur in the case of quinine. It is possible that too large a dose may, in some as yet undetermined manner, prevent the necessary action.

When giving more than five grains of quinine daily

I have always used the hydrochloride if obtainable, but if only five grains daily, the cheaper salts are just as satisfactory.

As soon as the patient can get about, some combination of arsenic in increasing doses and iron in small doses has proved most useful.

A COMPARISON OF TWO METHODS OF ADMINISTERING ARSENO-BENZOL COMPOUNDS IN SYPHILIS.

BY
H. E. GIBSON, M.D.Oxon.,
LONDON.

A COMPARISON of the two following courses of antisyphilitic treatment may be of interest.

In the first series of cases treatment consisted of one dose of 0.3 gram of kharsivan, followed at weekly intervals by doses of 0.6 gram, until a total of 2.7 grams had been given, but an interval of fourteen days was allowed between the third and fourth doses. This course was thus completed in five or six weeks. For want of a better term it may be termed a "concentrated course" of treatment. If further treatment was necessary owing to the Wassermann reaction being still positive, potassium iodide was given for fourteen days, and then two more injections of 0.6 gram, with one week's interval between them. All the above was given intravenously by the gravity method.

In the second series of cases, which may be termed a "prolonged course" of treatment, three injections of 0.3 gram were given, then two of 0.4 gram, and two of 0.5 gram, all at weekly intervals, except for a fortnight's rest between the third and fourth injections, as in the other series. This course lasted about eight weeks, and the drug was given intravenously as in the "concentrated course." Further treatment, if necessary, consisted as a rule of intramuscular injections of nov-arsenobillon, at weekly intervals, generally one of 0.3 gram and one of 0.6 gram. Occasionally there were slight variations in the above prolonged course, as, for example, by substituting a dose of 0.5 gram for the second dose of 0.4, or a dose of 0.6 for the last 0.5 gram.

The criterion of cure was a negative Wassermann reaction.

The total amount of arseno-benzol compounds given to any one case was limited to 4 grams, the first course, at the end of which the Wassermann reaction was tested, consisting of a total of 2.6 to 2.8 grams; the second, or "follow up" course, before a second Wassermann test was made, bringing the total to an average of 3.9 grams in the concentrated course, and 3.6 in the prolonged series.

It will be seen from the table that, although the percentage of primary cases which showed a negative Wassermann reaction after the first course is slightly higher in the "concentrated" series, the results are completely changed when we consider the cases showing secondary symptoms.

Cases Treated with a Concentrated Course.

No. of Cases.	Negative with up to 3 grams.	Negative with up to 4 grams.	Positive after 4 grams.	Doubtful after 4 grams.
Total cases... 89	47=52.80%	14=15.73%	23=25.84%	5=5.61%
Primary ... 34	28=82.35%	3=8.82%	2=5.88%	1=2.94%
Secondary ... 52	17=32.69%	10=19.23%	21=40.32%	4=7.68%
No symptoms ... 1	0	1=100%	—	—
Previous treatment 2	2=100%	—	—	—

Cases Treated with a Prolonged Course.

No. of Cases.	Negative under 3 grams.	Negative under 4 grams.	Positive after 4 grams.
Total cases ... 100	68=68%	38=38%	4=4%
Primary ... 65	52=80%	12=18.46%	1=1.53%
Secondary ... 35	16=45.71%	16=45.71%	3=8.57%

All cases also received 1 grain of mercury, given intramuscularly, at weekly intervals.

Apart from the question of cure, one great disadvantage of the short concentrated course is that the reactions are much more pronounced than in the prolonged course. These include diarrhoea, vomiting, and sometimes a violent and immediate vaso-dilator effect, in which the patient's face becomes "blown up," and the pulse weak and irregular; this is most alarming when seen for the first time, but apparently it always passes off without active treatment. Opinions differ as to the benefits of adrenalin in this condition.

Although slight reactions, such as diarrhoea, vomiting, and headache, may occur with the prolonged course, they are not so marked as in the other, and I have so far not had a single instance of the vaso-dilator phenomenon.

It has been impossible to follow up these cases to see what percentage of relapses, if any, occurred in the two series.

Conclusions.

Advantage seems to lie with the prolonged course, partly owing to the lesser incidence and violence of reactions, and partly because the total results are better than with the concentrated course; this especially applies to secondary cases.

Should a patient showing only a primary sore have urgent reasons for wanting a short course, the concentrated course may be given, but otherwise it is better to spread the treatment out over a longer period.

Memoranda:

MEDICAL, SURGICAL, OBSTETRICAL.

TREATMENT OF DELIRIUM TREMENS.

DR. STARKEY has sounded a note of warning, in the *BRITISH MEDICAL JOURNAL* of January 10th, with regard to Dr. Wyatt-Smith's remarks, under the above heading, in the *JOURNAL* of December 6th, 1919, which I must support. Prior to the war I made extensive observations on the action of various hypnotic drugs with a view to forming some ideas as to their selection in different types of mental disease. My results were published in the *Journal of Mental Science* (July, 1914), but I feel that it may be of some interest if I record here some of my observations on the drugs mentioned.

Hyoscine.—My opinion is that this drug is a very powerful cardiac depressant and that it ought not to be prescribed in patients who suffer from "heart disease." I laid considerable emphasis on this in my article referred to above and expressed the view that it should not be given to anyone who was not in robust physical condition. In states of extreme maniacal excitement, in strong healthy people, however, when one wishes to obtain the influence of a hypnotic as quickly as possible, it is a most useful drug, and given by hypodermic injection it is very rapid in its action. I have frequently given doses of gr. $\frac{1}{10}$ with benefit; in fact, large doses must be given in these cases and in this way the repetition may be avoided. I have found it especially useful in cases of delirium tremens, as Dr. Wyatt-Smith states, and especially so if combined with atropine and morphine. In milder cases it is of some use in doses of gr. $\frac{1}{10}$ three times a day, but I prefer not to use it for repeated administration, and the warning given by Dr. Starkey is fully justified from the experience I have had of the drug.

Sulphonal.—Most of those who have done service in mental hospitals have considerable experience of the use of this drug, and although Dr. Wyatt-Smith states that he has not yet met anyone who had seen a case of sulphonal poisoning, I must add my name to the list, with Dr. Starkey, of those having had this experience. I have, however, only seen one case in fourteen years' asylum work. It is usually associated with vomiting, diarrhoea, and abdominal pains, and the patient slowly passes into a state of collapse. The warning signal is haematoporphyrinuria, and it seems to me, therefore, of the greatest importance when prescribing this drug to have careful observations made on the passage of urine, and the report of any tendency for it to change to a reddish tinge. It has been my invariable rule to have these instructions written on the box in which sulphonal is dispensed, and this may account for my having seen only one case of poisoning from the drug, but I realize the