PSYCHOPATHOLOGY AND DISSOCIATION.*

BY

WILLIAM BROWN, M.A., M.D. OXON., D.Sc. LOND.,

READER IN PSYCHOLOGY, UNIVERSITY OF LONDON (KING'S COLLEGE); LATE MEDICAL OFFICER IN CHARGE OF CRAIGLOCKHART WAR HOSPITAL FOR NEURASTHENIC OFFICERS.

I HAVE chosen this subject because it seems to me an especially interesting part of psychology, and to those who are beginning the subject it opens out prospects of advance in the science which are extremely attractive and stimulating.

Psychopathology is the science of the abnormal or pathological working of the mind. Psychology considers the laws of working of the normal mind, the mind that is in adequate adaptation to its environment, physical and mental. Psychopathology considers the laws of working of the mind that is out of harmony with its environment because of some degenerative process.

The difficulty is to find one guiding idea to carry us along in this very complicated subject. The hint comes from normal psychology. Not many years ago our own English psychologists developed a system of association psychology, explaining the mind in terms of association of ideas—that is, regarding the mind as something gradually built up on the basis of what is given through impressions from without and of what is inherited within. So in psychopathology we might begin our discussion by considering facts of dissociation—indeed, that is the word to conjure with in our subject—dissociation, splitting up of the mind, the antithesis of association. We shall take as examples of pathological states the so-called borderline cases—hysteria, neurasthenia, psychasthenia, and compulsion neurosis. If we go beyond these we come to what are called the psychoses, which are recognized forms of insanity. Much the same kind of explanation will cover all types of mental disease, but the attempt to produce a satisfactory science of mental dissociation was begun within the domain of borderline cases, otherwise known as the psychoneuroses. These diseases may also be called functional nervous diseases. In them there is no demonstrable lesion of the nervous system. The nervous system is apparently intact, but is functioning wrongly or inadequately.

This lecture will be devoted to the problem of hysteria.

Hysteria.

In hysteria we have very obviously a case of dissociation; certain parts of the mind, certain sensations, certain powers of movement, certain memories, are lost to the main consciousness, but still exist somewhere or other, in some form or other, and can be restored under appropriate conditions. The anaesthesias that hysterical patients suffer from are a good instance of the kind of dissociation that occurs. A hysterical patient may suffer from anaesthesia of the hand, or of the hand and lower arm, or of the hand and arm up to the shoulder, ending in a straight line round the shoulder, or, again, from hemianaesthesia extending down the right or left side of the body and ending accurately in the middle line. Many other forms of anaesthesia are possible. All these types correspond in their extent to ordinary, uneducated concepts of the divisions of the body; for example, the hand, in ordinary unscientific thought, is considered as one entity, the whole arm is considered as an entity, ending in a straight line round the shoulder. This holds good even for purposes of tactile association, but from anatomy we know that the nerves supply irregular areas, consisting, for example, of a strip down the forearm, extending towards the fingers of the hand, so that, if there were an organic disturbance of a cutaneous nerve, we would find an irregular area of anaesthesia—the more irregular the smaller the number of nerves affected—we would not find an area of anaesthesia corresponding to the hand, or the arm. Pierre Janet emphasized these distinctions between what are called functional anaesthesias and organic anaesthesias, and pointed out that the anaesthesia illustrates the state of mind of the hysteric, who has lost the power of holding together in one unity all the sensations

from the different parts of his body. A patient may receive a blow on the arm, causing a mental shock, as a result of which his whole arm may become anaesthetic, and the anaesthetic area may end in a straight line round the shoulder-joint. That area does not correspond to the distribution of any set of nerve fibres, but it does correspond with the patient's own conception of his cutaneous sensibility. Moreover, it can be proved in directly that the sensations in this part of the body still exist. Sometimes we can do this by a trick, as Janet pointed out: We can ask the patient to say "yes" when he feels a pin-prick, "no" when he does not. We get him to close his eyes, and then silently explore different parts of his body. Whenever he is pricked on the normal part of the skin he will say "yes." When he is pricked on the so-called anaesthetic area of the skin he will often say "no," whereas if the area were really anaesthetic he would not know that it had been touched at all. Yet he is probably quite genuine in his belief that he has not felt the pin-prick—though of course we must be certain that he is not malingering. It is very important in considering hysteria to be familiar with the various types of malingering. Cases like this, of anaesthesia combined with functional paralysis of the arm, occurred frequently during the late war.

I have dwelt on hysterical anaesthesia, not because it is very important, but because it well illustrates this theory of functional dissociation. There is dissociated from the entire mind one psycho-physical power, in this particular case cutaneous sensitivity—the power of reacting to stimuli presented to some part of the body. Again, the patient may have loss of motor power: he may be suffering from loss of power in the legs—cannot stand or walk; yet if we investigate his nervous system we may find that it is perfectly normal. Nevertheless, there is loss of sensory power, or loss of motor power, and also, very frequently, a greater or less degree of amnesia, or loss of memory. To sum up the state of mind of the hysterical individual, we may say that it is a state of mental dissociation—dissociation of certain psycho-physical functions from the main personality. Yet these dissociated functions remain, and under appropriate conditions can be recovered—for example, under hypnosis. Hysterical patients are easily hypnotized; in fact, Charcot and Janet hold that any person who is easily hypnotizable is *ipso facto* hysterical, and that, if an ordinary person be hypnotized, he is made hysterical by dissociating his mind—a form of dissociation has been started. With this view I agree.

It is worth while considering this matter a little further. To hypnotize a person we get him to fix a bright object with his eyes, to relax his muscles, and to give his mind up to sleep—to turn his mind away from everything active. We continue to talk to him all the time, telling him to think of sleep, and saying that his eyelids will begin to get heavy, that he will lose feeling in his limbs, and that he will gradually become more and more drowsy. After he has fixed the bright object for a short time and his expectation of sleep has been encouraged in this way and his thoughts fixed on sleep, which (though we have as yet no satisfactory theory of sleep) may be provisionally said to be a quieting down of the mental powers, he will appear to lose consciousness. But there is one power of the mind that is not quieted down, and that is the power of hearing the physician's voice. The physician is talking almost the whole time, and whenever he speaks the patient, until he gets very deeply hypnotized, can respond. Thus we produce dissociation of the power of hearing a particular kind of sound from the rest of the mind. The rest of the mind is lulled to sleep; it is a case of partial sleep. But more than that, there is an emotional element at work. A patient cannot be hypnotized by a gramophone, unless he has been hypnotized before and has an extreme tendency to yield to this kind of suggestion. The patient has to feel an interest in the physician, either of fear or of confidence. Those are the two forms of mental reaction. Different people hypnotize in different ways; some have a tendency to hypnotize through fear, others through arousing confidence—one might even say affection—in the patient. There is always an emotional element involved. We shall have later to consider a theory that makes a great deal of this emotional element. In the case of a normal person such an emotional appeal must be very strong; in the case of a hysterical person it need not be so strong, because the patient already shows

* A lecture delivered to Psychology students at King's College, Strand, December 15th, 1919.

a tendency towards dissociation; the hypnotic effects appear more readily. Charcot considered the hypnotic state a form of artificially induced hysteria. Janet agreed with him, and so do I, though many people think that, because almost everyone can be hypnotized to some degree, this theory must be wrong. But there is a tendency to mental dissociation in all of us, based on emotional conflicts. No one is a complete mental unity.

Finally, as an example of a hysterical patient, let me take the condition of almost any bad case of shell shock on the Western front, or any of the other fronts, during the past five years—any of the cases as they reached the casualty clearing station. The patient's condition was always that of dissociation. All such cases agree in showing loss of memory of greater or less extent. They may have forgotten everything that has happened since the shell burst, and they will exhibit other losses of function, greater or less, according to circumstances. They may show loss of voice, of powers of walking, of powers of hearing, of the power of voluntary control. I would emphasize their loss of memory. It will be found that such a patient is very easily hypnotized; if it be suggested to him that he will remember the circumstances of his injury with hallucinatory vividness, he will act again the whole circumstances, and in that process his various dissociated functions will return. If he has been dumb, he will now speak, without the necessity of suggesting this. If he has been paralysed, movement will occur in his limbs. I do not say that he will at once be able to take up his bed and walk, but his limbs will move about, showing that there is power in them, and that the power is linked up with the lost memory. What has been done here? We have re-associated him by bringing up these lost memories. With the memories we have brought up the lost functions. But we have done more than that, we have given him an outlet for an emotion which was originally experienced by him with too great intensity—with so great an intensity that he could not do full justice to it, and his mind split in the attempt. The way in which these patients live again through their experiences shows what terrible sensations they must have had. They roll about, gripping at the sides of the stretcher, or rolling on the floor, tearing at their hair with their hands, contorting themselves in every possible way, foaming at the mouth, becoming purple in the face, their eyes starting out of their head, all their muscles tense. While under shell, the conscious personality was trying to suppress the emotion of fear, partly from a sense of duty—the duty of a soldier—partly in self-defence—anything to get away from it. There occurs a mental conflict—an attempt to get rid of the painful emotion. The attempt succeeds at the expense of the mental machinery; clouding of consciousness supervenes, the patient goes into a state of stupor, where he is not completely unconscious, but where he is dazed. He gradually recovers from this, but has no memory of it later. We see here the mechanism at work explaining the dissociation. The dissociation is obviously a fact, but it is not an ultimate fact. There is a cause for it, and that cause is mental conflict, an attempt at repression, an attempt which has succeeded to a certain extent, but only at the cost of the production of a pathological symptom or set of symptoms. In curing the patient we bring up the repressed experience once more, we encourage him to work off the emotion involved in it. Just as a person who grieves for someone he has lost finds relief in tears, so we let these patients work off their fear. Thus we are concerned not only with the linking up of the dissociation, but also with the working off of fear, of a certain amount of emotional energy that has been bottled up in the patient.

This illustrates the advance that Sigmund Freud made on Janet in explaining hysteria. Janet explains hysteria simply in terms of dissociation. He says that the patient suffers from weakness of mental synthesis. He compares the hysteric to an old lady who has gone shopping and has made too many purchases. She comes back with her arms laden with parcels. Some of these fall to the ground; she stoops to pick them up, and others fall. So, he says, is it with the hysteric after a shock. He loses the power of holding all his psycho-physical functions simultaneously in his mind so that some of them are lost, and he suffers from an anaesthesia and paraplegia. If you cure him of this by suggestion, later on he will, perhaps, as the result of some slight shock, show some other

functional symptom—he may lose his voice or his hearing. Some other hysterical symptom will take the place of the original one unless he is cured of his general mental state. But Janet did not undertake to explain why the dissociation took place. Freud found, after an investigation of some hysterical cases under hypnosis, that there was very frequently loss of memory, that such memories were always of an emotional character, and that the symptoms disappeared if the memories were brought to the surface of the mind with their original vividness. Breuer and Freud published their first article on the mechanism of hysteria in illustration of this process—this effect of repression of painful emotional experience: loss of memory and dissociation, and the recovery of these by means of hypnosis and the dissipation of the symptoms themselves through encouraging the patient to live through the emotion again. This living again through the emotion and working it off was called "abreaction." At this early stage of his theory, then, Freud regarded dissociation as due to a preliminary mental conflict and a repression of one of the conflicting mental tendencies. Later on he developed this view further. He found that in the case of the dreams of normal persons there was also evidence of mental repression, and of the production of mental symptoms as a result of the repression. His view of the dreams of normal persons was, and is, that they correspond exactly with the symptoms of the hysteric, that they are "disguised fulfilments of repressed wishes," to use his own words. I do not want to go into his theory of dreams in detail, it would take us too far afield; I wish only to recall this part of his theory.* Later on he found, after analysing a number of his own cases, that the emotional tendencies which were repressed were all of a sexual nature, and so he enunciated his dictum that no neurosis is possible in a normal sexual life, and that, if a patient is suffering from neurosis, there must have been some disturbance in his sexual life.

That is the extreme form of the Freudian theory, which I do not accept, though the Freudian school make it an essential part of their doctrine. Freud thinks that we can find evidence of sexuality in the early years of childhood—namely, feelings of attraction of the boy towards his mother and of the girl towards her father, combined with hatred and jealousy felt towards the parent of the same sex—and that these feelings are shortly afterwards repressed. A mental conflict takes place in the child's mind, and sexual repression occurs, partly through the further normal development of the child's mind—other interests arise and crowd out these feelings—partly through the fact of development of feelings of sympathy and of morality, and these repressed tendencies only produce a competing factor if shocks occur later in the child's life.

I have already referred to the production of hysterical symptoms as being due to an emotional shock. That seems to be the general rule—an emotional shock will produce hysteria in persons whose inherited make-up is suited for it. The hysteric is born, not made, but the hysterical symptom is originated through some emotional shock. Freud considered that these emotional shocks were repressed and dissociated from the mind, and continued to act as a foreign body in consequence, but that these repressed memories could be brought to the surface either by hypnotism or by his method of psycho-analysis. In his earlier work he thought that he could prove in every case the actual occurrence of a shock in early childhood of a sexual nature. Later on, however, he discovered that some of the memories brought up by these methods were false memories—that the patient had imagined them—but this did not alter his theory very much. He said that the fact that the patient had the power of imagining such events showed that there was maldevelopment of the sexual life at that time, so that he then explained the neurosis in terms of the process of development of the psycho-sexual life of the child—a theory into which I cannot go now. To be a thoroughgoing Freudian now one must be able to accept this theory, and for many of us it is because we cannot subscribe to it that we should not call ourselves Freudians.

C. G. Jung of Zurich was Freud's most brilliant pupil, and at first an enthusiastic supporter of his views. Later, however, Jung's researches led him to believe that these

* See my articles on "Freud's Theory of Dreams," *Lancet*, April 19th and April 26th, 1913.

earlier emotional memories of the child were not an essential factor in the production of neurosis, but that the principal factor was the present condition of the patient. When a patient falls ill of a psycho-neurosis it is because he is not adequately adapted to his present social and physical environment. Life is too great a task for him; he cannot hold his own; he has not sufficient courage to face facts, with the result that, if he is of a hysterical temperament, his mind turns back to childhood's memories and to more childish modes of conscious activity. What Jung calls "regression" takes place. The mental energy of the individual, which Jung has called the "libido" (corresponding very closely to Bergson's "*élan vital*"), is reflected back, and revives earlier memories of childhood, not only memories of actual occurrences in childhood, but also early fancies. These fancies may be of a sexual nature or they may not. The libido in that way becomes linked up with earlier memories, and is no longer of use to the individual in his present mental situation—thus he becomes still less competent to deal with his environment.

But Jung would use the same method as Freud uses in treating the patient—the method of psycho-analysis, because, he says, we want to get back these earlier memories for the sake of the libido that is linked up with them. Psycho-analysis is simply the method of free association. If the patient is suffering from some hysterical symptom you get him to let his mind pass into a passive state—to recline on a couch, say, with his eyes closed—and passively follow up the sequence of ideas as they occur to him one after another, without any prompting. He begins, say, to think of the time when the symptom first developed, and then just watches his mind drift as it will—gets into a state of meditation—and tells the doctor from moment to moment what he is thinking about without reservation or criticism. His thinking seems to be undirected, to be just determined by free association. The result is that more and more ideas and memories from the past come to his mind, and perhaps eventually memories that are of great significance for his symptom. This occurs because the method is not really a method of free association at all. The series of ideas is, it is true, not determined by conscious mental process; it is determined by unconscious mental process. By letting the conscious selective activity fall into abeyance trends of unconscious mental activity come nearer to the surface of the mind and guide the ideas, and these belong to the same world as those mental tendencies which are at the basis of the hysterical symptom.

The difference between Freud and Jung, then, is, that Freud says that earlier memories and earlier imaginations are directly responsible for the hysterical symptom; Jung says that failure of adaptation at the present moment causes a regression or reflection of mental energy back to the past, and that this revives earlier imaginations and memories and so produces the hysterical symptom. Both agree in adopting the method of psycho-analysis, which proceeds to recall these earlier memories and fancies.

These, briefly, seem to be the fundamental ideas of Freud and Jung on hysteria, as contrasted with the ideas of Janet. They go a step further to the cause of the dissociation—that is, mental conflict and repression. Link up this dissociation and remove this repression and the patient is cured.

Freud, who originally used hypnosis, gave it up, partly because he found that he could only hypnotize about one-third of his patients, whereas psycho-analysis, though it took much longer, could be applied to every case, and produced eventually much the same results. His second objection to hypnosis was that certain resistances are overcome by the recall of particular memories, but in this process the resistance in other directions is increased, so that the mind is limited rather than expanded. He also suggested the germs of the theory that Ferenczi worked out in detail afterwards—namely, that in hypnosis there is really a transference of emotional feeling, of the early sexual feelings of childhood, to the person of the physician. Ferenczi points to the fact that there are two main methods of hypnotizing: the method of command, corresponding to the paternal method with the child, and the method of coaxing and soothing, which corresponds to the attitude the mother adopts towards the child. He then argues that in both types of hypnosis there is a revival in the patient's mind of its early modes of response. When a patient is hypnotized he responds to the physician just

as he responded to his father or his mother in early years—that is, by a reaction that is of a sexual nature—hence the patient's early sexual feelings are fixed on the physician. What has thus really been done is to replace one set of symptoms by another. Hysterical symptoms, like loss of the power of walking or loss of voice, may thus be got rid of, but in the place of these the psycho-sexual dependence of the patient upon oneself has been produced, and this also is bad. What evidence has Ferenczi for this view? He says that it is taken from actual experience of patients whom he originally treated by hypnosis, but later on psycho-analysed. He found evidence of such transference of feeling in them.

These, then, are Freud's three objections to the method of hypnosis: (1) Failure of the method in certain cases. (2) Its tendency to produce other symptoms in place of those of which the patient is cured. (3) Fear of the transference of sexual feeling to the person of the physician.

With regard to the last objection we might urge that Ferenczi's theory is a *petitio principii*; the patient may respond to the physician as he did to his father or his mother in childhood, not necessarily, however, because of sexual feelings, but perhaps simply because at that time he was more suggestible than in later life. A young child has only a few facts in his mind, and if any suggestion is made to him he tends to believe it unquestioningly. He comes into the world with a tendency to what is called "primitive credulity"—a very important attribute. Most of what he learns in early life is learnt through suggestion, not by means of logical reasoning. We like to encourage a child to reason, but if we made it think out everything for itself we should only confuse it and hamper its development, instead of forwarding it. And if that is true of the intellectual side of the child's life, it is much more important on the moral side. We can explain the child's response simply in terms of greater suggestibility. We get our patient into a state in which he is willing to accept the suggestions that are made to him in the same way as he accepted them in his early childhood.

This method is much quicker than that of psycho-analysis, and can produce equally satisfactory results in simple cases. I will give two illustrations.

One is that of a signaller in the Flying Corps who was blown up by an aeroplane bomb whilst taking refuge in a disused trench in France. He became unconscious, and on coming to be found that the trench seemed to be turned round at right angles to the position in which he had expected it to be. When he arrived at the barracks the same thing had occurred—everything seemed to be twisted at right angles to its ordinary position. This feeling of disorientation, as we may call it, persisted for many months. When he came home to England it still showed itself in the following ways: When walking along a street which bent roughly in the form of a semicircle, thus $\cup$, he would have the irresistible feeling that he had been walking like this $\cap$. If he were riding on the top of a bus, and the bus turned at a right angle, he would have the feeling that he was continuing to go in a straight line, although he knew that the bus had turned a corner. He would think that the whole room he was in, everything, had twisted round to a right angle. I hypnotized this patient and put him through his experiences again, with the result that the occasions on which he got this feeling of disorientation became roughly about half as frequent as before; but the disorientation continued to occur from time to time. He was better, but by no means cured. A few days later he came to tell me that his mother had informed him that at the age of six, as the result of a fall, he had shown much the same symptom; while crossing the Tower Bridge one day he had told her that the bridge was turned "the other way round." I hypnotized him, and put him through this experience again. He lived it through with much vividness. He is sitting on a wooden horse (white with red stripes) in the dining-room. His aunt comes into the room and wishes to wash him; he declines, and edges away from her; the horse tips up on the edge of the hearthrug, and he falls on to the fender. He knows no more until he wakes up and finds himself in bed, with his aunt bathing his face. He continues to recall his

* But all Freudians regard such "transference" as an essential stage in the process of cure, and expect to produce it, in every case, in the course of psycho-analysis. They subsequently "resolve" the transference by further analysis.

experiences, and remembers crossing the Tower Bridge and finding it turned at right angles. After recalling this incident the patient was much better, but a few days later he went to the theatre to see *Going Up* (it is true that there is an aeroplane in *Going Up*), and he suddenly found that the theatre was the wrong way round. He went home, and in the middle of the night he woke up with the feeling of disorientation, got out of bed, and went towards what he thought was the window, and put his hand through the looking-glass. Thus he was by no means cured. I pointed out that there must be some memory in the first six years of his life which was the cause of the trouble. He tried to recall it, but did not make much progress. I then put him through his fourth birthday under hypnosis, and he remembered a frightening dream in which he had thought that there was an animal under the table. He had crawled downstairs and waked up half-way down. The next morning he had thought that the pier was the wrong way round, and had told his mother so. I interpreted his dream for him while he was in the hypnotic trance. He had seen the Giant Ape in a glass case at the Natural History Museum, South Kensington. This had frightened him, and it was this creature which he had thought was under the table in his dream. I took the precaution to get another doctor (Captain P. A. Galpin, R.A.M.C.) to come and see me hypnotize this patient the next time. I then suggested to him that he would go through his first experience of disorientation. When I put my hand on his forehead he shouted out, "Hot coffee!" It appeared that, when a child of nearly three, he had gone into the kitchen one morning, and had pulled the coffee-pot, which was on the table, towards him by pulling at the tablecloth. The coffee upset and poured down his right arm. He felt it as a pain in his left side, over his heart. He knew no more until he woke up in bed, when he saw his father come into the room, and he felt the bed to be the wrong way round. That was his first experience of disorientation—the first he could remember. It occurred to me, as a possible explanation of the dissociation, that the scalding of his arm produced a fainting fit, in which he fell to the left and everything twisted round to the right—he fell through a right angle. That was the beginning of his disorientation.

To make quite sure of this I hypnotized him again to bring up an earlier memory still. I suggested to him that he should remember his second birthday. He suddenly began to shout out: "He has bit me—Gordon has bit me!" Before I woke him up, I asked him all about it, but he could not tell me very much whilst in the hypnotic state—he had gone back to a period when he was too young to be able to describe his experiences. I told him that he would continue to remember all that he had just gone through, and I then woke him up. He told me that Gordon was not a dog, but a little cousin of his. They had been in bed together, standing up in bed, and it appears that he had punched Gordon in the face, and that Gordon had retaliated by biting him in the left arm. He could remember that at that time he was living near a Fire Station—could remember all the details of the incident, and could remember definitely that he had had no feeling of disorientation. The revival of this incident brought no memory of any earlier disorientation. The next day after this treatment he felt more disorientated than ever, for he had been going through these experiences very vividly. I let him stay in bed, and after two days he recovered completely, and very soon applied for his discharge. I let him go, but asked him to write to me if the feeling of disorientation ever recurred. I have never heard of him since.

This case seems to me to show that such memories date from very early life, and more than that, it shows that the memories need not always be of a sexual nature, because at the age of 2 even Freud would admit that there can be very little sex, and it is very difficult to see where the sexual experience could come in in such a case.

The other case is a much simpler one, but it is what I would call a crucial case of the value of abreaction, or the working off of emotion under hypnosis. It is the case of a gunner who was admitted to the hospital where I was working, after he had spent two years in military hospitals of different kinds. He was suffering from a tremor of the right hand, dating from the time when he had been blown up at Ypres. He did not remember anything more until he reached hospital, and the memory of this interval

had never been recalled to him by any of the doctors he had previously seen. I sent him to sleep—that took just about three seconds—and then suggested to him that he should live again through the experience of Ypres. He did so, and began to shout out all sorts of things which showed what had happened at the time. Bosch shells were falling nearer and nearer to the gun-pit. He was apparently serving the gun, and someone else was handing him ammunition, and this person had evidently lost his head, for my patient shouted out: "What the — do you mean by pulling the — pin out of that — fuse?" Then I noticed that he was moving the handle with his right hand; his hand began to shake violently, and soon he was shaking all over, but especially in his right hand. Then he suddenly became absolutely still. I suggested to him that he would continue to remember all that he had just gone through and then woke him up. He looked at his hand, which was absolutely still, with amazement, and expressed his gratitude, but his mind still appeared somewhat confused, so I told him to go and sleep it off. An hour later he came back and told me that he had not been to sleep, but that he had been thinking it all over. He knew everything that had happened, and told me that he had not been suffering from shell shock but from gun shock. His gun had been blown up, and the emotion which this experience had excited in him had been bottled up for two years, with the result that he had suffered from this tremor in his hand. The next morning he was able to shave himself with an ordinary razor, for the first time since his illness.

Now, was this working off of the emotion the cause of the recovery? The alternative explanation is that the recovery is due to suggestion. When this man came to me, however, he certainly did not expect me to cure him. I saw him the first day that he came to the ward and treated him at once, and he was not likely to have more confidence in me than in anyone else who had treated him. As regards the hypnosis, he went off straight away without realizing that he was being hypnotized—it was not that that impressed him. As far as one can make out, there was no expectation worked up—it was simply the working off of the emotion that cured him. That is why I call this a crucial case. In many other cases it may be argued that the cure is the result of suggestion—for example, in the case of the disorientated man—but this case of the gunner was not of that sort. I do not for a moment wish to deny the working of suggestion in curing such cases, but I wish to emphasize the curative effect of the working off of emotion as a fundamental factor. It is a more causative treatment than suggestion. Suggestion removes the symptom; abreaction removes the cause of the symptom.

The explanation of psychasthenia, compulsion neurosis, and some of the psychoses, in terms of the same concept of mental conflict and dissociation must be postponed.

In conclusion, I should like to express my indebtedness to Major R. Worth, O.B.E., R.A.M.C., of Springfield War Hospital, for his courtesy in allowing me to report the above two cases, and for his unfailing help and encouragement while I was working in that hospital.

THE IMPORTANCE OF THE HOUSE-FLY AS A CARRIER OF *E. HISTOLYTICA*.

BY

P. A. BUXTON, M.R.C.S., L.R.C.P.

FELLOW OF TRINITY COLLEGE, CAMBRIDGE; CENTRAL LABORATORY,
AMARA, MESOPOTAMIA.

THE investigation here described was undertaken to determine as far as possible the extent to which the house-fly is a carrier of the organisms causing diarrhoea, dysentery, etc. No one would now attempt to deny that the fly can carry a very large number of different species of pathogenic organisms; the present paper tries to afford some actual statistical evidence of the extent to which it actually does carry them. So far as I know, we have very little real knowledge of the relative importance of water carriage, flies, carrier cooks, etc., either in this country or in any other. The investigation has been carried through in one small area in Amara, on the lower Tigris, between February 1st and November 3rd, 1918,