to be found after treating the iris with silver nitrate. They are also, though not commonly, seen in the early stages of cataract. It must not be forgotten that there are apparently healthy eyes which habitually see haloes round lights in the dark; this phenomenon is probably connected with a faulty state of the patient's refraction. Transient haloes round lights are often complained of by those who suffer from conjunctivitis associated with mucous discharge. Wiping or washing away the offending flake of mucus from in front of the cornea will abolish the symptom.

CONCLUSION.

It is not pretended that this lecture is exhaustive; it merely touches the fringe of a large subject, but it is an effort to help you, by drawing your attention to matters which may too easily be relegated to the background of your mind, with harmful results to your patient's sight and with damage to your own reputation.

NEUROLOGICAL JOTTINGS.*

BY

JAMES TAYLOR, M.A., M.D., F.R.C.P.,

PHYSICIAN TO THE NATIONAL HOSPITAL FOR THE PARALYSED AND EPILEPTIC, QUEEN SQUARE; CONSULTING PHYSICIAN TO MOORFIELDS EYE HOSPITAL, AND TO THE QUEEN'S HOSPITAL FOR CHILDREN.

I WISH in these notes to direct attention to some points of interest that have come under my notice in the course of neurological work.

Complications of Epilepsy.

When I was seeing, through a long course of years, many epileptics every week at Queen Square, I was not infrequently surprised by curious developments. I remember, for example, one man, treated for many years before I saw him as a chronic epileptic. He had been attending the hospital for seventeen years, and there was nothing in his history to suggest that his case was anything more than one of chronic epilepsy. But one day he appeared at the out-patient room complaining of severe headache, and ophthalmoscopic examination revealed the presence of marked optic neuritis. He was admitted to the hospital, and died in a few weeks of a large tumour of the right hemisphere. Another case which surprised and interested me was that of a healthy looking young man, a railway signalman, who came as an out-patient one afternoon with the history that he had had an attack characterized by loss of consciousness, convulsions, and tongue biting. There was no family history of any such illness, and his personal history was uneventful. He had never had syphilis, and he had no optic neuritis. There was no doubt that the attack was of the nature of epilepsy, and naturally he could not remain at his work as a signalman. He became a barrier ticket collector, and for over four years attended the hospital regularly, and was treated with bromide. At first the attacks occurred about once a month, but they gradually disappeared under treatment. He continued to attend, and one day complained of severe headache which he said he had had for two or three weeks. He had not been subject to headaches before. I examined his eyes then (they had been carefully examined from time to time before, but nothing abnormal discovered) and I found a flaring optic neuritis in each eye. He was admitted, and a decompression operation done to relieve his headaches. He had a large growth in the left hemisphere, from which he died in a few weeks.

I could record a good many similar cases, and I feel sure my colleagues at Queen Square have had similar experiences. I suspect they are cases characteristic of a large group, and the existence of such cases shows how necessary it is not to be content with the diagnosis of "ordinary epilepsy." I remember Dr. Hughlings Jackson, remarking on one occasion that there is no such thing as a case of "ordinary epilepsy," and that every case of convulsion or so-called fainting ought to be investigated most fully and carefully, and kept under observation. These cases would seem to "point the moral," but it is obvious that diagnosis of the true cause of the convulsion is in the first instance impossible.

*Part of this paper constituted the Presidential address to the Harveian Society.

Variability in Response to Treatment in Epilepsy.

I have frequently been struck with the variability in response to treatment of some cases of recurrent loss of consciousness with or without convulsion. One case will not be influenced by bromide until he is getting up to something like 30 grains thrice daily. It may then be found that such a patient not only remains free from attacks with such treatment, but improves greatly in general health. I have a patient I see now about twice a year who for the last seven years has taken 35 grains twice a day, and has remained well, and able to carry out responsible duties in a City office. I have another patient, a lady, who for five years has remained well although taking 30 grains thrice daily.

On the other hand, one has seen patients who, with large doses of bromide, become progressively worse, but who improved on different treatment. One little girl was brought to me at the hospital some years ago who suffered from frequent attacks, several a day, of loss of consciousness without convulsion. She was only 6, and I put her on 10 grains of bromide thrice daily. In a month I saw her again. The attacks had become more frequent. I stopped the bromide for a month and saw her again. The attacks were not so frequent as while she was taking the bromide, but as frequent as when I saw her first. I then put her on 5 minims of tincture of belladonna thrice daily and saw her again in a month. There had been no attacks during that time and she remained well. Of course, I cannot explain this. I only state the facts. I understand that a similar result not infrequently follows the administration of *Adonis vernalis*.

Another interesting point I have observed about so-called "nocturnal" epilepsy. I may say, in passing, that this is an unfortunate term and really incorrect. It would seem to indicate that in these cases the fits occur only at night, but this is not the case. It is really sleep which determines the onset of an attack, for in many of these cases of so-called nocturnal epilepsy inquiry will elicit the fact that a fit occasionally occurs during the day, but only if the patient has happened to go to sleep during the day. The point, however, which I wished to emphasize is that while bromide alone is very useful in these cases, its effect seems to be reinforced by the addition of tincture of digitalis, and 30 grains of bromide with 5 minims of tincture of digitalis taken nightly are, as a rule, effective in controlling the attacks.

Dislocation in Epilepsy.

There is one point about epilepsy which is worth remembering. As is well known, after a Jacksonian fit affecting one side and starting in the arm there is frequently temporary weakness of the affected side, usually more marked in the limb in which the convulsions started—that is to say, in the arm in such a case as I refer to. But it must also be remembered that dislocation at the shoulder-joint not infrequently occurs in the course of a convulsion. I remember one case in which one of these conditions was remembered by the observer but the other forgotten. A junior colleague of mine, in my absence on one occasion, saw my out-patients. One of them was a young man subject to occasional fits. He came up on this occasion saying that he had had a fit the previous week and found that after it he could not use his arm. There was a note made to the effect that weakness of the right arm was present, and that although the right leg was not complained of there seemed to be an exaggerated knee-jerk on that side. Obviously my colleague, who had not seen the patient before, thought he was dealing with a Jacksonian fit with weakness of the affected side following the fits. But when the patient came up the following week it was found that he had only a dislocation at the shoulder-joint, and when this was reduced the weakness cleared up.

Recurrent Attacks of Herpes.

I have been considerably interested in herpes, and one or two jottings on this disease are, I think, of interest. I rather think we were all taught that a second attack of herpes is practically unknown. I think I am right in saying that that great observer, Jonathan Hutchinson, was in the habit of telling us that second attacks of herpes did not occur. I have seen three cases which certainly controvert this. One is the case of a man whom I saw at the Moorfields Eye Hospital for some eye condition unrelated

to herpes, who came up one day complaining of an eruption on his neck. I told him it was an attack of "shingles"—cervical herpes—and he told me that he had had an attack on his chest some years before. On stripping him it was obvious from the marked scarring that he had had a severe attack of herpes zoster. The second case was that of a friend of mine—a medical man—whom I saw about twenty years ago with herpes frontalis. About two years ago he had another attack of herpes frontalis on the other side, and he told me that in the intervening years he had had two other attacks of herpes, one in the deltoid region and one of herpes zoster. So that this patient had had in all four separate distinct attacks of herpes, two of which I saw myself.

In a patient whom I saw with Dr. Melvill Green a few years ago—an old lady of 80—a condition was present which I have never seen before—namely, herpes of three different distributions. She had had, about two years before I saw her, not only herpes frontalis but also herpes in the deltoid region and herpes zoster. It was on account of the very severe post-herpetic pain that I was asked to see her, and the scarring in all three regions was extremely severe. The eruption in all three distributions in this case occurred at the same time. She died soon after I saw her, exhausted by the severe and intractable pain. I have also recently seen, with Dr. Robertson-Fullarton, a lady with facial herpes and severe post-herpetic pain who a year ago had an attack of herpes zoster.

This leads me to remark on another point—namely, the curious variability in the occurrence of post-herpetic pain. Such pain in children is, in my experience, practically unknown; in the majority of adults it is severe, but sometimes of comparatively short duration. In a certain number it is not only severe but persistent, and not infrequently leads to a great impairment of health. In old people especially post-herpetic pain is always fraught with a certain amount of anxiety because of the exhaustion to which the pain and consequent loss of sleep give rise. On the other hand one has met some adults, by no means young, who have experienced no pain at all after an attack of herpes.

Herpes and Chicken-pox.

A very interesting point about shingles is its relationship to chicken-pox. Of this relationship the evidence seems convincing, but its explanation is still very difficult. In the *British Journal of Dermatology and Syphilis*, October to December, 1917, there is a most interesting paper by Dr. W. P. Le Feuvre, in which he actually urges that "shingles should be recognized as an infectious disease, and one under certain unknown conditions likely to become a starting point for a chicken-pox epidemic." Reference to the possible connexion of these diseases has been made, especially in the correspondence columns of the *BRITISH MEDICAL JOURNAL*, and evidence in favour of it adduced. The points common to all the cases brought forward were three in number—

1. Chicken-pox in one individual followed shingles in another within the ordinary incubation period of chicken-pox—namely, twenty-one days.
2. No other source of infection was discovered; and
3. Chicken-pox in a child followed an attack of shingles in an adult.

In Dr. Le Feuvre's paper fifty cases in all are referred to. Before seeing this I myself had been struck by the curious coincident occurrence of these diseases, and I had notes of four instances. One of these—a lady with a very severe attack of herpes of the fifth nerve affecting all its branches—I saw in consultation with Dr. Woollerton. A fortnight later I was asked to see the same lady again, and the doctor asked me also to see her daughter, a young lady of 28, who had such a severe attack of chicken-pox that she had been afraid it might be small-pox. I also saw, in consultation with Dr. Haward, a patient suffering with severe neuralgia after herpes zoster. Being interested in the matter, I asked Dr. Haward if there had been any cases of chicken-pox in the house at or about the same time. He was surprised at the question, but on my explaining the reason for it, he said that three children in the house certainly had had chicken-pox, but he could not remember whether it was about the same time. He promised to look into the matter of dates, and was kind enough to write me to the effect that chicken-pox in two children in the house

had appeared about a fortnight after the shingles eruption in the father. Another patient I saw in consultation with Dr. Hewetson on account of pain and actual muscular wasting following a severe attack of herpes in the deltoid region. He also was surprised at my question, but was able to tell me that two children in the house had suffered from chicken-pox within a week or two of the attack of herpes in the adult. Of course I cannot explain this curiosity. The vesicles of herpes are almost, if not quite, indistinguishable from those of chicken-pox except that in the former disease they are so often confluent. Yet one frequently sees in cases of herpes isolated vesicles in parts of the body even remote from the herpetic area—a fact which was brought to my notice some years ago by Dr. Pernet. Dr. Pernet saw with me the medical man to whose case I have already referred, who had four attacks of herpes, and this doctor's only daughter within three weeks of her father's fourth attack of herpes had an attack of chicken-pox. It is stated in the paper I have referred to that shingles may occur in an individual in the house in which chicken-pox is present, and that shingles and chicken-pox may occur coincidentally in the same individual. I have not myself seen either of those occurrences, unless the isolated vesicles in remote parts of the body in cases of shingles are really vesicles of chicken-pox.

Lethargic Encephalitis.

Those of us who are working amongst neurological cases are unfortunately only too familiar in these days with cases of lethargic encephalitis. In some of them the condition is fatal, and the changes in the central nervous system now regarded as characteristic are found. In another class of cases recovery takes place, but the patients are gravely altered, and this alteration remains permanent. In most of them the condition established by the disease is practically indistinguishable from paralysis agitans, although when tremor is present it is not of the regular rhythmical type usually met with in paralysis agitans. In many, however, tremor is absent, but the stiff gait, the mask-like immobile face, the low monotonous voice, and often the everted, dribbling lip, make the resemblance to paralysis agitans very striking.

I wish, however, to call attention to what I believe are examples of this illness of a very mild type which apparently make a good recovery. I had recently under my care at Queen Square a young woman admitted on account of diplopia who on examination presented no abnormality except the presence of nystagmus and some general weakness. All the reflexes were normal, and this, of course, excluded the diagnosis first thought of—namely, disseminated sclerosis. And the history was significant; it was to the effect that a few weeks before admission she had become ill—very drowsy, but could be roused to take her food. After partaking of it she at once went to sleep again. This sleepiness had passed off before she came to the hospital. The diplopia gradually ceased to trouble her, but a little nystagmoid jerking still remained when she left the hospital. I cannot but think that this was a mild case of encephalitis lethargica.

Another case I saw in private, a young healthy man of 35 who had been working very hard for a long time. Diplopia in his case was the first trouble, and when I saw him he had obvious weakness of one sixth nerve. He had been sleeping badly. I advised his going to bed. At first his sleep was not good, then it became unusually good, and there was rise of temperature and slight delirium. When I last saw him, however, he was improving in the most satisfactory way. His father was a medical man, and I think I should have heard had there been any relapse. This also I am inclined to think was probably of the same nature, although not so definitely as the other cases I have related.

A SERIES of congresses will be held at Monaco on April 15th and following days. They include congresses of hygiene and climatology, mineral spas, and seaside resorts.

THE Council of l'Assistance Publique has drafted a scheme for the improvement and extension of the hospitals of Paris. Extensive alterations in existing hospitals are proposed in order to bring their various departments up to date, both for the treatment of patients and for clinical instruction; it is intended also to provide several new hospitals, built and equipped in the best manner.