fibroblastic cells which are found in certain areas. The presence of stellate cells of pre-adipose tissue type is also of interest.

The fourth specimen of a free body was removed by Major Hume from the knee-joint of a patient who gave a history of recurrent attacks of pain in the joint for many years. The specimen is an irregular round flat body, 4 cm. in its largest diameter, 3.5 in the smallest, and about 0.5 cm. thick. The edges are rounded, and the surfaces are nodular and fissured but covered with a layer of shining, membranous-like tissue. On section the structure resembles modified hyaline cartilage. There can be little doubt that this specimen altered in shape and increased in size after separation.

Microscopic examination of this specimen shows changes which differ somewhat from those described in the previous specimen. No typical articular cartilage is seen. Part of the specimen is composed of a form of young hyaline cartilage mixed with which are areas of atypical fibrocartilage. In some parts of this tissue a deposit of calcium salts is found (Fig. 6). Scattered throughout the section, especially towards the surface, are groups of cells which, on staining by Schmorl's method, are found to be typical bone cells. These do not, however, show the structure of true bone when stained with haematoxylin and eosine, or with picrocarmine. In the centre of the specimen there is an area of degenerating tissue in which lies a small mass of true bone; this appears to be a portion of a fragment of the original bone showing typical bone trabeculae, with Haversian canals surrounded by their system of concentric lamellae. The fat cells of the bone marrow and the blood vessels of the marrow and bone are necrotic. The bone cells do not stain well.

In this specimen again there is evidence of proliferative change both in the cartilage and bone after separation, but this change is more advanced and the resulting structures are more atypical than in the other specimens described.

It seems fair to assume that the changes in structure found in the detached cartilage and bone of the four specimens described occurred after separation from the articular surface; even if this is not admitted, the long history of pain and inconvenience—which ceased in two of the patients after removal of the free body—is evidence that these free bodies had been present in the joint for some years. The staining properties and the appearance of the cells on section show that they were alive at the time of their removal from the joint cavity.

CONCLUSIONS.

It may thus be assumed that nutritive processes were being carried on after separation. That these processes were of an active type is shown by the proliferative changes described.

There are two possible sources from which this nutritive material might have been derived:

1. From the death of cells in the fragment. From these dead cells the surviving cells might have obtained their food supply. Nutrition of this type has been described by Burrows and Neymann⁸ as occurring during the growth of cells in isotonic salt solution. This explanation, however, cannot be accepted, as there is evidence that such fragments increase in size after separation while they are lying free in the joint cavity.

2. The other explanation is that the nutritive material is derived from the synovial fluid. If this is so, then the possibility of cartilage cells and of bone cells also being nourished by the synovial fluid must be allowed.

It follows, therefore, that the articular cartilage of the joints may derive some, if not the greater part, of its nourishment from the synovial fluid.

If this hypothesis is true, then the changes which are found in the degenerative types of arthritis can also be explained. The primary cause of these degenerative types will be found not in the cartilage or bone, but in changes in the nutritive value of the synovial fluid. There is evidence, which I hope to produce later, that changes in the nutritive value of the synovial fluid may be due to changes in the vessels of the capsule of the affected joints.

REFERENCES.

- ¹ Klein: *Virchow's Archiv*, 1854, vol. lxi, p. 190. ² Weichselbaum: *Ibid.*, 1873, vol. lvii, p. 127. ³ Poulet and Vaillard: *Archiv. de physiol. norm. et path.*, 1885, vol. xvii, p. 266. ⁴ Rea: *Deutsche Zeitschrift f. Chirurgie*, 1894, vol. xxxviii, p. 1. ⁵ König: *Verh. d. d. Gesellschaft*, 1899, vol. ii, p. 1. ⁶ Rimann, H.: *Virchow's Archiv*, 1905, vol. clxxx, p. 446. ⁷ Ludloff: *Arch. f. klin. Chir.*, 1908, lxxxvii, p. 552. ⁸ Burrows and Neymann: *Journ. Exper. Med.*, 1917, vol. xxv, p. 93.

ON THE USE OF POLARIZED LIGHT IN THE DETECTION AND INVESTIGATION OF SUTURE MATERIALS EMBEDDED IN THE TISSUES.*

BY

M. J. STEWART, M.B.GLAS., M.R.C.P.LOND.,
PROFESSOR OF PATHOLOGY IN THE UNIVERSITY OF LEEDS.

[With Special Plate.]

THE use of the polarizing microscope in pathology has hitherto been mainly confined to the study of the doubly refracting lipoids, "myelin," cholesterol, cholesterol esters, and the like. Such investigations have been made by Adami and Aschoff (1906,¹ 1910²), Powell White (1908,³ 1909⁴), Stewart (1915⁵), and others. Its use as a means of detecting microscopic foreign bodies (fragments and threads of clothing, particles of wood, etc.) in the depths of wounds has been pointed out by Policard and Desplas (1917⁶). Shattock (1917⁷) has shown its value in the elucidation of a unique case of foreign body tumour formed round crystals of silica which had been forcibly intruded into the tissues of the lip eleven years before.

The present communication will show that many of the suture materials in common use by the surgeon are optically active, and that the polarizing microscope affords a ready means for their detection in the tissues as well as for their subsequent investigation. Since the optical activity of these substances is unaffected by treatment with the usual dehydrating and clearing agents, paraffin sections are all that is required for the demonstration of the phenomenon. If such a section be viewed microscopically between crossed Nicol's prisms, the particles of suture material present will, in certain instances, show up brilliantly illuminated on a dark background, and in this way they can often be picked out with great readiness, even when their presence has not previously been suspected.

It is important to notice that certain tissues are normally optically active, even in paraffin sections. Thus, the collagen-fibril bundles of dense fibrous tissue show various degrees of luminosity, at least when they are cut longitudinally or obliquely, and the same is true of the stratum corneum of squamous epithelium and of the hairs in their root follicles. The muscle bundles of striped muscle are also doubly refracting under certain conditions (? of internal strain), and even the stratum lucidum and rete Malpighii may show the phenomenon to a slight extent. In these situations it is sometimes noticeable that luminous and non-luminous segments alternate. Elastic tissue, on the other hand, would appear to be optically inactive.

THE OPTICAL ACTIVITY OF SUTURES.

The suture materials used in surgery fall into two main categories, (1) absorbable (the various kinds of catgut), and (2) unabsorbable (silkworm gut, silk, linen and "celluloid" thread, and horsehair). Kangaroo tendon occupies an intermediate position. These substances have all been examined with the polarizing microscope in the freshly prepared state, before use, and it has been found that silkworm gut, silk, linen and "celluloid" thread show intense optical activity, whereas catgut, horsehair and kangaroo tendon show the phenomenon only to a very slight degree or even not at all. This difference is accentuated by residence in the tissues. Catgut tends to lose very speedily what little optical activity it possesses, whereas that of silk and silkworm gut persists for months and years.

Before discussing the morphological appearances of the more common suture materials as seen in paraffin sections it is necessary to indicate briefly the different purposes for which they are employed.

1. Silkworm gut is now used almost exclusively for skin sutures, whose function is temporary and which are removed by the surgeon after a certain number of days.

2. Catgut is used where absorbable sutures are required, but these must be able to hold the parts in apposition

* Communicated to the Pathological Society of Great Britain and Ireland, January, 1920. The expenses of this research were in part defrayed by a grant from the British Medical Association, for which I beg to express my thanks.

sufficiently long to allow natural union to take place; for example, in stitching the deeper layers of the anterior abdominal wall after laparotomy.

3. Silk, linen, and "celluloid" thread are used under two conditions, in both of which it is necessary that the mechanical action should be exerted over a considerable period of time, namely, (a) where the suture will ultimately cut out spontaneously and escape from a free surface, as in closing off the duodenum after partial gastrectomy, and (b) where it will probably remain permanently in the tissues, as in arterio-venous anastomotic operations and in ventrifixation.

MATERIAL STUDIED.

The material on which this study is based consists of twenty-six specimens containing suture relics. Most of them are from portions of tissue which had been excised during life by the surgeon; a few were obtained *post mortem*. Eighteen specimens were found to contain silkworm gut, five catgut, and six silk. The age of the various specimens, that is, the length of time the sutures had been in the tissues, varied between wide limits. The silkworm gut periods varied from three weeks to six years, the catgut from four days to five months, and the silk from three weeks to three years.

1. *Catgut*.—This is met with in the form of irregular masses, homogeneous or very slightly fibrillated, and occasionally broken up into a few irregular pieces, never into definite discrete fibres. If the stitch has become infected it will be found embedded in pus, otherwise it is shut off by a fibrous wall which is lined internally by cellular tissue containing many endothelial cells and foreign body giant cells. None of the latter show small portions of suture material in their cytoplasm such as occurs in the case of silkworm gut. The catgut takes the acid dye, and it may or may not show a slight degree of luminosity when viewed between crossed Nicol's prisms. The rate of absorption varies greatly in different cases, and is accelerated by the presence of sepsis.

2. *Silkworm Gut*.—This is the common variety of suture relic met with in connexion with old operation scars, and the one which provokes the most characteristic and striking foreign body giant-cell reaction. The appearances in these cases suggest that small fragments of the gut become detached and remain behind in the track of the suture when the latter is removed.* They are seen as small highly refractile bodies of various shapes and sizes, often acicular or rod shaped, sometimes unstained, sometimes taking the basic dye, and always showing intense optical activity when viewed with polarized light (Figs. 1A and 1B). The smaller particles are frequently contained within the cytoplasm of giant cells (Figs. 2A and 2B); the larger have usually a number of giant cells in close relation to them. They tend to occur in small foci, with many giant cells, endothelial cells, fibroblasts, and lymphocytes, the whole often surrounded by a wall of dense fibrous tissue. In this way they may lie locked up in the tissues for years. I have never been able to convince myself that there is much evidence that foreign body giant cells absorb these silkworm gut relics. They seem rather to protect the tissues by simply enveloping or rounding off the corners of mechanically irritating particles. In some of the cases the particles of silkworm gut are both few in number and minute in size, and might readily escape detection† if looked for in the ordinary way. The use of the polarizing microscope may then greatly facilitate the search for such relics or confirm their presence (Figs. 2A and 2B).

3. *Silk*.—Silk and linen thread (including Pagenstecher's celluloid thread) are somewhat similar morphologically, and all show intense optical activity. They can be distinguished by the fact that linen fibres possess transverse cracks or striae, whereas silk fibres do not. The specimens which I have examined—six in number—all contain silk sutures. They are composed of fibres, usually tightly packed, which are triangular or polygonal in transverse section, and which, when they are cut obliquely or lie longitudinally, have often a prismatic as well as a doubly

refracting action on polarized light (Figs. 3A and 3B). In haemalum and eosin sections the fibres remain unstained except for a narrow rim at the margin, which stains a purplish hue. The whole fibre is stained bright yellow by picric acid, as in van Gieson's stain. In two of the six cases examined the suture is embedded in pus, doubtless as a result of bacterial infection; in two it is surrounded by a zone of fibroblasts, endothelial cells, and foreign body giant cells, and in one it is simply encased round with dense fibrous tissue. In the two purulent cases, the marginal fibres of the suture, which are becoming separated from one another by pus cells, show a very definite and peculiar morphological change, associated with loss of optical activity (Figs. 4A and 4B). The change affects first the peripheral part of the fibre, which becomes swollen, granular, and intensely eosinophilic. The outline changes from triangular or polygonal to spheroidal, and there is a definite increase in size. The central zone, at first unaffected and still optically active, steadily diminishes in size until finally the whole fibre becomes changed in the manner indicated, and optical activity entirely disappears. It would appear to be a preliminary process to absorption, since such fibres may sometimes be seen in course of disintegration, and it is clearly related to the presence of pus, being perhaps due to a digestive action on the part of the leucocytic enzymes.

Apart from this physico-chemical change in silk, it is very striking how both silkworm gut and silk retain for long periods the property of optical activity when embedded in the tissues. In some of the cases which I have examined the suture materials have persisted in an optically active form for three years (silk) and six years (silkworm gut), the latter even after it has been broken up into tiny fragments and ingested by foreign body giant cells.

FALLACIES.

1. The only serious source of fallacy is the presence of small, air-borne particles of cotton-wool and the like on the surface of the section, since many of these—for example, cotton-wool itself—are optically active. It is usually easy to differentiate. Suture fibres are obviously embedded in the substance of the tissue; extraneous materials lie in a plane above (or below) that of the section, and are seen to be overlapping the cellular structure of the part. Sutures tend to be focally distributed more or less in relation to scar or granulation tissue, possibly with giant cells in their immediate neighbourhood; air-borne fluff may be found on any part of the slide.

2. As already mentioned, very dense fibrous tissue, striped muscle, and the stratum corneum of squamous epithelium, all possess slight optical activity, but the degree of luminosity is so slight and the anatomical arrangements are so obvious that no difficulty should arise.

SUMMARY.

1. Sutures or ligatures of silkworm gut, silk and linen thread—that is, all the ordinary suture materials except catgut—are intensely optically active, and the polarizing microscope affords a ready means of determining their presence in tissue sections.

2. These various materials may retain their optical activity in the tissues for years, and in the case of silkworm gut even after the fibres have been broken down into tiny pieces and ingested by foreign body giant cells.

3. Catgut is much less optically active than the other substances named, and the property tends to be speedily lost in the tissues. For this material, therefore, the method may be inapplicable.

4. Ordinary paraffin sections are all that is required for the demonstration of the phenomenon.

Addendum.—Since this paper was written Dr. J. G. Greenfield has very kindly lent me the histological preparations relating to the experimental investigation of nerve suture materials carried out by him in conjunction with Mr. Percy Sargent.⁸ The experiments were made on the great sciatic nerve of rabbits, and extended over periods of from twenty-one to eighty-one days. Suture relics were visible microscopically in 13 cases, and in these the materials used were catgut of various kinds (6 cases), silk (3 cases), linen thread (3 cases), and Japanese silkworm gut (one case). The last three show very intense optical activity in every instance. Catgut is variable: some specimens are optically quite inactive;

* I am informed that this is quite in keeping with clinically observed fact—namely, that silkworm gut sutures are often rough and eroded at the time of removal. Sutures which have been repeatedly sterilized by boiling are much more liable to undergo this change, and are, indeed, sometimes frayed when inserted.

† Some were in fact so missed and only discovered when the sections were re-examined with polarized light.

most show fairly definite luminosity with crossed Nicol's prisms, at least in parts—even a specimen which had been eighty-one days in the tissues.

I beg to express my great indebtedness to Professor J. H. Teacher and Dr. J. G. Greenfield for the loan of valuable material, and to Mr. E. R. Flint for useful criticism and suggestions.

REFERENCES.

¹ J. G. Adami and L. Aschoff: On the Myelins, Myelin Bodies, and Potential Fluid Crystals of the Organism, *Proc. Roy. Soc. Lond.*, Ser. B, 1926, 77, 359. ² L. Aschoff: Zur Morphologie der lipoiden Substanzen, *Ziegler's Beitr.*, 1910, 47, 1. ³ C. Powell White: Cholesterol, Fluid Crystals, and Myelin Forms, *Med. Chron.*, Manchester, 1908, 47, 403. ⁴ C. Powell White: On the Occurrence of Crystals in Tumours, *Journ. Path. and Bact.*, Cambridge, 1909, 13, 3. ⁵ M. J. Stewart: On the Cellular Reactions Induced by Local Deposits of Cholesterol in the Tissues, *Journ. Path. and Bact.*, Cambridge, 1915, 19, 305. ⁶ A. Policard and B. Desplas: "A propos de la mise en évidence des corps étrangers microscopiques dans les plaies de guerre. Utilisation de la lumière polarisée." *Compt. rend. Soc. de Biol.*, Paris, 1917, 80, 248. ⁷ S. G. Shattock: Pseudo-tuberculoma Silioticum of the Lip, *Proc. Roy. Soc. Med.*, 1917, vol. x, Sect. of Path., p. 6. ⁸ P. Sargent and J. G. Greenfield: An Experimental Investigation of Certain Materials used for Nerve Suture, *BRITISH MEDICAL JOURNAL*, 1919, ii, 407.

A Clinical Lecture ON GALL STONES.

DELIVERED AT GUY'S HOSPITAL

BY

R. P. ROWLANDS, M.S., F.R.C.S.,
SURGEON TO THE HOSPITAL.

I PROPOSE to make some general remarks upon gall stones and to illustrate my lecture by three interesting cases.

Nearly all gall stones form in the gall bladder, and are due to:

- (a) Infection of the gall bladder.
- (b) Over-concentration of the bile.

(a) The infection may occur through the blood stream, through the lymphatics, or ascend along the ducts from the duodenum. The researches of Rosenow indicate that infection by the blood stream is by far the most important, and cultivations obtained from infected human gall bladders often produce similar lesions in animals. It is very probable also that infection spreads to the gall bladder along the lymphatics from diseases of neighbouring organs, as, for instance, the appendix, duodenum, and stomach. The most common organisms to be found are the streptococcus, *B. typhosus* and *B. coli*. These have been cultivated from the centres of gall stones, from the bile, and from the wall of the gall bladder. It is well known that infection of the gall bladder often occurs in enterica and that gall stones frequently follow it.

(b) That concentration of the bile, leading to the deposition of mucus and crystals, is an important factor is clear, for nearly four-fifths of the patients suffering from gall stones are women who have had children, and in them symptoms often date from the time of pregnancy, when the bile is very concentrated. Stasis of the bile in the gall bladder is an important additional factor, therefore in a pendulous weak gall bladder, or in one the duct of which is linked or obstructed in any way, gall stones are likely to form.

The cystic duct is very narrow and tortuous and its mucous membrane is raised into valvular folds, so that it is very difficult for any but very small stones to pass through it (Fig. 1). Moreover, it is much narrower than any part of the common bile duct, except the termination of the latter at the duodenal papilla, therefore the large majority of sufferers from gall stones never develop jaundice.

It is also important to remember that the part of the common bile duct traversing the pancreas is not as easily distended as the first part of the duct; hence it often happens that the common bile duct is shaped like a funnel, and stones become impacted in the upper part of the pancreatic portion of the duct. The duodenal papilla is very narrow, and minute stones which will pass freely through the other parts of the ducts become arrested here. A stone smaller than a pea may thus lead to death; when it is impacted at this point it dams up both the bile and the pancreatic juice.

SYMPTOMS.

1. *Stones in the gall bladder* may cause (a) recurrent attacks of pain in the right hypochondrium associated with vomiting and local tenderness. These attacks are often mistaken for gastritis. It may be possible to feel the gall bladder during the colic, or on deep inspiration there may be "a catch" or pain in the gall-bladder region, and local tenderness on palpation. Or (b) a dull aching pain in the right hypochondrium and right shoulder, with a distaste for food and flatulent dyspepsia. It may not be possible to feel any swelling unless the cystic duct be obstructed or cholecystitis with local peritonitis develop. There is no jaundice unless the common duct also is obstructed.

2. *A stone in the cystic duct* may cause colic at intervals, with a swelling having the characteristics of a distended gall bladder. Usually this is in the right hypochondrium, but it may reach the right iliac fossa. Jaundice is absent unless the stone is junctional and presses upon or otherwise obstructs the common or hepatic duct. The stone usually falls back and the colic abates after a few hours, but it may remain impacted and cause inflammation with suppuration, or even sloughing of the gall bladder. Grey patches appear at the fundus of the gall bladder or below the stone in the duct. Perforation may take place and local or general peritonitis ensue, or adhesion to the duodenum, colon, or stomach may take place, with the formation of a fistula. It is still a common error to believe that gall stones are unlikely without a history of jaundice, but it should be remembered that as the cystic duct is narrower than the common duct, except at the papilla, more stones get impacted in the cystic duct than elsewhere, and that jaundice is really an exceptional and late sign of gall stones. This error has condemned many patients to years of unnecessary suffering, a diagnosis of gastritis or hysteria being commonly made.

3. *A stone in the common bile duct* usually causes colic and jaundice sooner or later, and fever due to an ascending infective cholangitis is sometimes present during the attack of jaundice. Sometimes stones impacted in the common duct do not cause colic but only nausea. There is often tenderness on deep palpation over the common duct above and to the right of the umbilicus. Small stones in a dilated common bile duct, although they may cause frequent colic, may not obstruct long enough to produce jaundice. It is imperative, therefore, to examine the common duct carefully at the operation in every case, even when there is no history of jaundice. The gall bladder is usually contracted, shrunken, and matted

FIG. 1.—Anatomy of the biliary apparatus.

down by adhesions in these late cases, so that no gall bladder swelling is evident. In fact, if the gall bladder can be felt in a jaundiced patient, the obstruction is probably due to other causes, such as growth of the head of the pancreas or chronic pancreatitis.

DIAGNOSIS.

(a) *Stones in the Gall Bladder.*

The diagnosis of gall stones in the gall bladder when there is no history of definite biliary colic may be very difficult, for the symptoms may closely resemble those of

M. J. STEWART: USE OF POLARIZED LIGHT IN DETECTION AND INVESTIGATION OF SUTURE MATERIALS EMBEDDED IN TISSUES.

In each instance Figure A represents the ordinary microscopic appearances, Figure B the same field viewed between crossed Nicol's prisms.

(Drawings by Miss Ethel Wright.)

Fig. 1.—Low power. Case 2: duration six years: silkworm gut. The patient had been operated on six years before for appendicitis. The tissue containing the suture relics was removed from the wall of the gut and examined ever since. Note numerous small portions of silkworm gut, many of them ingested by foreign-body giant cells. The lesion is bounded by a layer of dense fibrous tissue (seen here on two sides) separating it from the surrounding striped muscle.

Fig. 3.—Low power. Case 6: duration three years: silk. From an old case of ventrifaction. At the second operation a small painful tumour was excised from the depths of the scar. To show numerous silk fibres cut transversely or obliquely, lying in the midst of necrotic debris and pus. Some of the fibres to the left hand side of the field show the pre-absorptive changes mentioned in the text.

Fig. 2.—High power. Case 8: duration six years: silkworm gut. From a case of recurrent renal calculus in which a portion of the old scar was excised at the second operation. To show a large foreign body giant cell containing two small portions of silkworm gut.

Fig. 4.—High power. Case 19: duration three months: silk. Old gastro-enterostomy case. To show pre-absorptive changes in silk fibres. The marginal portion first and ultimately the whole fibre swells up and becomes granular and strongly eosinophilic, at the same time losing its optical activity from without inwards. Note the small, optically active cores in the left half of the field.