cases of asthma and commented on surgical interference in the nose.

Dr. HAMILTON (Glasgow) regretted that there had been no combination with the Section of Psychiatry in this discussion, as suggestion often played so important a part in the cure, and he considered that asthma in general practice was far more a nerve condition than he had been led to believe that day.

Dr. GREENFIELD (Rushden) expressed the opinion that the cutaneous tests were too delicate for general practitioners, but thought peptone treatment would be useful, and asked for details.

Dr. JOHN STENHOUSE (Toronto) related several cases of asthma. He thought that the disease was probably due to toxic products.

Dr. JOHN F. WALKER (Southend-on-Sea) asked for information as to the utility of the ophthalmic reaction, and expressed the opinion that information obtained by general practitioners in this condition should be collected in some way for consideration by experts.

Dr. FREEMAN and Dr. GOW replied to various questions dealing with the ophthalmic reaction and peptone treatment respectively, and Sir HUMPHRY ROLLESTON summed up the main points of the discussion.

Demonstrations.

In the afternoon a demonstration of the skin reactions in asthma was given by Dr. MACKENZIE WALLIS in the Pathological Department. At the Royal Infirmary, in addition to ward demonstrations by the honorary medical staff, the following were demonstrated: The negative pressure in the pericardium, by Dr. G. ARBOUR STEPHENS; the treatment of tuberculosis with an attenuated tubercle vaccine, by Dr. NATHAN RAW; the value of tuberculin in the treatment of bone tuberculosis, by Dr. W. CAMAC WILKINSON; the technique of artificial pneumothorax, by Dr. F. G. COLEY.

ADOLESCENT TETANY AND ITS RELATION TO GUANIDIN.

BY

F. J. NATTRASS, M.D., AND J. S. SHARPE.

(From the Royal Victoria Infirmary, Newcastle-upon-Tyne, and the Institute of Physiology, University of Glasgow.)

IN 1917 Noël Paton and Findlay¹ showed that the symptoms of tetania parathyreopriva and of tetany are similar to those produced by the subcutaneous administration of salts of guanidin, while Burns and Sharpe² demonstrated that in the former condition in dogs the amount of guanidin in the blood and urine is increased, and that, in the latter condition in infants, a marked increase occurs in the urine. It was impossible to get sufficient blood from them for analysis. In 1918 Burns and Watson³ showed that parathyroidectomy produces similar effects on the heart to those produced by the administration of guanidin.

The conclusion seemed inevitable that the symptoms of tetania parathyreopriva and of idiopathic tetany of children are both due to some error in the metabolism of guanidin in the body.

Findlay and Sharpe⁴ in 1920 recorded a case of adult tetany in which the excretion of guanidin in the urine and the faeces was markedly above the normal. We are now able to record another case of the same kind in which the excretion of guanidin by the urine and the faeces was estimated. The clinical features of the case are of no little interest.

The patient, a girl of 15, first came under observation in January, 1920, complaining of attacks of painful spasm of the hands. From her description of the seizures tetany was suspected, and subsequently on many occasions typical attacks of tetany have been observed.

History.

The attacks commenced suddenly and without warning about a year before she came under observation. At first the spasms occurred at intervals varying from a week to a month, but they gradually increased in frequency until finally they occurred two or three times daily. The duration of each attack also became progressively longer, and the later attacks lasted for two or more hours.

In severe and prolonged attacks usually both hands, but sometimes one only, became red, hot, and painful.

The patient was occasionally awakened at night by cramp in the feet, but this was the only indication of any affection of the feet similar to that of the hands. Apart from the spasms she felt in perfect health. She was habitually constipated, but not severely. Menstruation commenced a month or two before the first attack in the hands, and was irregular and of excessive duration (frequently lasting for twelve days with intervals of only a fortnight), but she had felt no ill effects of the heavy losses. Her mother stated that the patient never had any similar attacks as an infant, nor any convulsions nor child-crowding. She had never had any serious illness.

Her father and mother were healthy, but her maternal grandmother suffered from generalized spasms which the mother (a very intelligent woman) described as being like the patient's. A great aunt on the paternal side was treated for some condition with thyroid extract. A sister, now aged 22, suffered from the age of 14 to 19 years from attacks exactly resembling those of the patient. The patient has also two brothers, who are healthy. No evidence of infantile convulsions or laryngismus stridulus in any member of the family was obtained.

Present Condition.

The patient is healthy in appearance, well nourished (weight 7 st. 7 lb.), and intelligent. There is a slight uniform enlargement of the thyroid gland, no greater than that often seen in girls at puberty, and no signs of hyperthyroidism. The left thumb is deformed, the terminal phalanx at the interphalangeal joint being deviated to the ulnar side at an angle of about 20 degrees. Examination reveals no other abnormality in any part of the body.

The attacks of tetany occur with no apparent cause, and do not appear to be influenced by nervousness nor capable of being produced by suggestion. An attack in one hand can be induced by pressure on the corresponding nerve trunks in the upper arm (Trousseau's sign), and a very severe attack in the right hand followed shortly after galvanic testing of the right ulnar nerve.

There is very marked twitching of all the muscles on one side of the face on percussion of the corresponding facial nerve near its exit from the stylo-mastoid foramen (Chvostek's sign). The irritability of the peripheral nerves to galvanic stimulation is much greater than normal, and the characteristic reactions to the opening stimuli are strikingly manifest (Erb's sign). The reactions of the right ulnar nerve, stimulated behind the internal epicondyle of the humerus, in the patient and in a control case at the same time under the same conditions, were as follows:

	Patient.	Control.
A.O.C.	0.6 ma.	3.5 ma.
K.C.C.	0.8 ma.	1.0 ma.
A.C.C.	1.0 ma.	1.4 ma.
K.O.C.	2.0 ma.	over 7-8 ma.

Examination of the Patient's Family.

The sister, who without doubt had tetany which ceased three years ago, appears in every respect normal. The electrical reactions are normal and Chvostek's sign is not present.

The younger brother, aged 17 years, shows Chvostek's sign of quite marked degree, but much less than in the case of the patient herself. The elder brother, aged 24 years, shows a slight degree of the same sign.

THE EXCRETION OF GUANIDIN.

The patient was admitted to the Royal Victoria Infirmary, Newcastle, on April 10th, 1920, under Dr. Beattie. She was placed upon a creatine-free diet for one week, during which time the whole of the urine and a watery extract of the whole of the faeces were collected, each being evaporated daily to a small bulk with the addition of a few drops of thymol in chloroform. They were sent to Professor Noël Paton at the Physiological Laboratory of the University of Glasgow and the guanidin content was estimated by one of us (J. S. S.).

During this week of special dieting the patient had three attacks of tetany of average severity. No laxative or other drug was given during the week, and the bowels moved naturally on six days out of the seven.

Chemical Examination.

The urine and extract of faeces, each amounting to almost a full Winchester quart, were treated by the method described (Findlay and Sharpe⁴). The results were calculated as guanidin in milligrams per diem.

TABLE I.

By urine...	...	...	305.0
By faeces	...	...	17.0
Total per kilo of body weight	...	...	6.7

The examination of the picrate precipitates gave the following results:

TABLE II.

	Found.	Calculated as Dimethylguanidin.
Total nitrogen	24.0 %	26.5 %
Melting point	226° C.	225° C.

This indicates that the guanidin is chiefly in the form of dimethylguanidin, as it was in the case of adult tetany previously recorded (loc. cit.). In that case the excretion by the urine was 4.8 mg. per kilo a day, against 6.7 mg. in the present case. In the case of active idiopathic tetany in children recorded in 1917² the excretion of guanidin by the urine (determined by the aurichloride method) averaged 0.49 mg. per kilo a day, in latent tetany 0.47 mg., and in a case recovered from tetany 0.12 mg., while in five cases free from tetany the average was 0.116 mg. per kilo a day. In dogs suffering from tetania parathyreopriva the average excretion in the urine (determined by the aurichloride method) was 1.0 mg. per kilo a day.

Later experience has shown that results obtained by the aurichloride method are lower than those by the picrate method, and that the yield depends largely upon the concentration of the solution from which crystallization takes place. While of use in giving comparative results in a series of simultaneous determinations it is somewhat unreliable, and it has been abandoned for the picrate method, which has proved more accurate and much more rapid.

Subsequent observations have shown no excess of ethereal sulphates in the urine when the attacks are frequent. There is no evidence of pancreatic deficiency as tested by Loewi's adrenaline mydriasis test; the amount of diastase in the urine, and the fat, starch, and protein content of the faeces.

Treatment.

A prolonged trial has been made of parathyroid tablets (Armour), gr. 1/20, by the mouth, the dose being increased gradually to three tablets three times daily. No curative action can definitely be ascribed to this treatment, for though remissions have occurred, similar remissions have been seen when the tablets were omitted. The addition of thyroid tablets, gr. v once daily, has so far produced no appreciable effect.

A definite remission followed the week in which the patient was in hospital on a meat-free diet. But in spite of limiting subsequently the amount of meat and meat extracts in the diet, and ensuring regular action of the bowels, the attacks continue to recur at very variable intervals. There is, however, no doubt that their frequency and severity are considerably less than they were. The administration of salol by the mouth has not appeared to influence the attacks. Trial is now being made of calcium salts.

The estimation of the urinary sulphates, and the examination of the faeces for evidences of pancreatic defect, were carried out by G. J. Crawford.

REFERENCES.

¹Noël Paton and Findlay: *Quart. Journ. of Exper. Phys.*, 1917, vol. x, p. 203. ²Burns and Sharpe: *Quart. Journ. of Exper. Phys.*, 1917, vol. x, p. 345. ³Burns and Watson: *Journ. of Physiol.*, 1918, vol. lli, p. 88, and 1920, vol. liii, p. 385. ⁴Findlay and Sharpe: *Quart. Journ. of Med.*, 1920, vol. xiii.

THE SACHS-GEORGI PRECIPITATION TEST FOR SYPHILIS

AND A COMPARISON WITH THE WASSERMANN REACTION
IN OVER 1,500 CASES.

BY

T. TANIGUCHI AND N. YOSHINARE.

(From the Pathological Department of the University and Western
Infirmary, Glasgow.)

THE formation of precipitate in a mixture of syphilitic serum with lipoids or other reagents has been frequently recorded. But it was not until the work of Sachs and Georgi, and Meinicke¹ had defined more precisely the conditions necessary for eliciting the phenomenon that it appeared to compare with the Wassermann reaction in point of specificity and constancy as a test for syphilis. It has been found that with a suitable antigen (alcoholic extract of human heart reinforced by the addition of cholesterol according to the principle for the Wassermann antigen established by Browning, Cruickshank, and MacKenzie), when appropriately diluted and incubated at 37° C. for eighteen to twenty hours, precipitates are produced in the presence of heated syphilitic serums, whereas non-syphilitic serums in similar conditions fail to cause precipitum formation.

Method of the Test.

Antigen.—1 gram of minced human heart muscle (nature of case immaterial), freed from fat, is extracted at room temperature with 5 c.cm. of absolute alcohol for forty-eight hours,

shaking at intervals, and finally filtered; the clear filtrate is kept in the dark at room temperature. To prepare an emulsion for use in the test 1 c.cm. of this extract is diluted with 2 c.cm. of absolute alcohol; to 1 c.cm. of this diluted extract 0.045–0.05 c.cm. of 1 per cent. alcoholic solution of cholesterol is added. Then 1 c.cm. of cholesterolized extract is placed in a 3 by $\frac{1}{2}$ in. test tube and 1 c.cm. of 0.85 per cent. NaCl solution ("saline") is rapidly poured into it from a similar tube (when greater volumes are required correspondingly larger tubes are used): after the mixture has stood for five minutes at room temperature, 4 c.cm. more of saline are added rapidly in the same fashion as before and the emulsion is used for the test after standing for five to fifteen minutes at room temperature. The antigen should appear opalescent, but without flocculi when examined with a hand lens. (When the antigen has stood by itself for a longer time—for example, more than twenty minutes at room temperature, visible precipitation sometimes occurs in the antigen control on incubation.)

Serum (within seven days after withdrawal of blood) is tested after standing for several hours at room temperature, subsequent to heating for thirty minutes at 55° to 56° C.

As controls for a series of tests (1) negative, weak and strong positive serums are used as in carrying out the Wassermann reaction; (2) a tube containing the lipid emulsion and saline is included as the antigen control.

The reagents are measured in the following order into 3 × $\frac{1}{2}$ in. test tubes in the amounts shown. The contents of each tube

Main Test.		Serum Control.	Antigen Control for each Series of Tests.
First Tube.	Second Tube.	Third Tube.	
0.9 c.cm. saline + 0.05 c.cm. serum + 0.5 c.cm. antigen	0.9 c.cm. saline + 0.1 c.cm. serum + 0.5 c.cm. antigen	0.9 c.cm. saline + 0.1 c.cm. serum + 0.5 c.cm. 16.6 per cent. dilution of alcohol in saline	0.9 c.cm. saline + 0.5 c.cm. antigen

are immediately mixed by gentle shaking and then placed at 37° C. for eighteen to twenty hours; the results are read on withdrawing the tubes from the incubator. The antigen and serum controls and also the mixture of known negative serum with antigen should all show no flocculation.

The reading of results is carried out as in the case of bacterial sedimentation tests. In order to detect weak reactions it is advisable to use a hand magnifying lens; but in estimating such weak reactions it is essential to compare carefully the control with a known negative serum, as slight flocculation may occasionally occur in the latter. Provided the negative control and the serum control show no flocculation, then precipitation which is just visible with the magnifying glass constitutes a suspicious reaction, while flocculation visible to the unaided eye constitutes a positive result.

Results Compared with the Wassermann Test.*

In a series of 1,575 serums (including 296 cases already investigated by Taniguchi), comprising cases submitted to the laboratory from the wards of a general hospital, as well as the venereal diseases centre, the following results were obtained:

Parallel results:			
No. of cases	...	...	1418 = 90%
Reaction positive in both tests in	...	...	553
" negative in both tests in	...	...	845
" doubtful in both tests in	...	...	20
Discrepancies:			
No. of cases	...	...	157 = 10%
Positive Wassermann, but negative or doubtful Sachs-Georgi in	...	...	23
Negative or doubtful Wassermann, but positive Sachs-Georgi in	...	...	77

When cases of syphilis which have undergone specific treatment are grouped by themselves it is seen that the proportion of positive results with the Sachs-Georgi test is greater than with the Wassermann reaction.

Results in 316 Serums from Treated Cases.

Parallel results:			
No. of cases	...	...	256 = 81%
Reaction positive in both tests in	...	...	152
" negative in both tests in	...	...	97
" doubtful in both tests in	...	...	7
Discrepancies:			
No. of cases	...	...	60 = 19%
Positive Wassermann, but negative or doubtful Sachs-Georgi in	...	...	5
Negative or doubtful Wassermann, but positive Sachs-Georgi in	...	...	44

* The Wassermann reactions were carried out as described in Browning and Watson's *Venereal Diseases* (London, 1919) using human heart cholesterol antigen.