removal of the sac a single layer of ilio-tibial band was inserted in the way described and the wound was closed.

The patient replied to a letter of mine on June 16th, 1921, stating that he is a miner, and is able to do full work without any trouble; that he has never worn a truss; that the scar does not bulge on coughing or straining, and that he is most grateful for the result.

CASE II.

A. M. B.: double inguinal hernia. Left side operated on March 15th, 1917. The hernia was as large as in the first case. A double thickness of ilio-tibial band, produced by folding the piece removed diagonally, was inserted in the way described.

In answer to my inquiry the patient stated, on June 19th, 1921, that, although he has some cramp-like pains in the groin, especially after sexual intercourse, he has no bulging of the hernial region on coughing or straining, and does not wear a truss. He also expresses gratitude for the operation, which, he says, is quite successful.

I operated on the other side at a subsequent date. Here the conditions were much better, the base of the sac being smaller, and the internal oblique stronger. I did not consider a graft necessary on this side, and did an ordinary Bassini's operation, which is quite successful.

CASE III.

Bdr. F. C.: left inguinal hernia. Operation on May 2nd, 1917. The abdominal wall was very weak in the hernial region and the sac had a large opening. I could not have made a satisfactory repair by the usual method, and therefore inserted a folded graft as described.

The patient writes on June 19th, 1921, to say that the operation is quite successful. He has no pain whatever, and no bulging on sneezing, coughing, or straining. I quote his own words: "I am perfectly cured, and never felt so well before. The work I do is heavy, but I never feel any pain, so shall never want to wear a truss."

The operation has stood the test of six years in the first case and of four years in each of the others without recurrence of the hernia, and two of the patients have been, and are, doing heavy work. I am confident that repair by usual methods would not have been satisfactory, and I believe that this method may be of considerable value in difficult cases. I have not seen any trouble from the removal of the piece of ilio-tibial band. There is a slight bulging of muscle through the gap for a few days after the operation, but none of these patients had muscle hernia on leaving hospital.

For stitching the graft into position I used silk, because I thought it would remain unabsorbed for a considerable time, and would therefore hold the graft until it was firmly united to the neighbouring tissues, but, provided catgut which will last three weeks and is absolutely sterile can be obtained, I do not see any reason why it should not be used if desired. In each case healing was uneventful, and I had no trouble from the use of silk.

IONIC MEDICATION IN THE TREATMENT OF NEURITIC AND RHEUMATIC PAINS.

BY

W. F. SOMERVILLE, M.D., F.R.F.P.S.GLAS.,
LATE MEDICAL ELECTRICIAN, WESTERN INFIRMARY, GLASGOW.

CASES of neuritic and rheumatic pains are frequent events in general practice, and not a few of them prove rebellious to treatment. It may be questioned whether in such cases practitioners generally realize the benefit that may be obtained by the employment of ionic medication, especially in cases where ordinary drug treatment fails to give relief.

The treatment is easily carried out by means of the following apparatus, all of which may be obtained at small cost from makers of electrical requirements: namely, a switchboard connected with the electrical mains supply, with a sliding resistance, and with a milliampère meter in circuit; two insulated conductors connected with the poles of the battery and with flexible metal electrodes 4 in. by 3 in.; two pads, each of 16 to 20 layers of lint and of a size sufficient to prevent the metal electrodes from coming in contact with the patient's skin; 2 per cent. solutions of quinine hydrochloride and of sodium salicylate and 1 per cent. of potassium iodide for different conditions, while at the indifferent electrode a 2 per cent. solution of common salt is employed. The only other requirements are bandages to keep the pads in close contact with the patient's body and a suitable couch.

In all cases the pads, especially the one placed in connexion with the positive pole, must be thoroughly saturated with the warm solution employed. The current must invariably be applied and switched off very slowly and smoothly. A current of from 15 to 75 milliamperes is employed according to the locality of the pain and the age, sex, and condition of the patient. The séance should be given daily for twenty to forty minutes. Occasionally I have combined ionization with high-frequency massage to the general advantage of the patient and as a helpful agency in the encouragement of sleep.

Notes of a few cases lately under my care may induce others to test this method of treatment:

Trigeminal neuralgia. A lady, aged 52, sent to me in February, 1921, by Dr. Archd. Sloan, complained of neuralgic pain in the area of the left fifth cranial nerve, and particularly in the supra-orbital region. The pain had been present more or less continuously for twenty years, and was aggravated when the patient washed her face or when she was exposed to wind. Ionization with solutions of quinine hydrochloride (connected with the positive pole), of sodium salicylate, or of potassium iodide, each of the last connected with the negative pole, were employed on successive days. After thirty-three applications there was distinct improvement, and by the end of April the pain had completely ceased. I saw her on August 5th, when she informed me that since the end of April she had had no pain; and that the experiences which had previously provoked or aggravated the pain could now be enjoyed without any discomfort. In addition, her general health had greatly improved.

I have successfully treated many cases of neuralgia of the superficial branches of the circumflex and musculospiral nerves. Their intractability to ordinary methods is well known, yet, with very few exceptions, I find that relief can be secured by ionic medication, a solution of quinine hydrochloride being the most useful. One case as an example may be cited.

A solicitor, aged 52, otherwise in excellent health, sent to me by Dr. Walls Christie in January, 1921, complained of a sudden attack of pain of three days' duration in the right arm, the pain being so severe as to cripple the use of the limb. At the same time there had developed a sensation of numbness in the outer three and a half fingers. Seventeen applications of ionization with a solution of quinine hydrochloride effected a complete cure, and neither the pain nor the numbness has returned.

The average number of applications required is from fifteen to twenty, but many patients are relieved after a less extended course. A lady, 40 years of age, sent to me by Dr. R. O. Adamson on account of severe brachial neuralgia, was completely relieved after six applications. Another lady, aged 33, sent by Dr. W. McMillan, had her brachial neuralgia removed after nine visits.

Neuralgia of the smaller occipital nerve. An elderly lady, advised by Dr. Linton of Renfrew and Dr. George S. Middleton of Glasgow, came into my private nursing home in September, 1920, complaining of great pain at the back of the head and over the right parietal region and vertex. The pain had lasted for more than six months. Ordinary drug treatment and rest in bed had produced little result. Ionic medication with solutions of quinine hydrochloride and of sodium salicylate, however, gave a complete and lasting cure.

Neuralgia in the region of both tuber ischii. A young medical student was referred to me by Dr. Ernest G. Fortune on account of pain in the region of both hips. This had lasted for five years and had prevented him from engaging in any form of sport to which he had previously been devoted. He could stand, lie down, and walk slowly, but any hurried effort caused him suffering. I skiagraphed both hip-joints but discovered nothing abnormal. Quinine ionization was applied twice over each tuber on successive days with surprising effect. Three evenings afterwards he went to a dance "by way of a trial," as he remarked, and took part in every dance with no sensation of pain at the time or ill effect thereafter.

Neuralgia of the twelfth dorsal nerve. A colliery manager, aged 60, consulted me fifteen months ago on account of a severe and continuous pain over the area supplied by the twelfth left dorsal nerve. The pain had troubled him more or less for two years, and though he had tried many forms of treatment, including, he said, the surgical division of the nerve, he had derived no benefit. Ionization with quinine, however, relieved him very considerably and permitted him to continue his work, though this entailed exposure to heat and cold and to wet clothes.

I may, perhaps, be permitted to quote my own case. About nine years ago I was completely laid aside for three weeks with violent neuritic spasmodic pain in the region of the twelfth left dorsal nerve, extending to the thigh. Ionization with quinine gradually produced complete cure, and though I have since been engaged on home military service for four and a half years, have spent months under canvas, been exposed often to wet and inclement weather, and engaged constantly in field exercises and route marches, I have had no return of the pain.

In cases of pain, which for want of other explanation must be termed "rheumatic," the results of ionic medication are most gratifying. A solution of sodium salicylate connected by an electrode with the negative pole is applied to the part affected, while a solution of common salt is used at the indifferent electrode in connexion with the positive pole.

A young engineer engaged in work involving exposure to cold and damp came to me in March, 1921, with "rheumatism" in both feet. He had suffered much for several years, and could walk only with difficulty and pain. The knees and the finger-joints were also affected. After a somewhat prolonged course of ionization with sodium salicylate he made a complete recovery, and was able to walk easily and without discomfort. His swollen finger-joints became normal, and this enabled him to resume piano playing, from which he had previously been debarred.

Ionization with copper metal to the interior of the uterus gives satisfactory results in cases of endometritis and menorrhagia. The whole uterine mucous membrane is dealt with and no anaesthetic is required. The operation is of the simplest character. A patient of Dr. Alexander Thomson, suffering from a dilated meatus urinarius following childbirth, with inability to retain urine on any movement, was completely cured by the introduction of a copper probe through which a very gentle continuous current of from 3 to 5 milliampères was passed for five minutes at a time. The useful effect of zinc ionization in cases of carbuncles and boils is also well known.

It is with every confidence that I venture to encourage practitioners to test ionic medication as a simple and effective therapeutic agent in cases such as are here described.

A CASE OF CHOLECYSTGASTROSTOMY.

BY

C. A. MOORE, M.S.LOND., CH.M., F.R.C.S.,

SURGEON, BRISTOL GENERAL HOSPITAL; CONSULTING SURGEON,
COSSHAM HOSPITAL, KINGSWOOD.

The following case presents features of interest and seems worthy of record:

A married woman, aged 45, complained of ill health of seven years' duration. She suffered from attacks of abdominal discomfort half an hour after meals; they were never very severe, and she rarely vomited; she had had no melaena or haematemesis so far as she was aware. At intervals of a few weeks she had had attacks of nausea and diarrhoea. During the previous two months she had been worse and was losing weight. She was pasty-faced and sallow, but not jaundiced. The temperature and pulse were normal, as was the urine. Nothing of note was found on examination of the chest or rectum. There was a small, firm, rather fixed swelling just above and to the right of the umbilicus. The spleen was readily palpable an inch below the costal margin. The liver was not enlarged, nor the stomach dilated. A blood count showed merely slight secondary anaemia, and a barium skiagram gave no evidence of ulcer or pyloric stenosis.

Operation.

The abdomen was opened above the umbilicus. The swelling proved to be a hard nodular mass in the head of the pancreas. The liver, gall bladder, and ducts were normal to sight and touch. The splenic vein was dilated, and ran directly into the pancreatic mass, and the splenic enlargement was clearly due to venous engorgement. The diagnosis between chronic pancreatitis and carcinoma was uncertain. On the assumption that the swelling was probably inflammatory, some form of drainage of the biliary passages seemed indicated, and by cholecystenterostomy rather than cholecystostomy. In view of the great induration of the pancreas and enlargement of the spleen it seemed doubtful whether a few weeks' drainage would suffice.

Cholecystenterostomy, then, was decided upon in the hope that it would prove curative if the swelling were inflammatory. If it were malignant it would serve to prevent jaundice from the inevitable and early obstruction of the common duct. The pancreatic mass so overlapped and fixed the duodenum that no amount of mobilization would render it available. My experience has been that the use of the colon is preferable to that of the jejunum, and arrangements for cholecystocolostomy were being made when it was noticed the relation of the gall bladder to the stomach was such that an anastomosis could readily be made without disturbing their relations. This was done by simple suture in two layers and the abdomen closed. Recovery was uninterrupted.

Four months later the patient expresses herself as being completely relieved of her symptoms and having "never felt better in her life." The pancreatic mass was just appreciable to palpation, although distinctly smaller, and the spleen could still be felt.

Operations with the object of short-circuiting the common bile duct date from 1830, when von Winiwarter of Liège, at the suggestion of Nussbaum, performed the first. He used the colon and operated in six stages, lasting from July 20th, 1830, until November 14th, 1831. The first operation in this country was performed by Mayo Robson in one stage in 1889. The use of the duodenum was suggested by Cozi, and the first operations were those of Terrier and Bardenheuer. No doubt physiologically the best site for the junction is the duodenum, but as a matter of practical surgery it is not always available.

The objection to the use of the jejunum is that regurgitant vomiting, such as used to follow gastro-enterostomy with a "loop," is not unknown. This may be partially guarded against by the plan recommended by Mickulicz of performing a lateral anastomosis between the two limbs of the jejunal loop, but probably the best method is to make a retrocolic junction. An opening is made in the transverse mesocolon and the jejunum pulled through it. The operation is completed by drawing the junction down and suturing it to the edges of the aperture in the mesocolon, just as in posterior gastrojejunostomy. To the use of the colon there is the objection that there is an appreciable risk of suppurative cholangitis. In practice, however, this does not often occur, and the anatomical suitability of the colon has caused its frequent use with generally satisfactory results.

Attention seems to have first been drawn to the possibilities of the use of the stomach for this purpose by Oddi¹ in 1887. He experimented on dogs, and found that the diversion of the bile into the stomach was quite consistent with perfect health. His observations were confirmed by Cannac, Masse, Mocquot and others.

The first operation on the human subject was performed in Vienna on September 21st, 1892, by Wickhoff and Angelberger² in a case of gall stones and stricture of the common duct. The patient recovered. They were followed by Terrier³ on December 19th, 1895, in a case of carcinoma of the head of the pancreas. The patient survived ten months. Monod⁴ did a similar operation on June 1st, 1896, but the patient died the next day. Jaboulay⁵ recommended it as the operation of choice in tumours of the head of the pancreas. About this time cases were also published by Montagnon and Duchamp,⁶ and by Krumm,⁷ while Perrin⁸ in 1902 collected fourteen cases, including the foregoing, with generally satisfactory results. Kehr⁹ has several times strongly advocated the operation, and had, in 1914, performed it sixty-two times, but gives no results.

Cases in the literature since 1914 are naturally scarce, but the subject has been referred to in America by Barr,¹⁰ Jacobson,¹¹ and Erdmann.¹² The latter quotes Moynihan as having operated in this way twenty-one times with one death, and as having seen no disability from the presence of bile in the stomach. Moynihan advocates operation in all cases of obstructive jaundice, however positive may be the evidence of malignancy. The diagnosis is often difficult and never certain.

To sum up, it may be said that, while short-circuiting operations of this type will rarely be performed to-day—as were many of the earlier ones—for stones impacted in the common duct, indications will still be found, especially in cases of pancreatic disease. It is submitted that when the indication arises the use of the stomach for the anastomosis makes the operation technically easy, physiologically sound, and unlikely to be followed by undesirable complications.

REFERENCES.

- ¹ Oddi: *Arch. Ital. de Biol.*, ix, 138.
- ² Wickhoff and Angelberger: *Wien. klin. Woch.*, vi, No. 18, 325.
- ³ Terrier: *Bull. et Mém. de Soc. de Chir. de Paris*, xxii, 565.
- ⁴ Monod: *Ibid.*, xxii, 546.
- ⁵ Jaboulay: *Lyon Méd.*, lxxxix, 365.
- ⁶ Montagnon and Duchamp: *Loire Méd.*, xviii, 142.
- ⁷ Krumm: *Muench. med. Woch.*, xlviii, 826.
- ⁸ Perrin: *Thèse de Lyon*, p. 1-9.
- ⁹ Kehr: *Technik der Gallensteinoperationen, Ergebnisse der Chir. Orthopaed.*, 1914, vol. viii.
- ¹⁰ Barr: *Journ. Tenn. Med. Assoc.*, 1914-15, vii, 454.
- ¹¹ Jacobson: *Journ. Amer. Assoc. Obstet. and Gyn.*, 1914-15, xxvii, 172; *Amer. Journ. Obstet.*, 1914, lxx, 825.
- ¹² Erdmann: *Annals of Surgery*, 1918, lxxvii, 273 and 363.

THE Japanese Home Office is about to investigate the death rate of infants in Tokyo, basing itself on the police statistics of the deaths of infants under the age of 5 years and their causes from 1916-20. It is believed that in the poor quarters of Tokyo the infantile death rate reaches 37 per cent. of births.