dried bodies. At the same time, further rabbits are being prepared with autolysates of these virulent meningococci, and others with the aqueous extracts of the same cocci after being dried and powdered. The value is also being determined of sensitizing the cocci, or its contents, before injection. By this means it is hoped to define the relative value of each of these preparations respectively of the same meningococcus for exciting the production of anti-endotoxin. It may be of interest to state that one rabbit that had received intravenously a dose of 0.1 c.cm. of the aqueous extract of a virulent example of Type I meningococcus prepared in the standard manner described previously, followed a week later by a dose of 0.2 c.cm., in spite of considerable loss of weight, showed on the sixteenth day after receiving the first dose sufficient anti-endotoxin in 0.5 c.cm. of its serum to neutralize completely a lethal dose of Type I endotoxin for the mouse. Captain Tulloch also found this serum to contain a good yield of opsonin for Type I. Here the matter at present stands.

CONCLUSION.

It would seem, then, that, as regards anti-endotoxin, there is at the present time much room for improvement in, at any rate, a considerable proportion of the samples of antimeningococcus serum supplied for therapeutic use. But it should be stated that several of these serums, though deficient in anti-endotoxin, have been found by Captain Tulloch, who is making a special study of meningococcus opsonins, to contain a good amount of opsonin which in the eyes of Flexner and some other authorities is of almost, if not quite, equal clinical importance. However this may be, few, I think, will deny that it is very desirable that all antimeningococcus serum sent out for therapeutic use as "multivalent" should contain as high a yield as possible not only of opsonins but also of anti-endotoxins capable of neutralizing the toxic principles of the commonest members of the meningococcus group.

The chief obstacle hitherto has been the absence of a trustworthy and simple procedure for determining the amount of anti-endotoxin in antimeningococcus serum. It is hoped that the present method may be found to contribute towards the end in view, because it would seem that some of the best results in the way of lowering the mortality from cerebro-spinal fever during the present outbreak have been obtained by early intrathecal administration of serum that, in addition to a good yield of opsonins, has also been found to contain an appreciable quantity of antibodies capable of neutralizing the endotoxins of each of the two commonest types of the meningococcus.

BIBLIOGRAPHY.

- Flexner, S.: *Journ. Exper. Med.*, vol. ix, 1907.
 Dopter, C.: *Annales de l'Institut Pasteur*, vol. xxiii, 1910, pp. 96-113.
 Kolle, W., and Wassermann, A.: *Deut. med. Woch.*, vol. xxxii, No. 16, 1906, pp. 609-612.
 Besredka: *Annales de l'Institut Pasteur*, vol. xix, 1905, pp. 477-480.
 Ibid., vol. xx, 1906, pp. 149-154, and pp. 304-310. Also article "Endotoxines Microbiennes," *Bulletin de l'Institut Pasteur*, 1914, pp. 145-154, and 195-205.
 Wassermann and Leuchs, Kraus, etc., quoted from article "Die Wertbestimmung des Meningokokken-Serums," by R. Otto and K. H. Boehneke in Kolle and Wassermann's *Handbuch der Pathogenen Mikroorganismen*, 2nd edit., Jena, 1913, Bd. 2, pp. 1230-1233. See also article "Gewinnung des Meningokokkenserums," by St. Baecher, in Kraus and Levaditi's *Handbuch der Technik und Methodik der Immunitätsforschung*, Jena, 1911, Erster Ersatzband, p. 80.

NOTES ON ASTHMA.

BY

JAMES ADAM, M.D.,

GLASGOW.

SEVENTEEN years ago I put forward the view that asthma is in most cases essentially a toxæmia. More than once in this JOURNAL I have given arguments for this view and have amplified them in my book, and details will not be repeated, but some of the criticisms of that book and much recent writing show that that view is still far from being widely accepted, and yet the proof that asthma is a toxæmia is based not only on the clinical facts but on the best of all evidence—therapeutical success. Of 390 cases about 70 per cent. can fairly be reckoned as cured, or almost so. These cases have come from all parts of this country as well as from some of the colonies, so that "place" has less influence on asthma than was believed.

Among these 390 cases two groups become prominent clinically. One group is that of week-end asthma. The asthmatic attacks in working-class patients tend to appear at the week end, and are clearly associated with the over-feeding and under-exercise of the week end. I have

repeatedly dealt with this point. The only possible explanation of this periodicity is the upset metabolism resulting in a form of toxæmia which induces asthma. This class of case is usually, and for obvious reasons, readily amenable to treatment. This week-end periodicity is not confined to asthma or to man. It is found in some forms of albuminuria, angioneurotic oedema, and epilepsy; also in the "Monday morning disease" of horses (Woods Hutchinson). The toxæmia in this group is interesting as helping to explain the next group.

The second large group shows a clinical history of bronchitis in children, recurrent year after year, until ultimately what is called spasmodic asthma is diagnosed. The patient may be adolescent or adult before he realizes that he has asthma. In this class eczema is a prominent concomitant. Treatment proves this eczema and this asthma to be due to coddling, to confinement, and to carbohydrate excess, chiefly in the form of milk puddings; for alteration of these conditions means immediate improvement and ultimate cure of the diseases. Here again we have an imperfect metabolism leading to toxæmia, which, in turn, reveals itself in catarrh—of the skin as eczema: of the respiratory tract as various forms of rhinitis, enlarged adenoids, laryngismus stridulosa, bronchitis. This toxæmia acts not only on skin and mucous membrane, but, sooner or later, on the nervous system. And so the catarrhal condition may be accompanied or followed by a spasmodic state, hence—sometimes in combination—urticaria, angioneurotic oedema of the mucous membranes, laryngismus, paroxysmal sneezing, asthma.

Two points are of great importance in this connexion. First, the nose as the respiratory orifice, and the most sensitive area of the whole respiratory tract. If the nose is not right the lower mucous membrane is sooner or later likely to become affected by spasm as well as catarrh: hence, in treatment, attention must be directed to the nose as well as to the toxæmia. Secondly, the widespread connexion of the vagus with the digestive as well as with the respiratory tract has ever to be kept in mind. It is possible the vagus may be directly affected by the toxæmia. All this points to the importance of dietetic treatment, which Hyde Salter so rightly emphasized. Treatment of these two classes of cases is usually soon and absolutely successful; but where the toxæmia has lasted many years, and the mucous membrane has become so altered that *restitutio ad integrum* is not to be expected, the most we can hope for is improvement, though even here the spasmodic attacks may be abolished. I have seen recovery after twenty, thirty, and even forty years of asthma.

Consideration of the circumstances under which asthma arises in domesticated animals goes to support the view that it is a toxæmia. The subject is disappointingly dealt with in British veterinary textbooks, but the account in the American one by Law (1896) is excellent. What he says as to the origin of asthma in dogs and horses exactly bears out what I have written here and elsewhere as to its origin in man. Asthma in dogs "is almost exclusively confined to those kept indoors, over-fed, without exercise, plethoric, and constipated." Everyone is familiar with the wheezy, asthmatic pet pug, but is apt to miss the fact that it frequently belongs to a pampered mistress—fat, wheezy, and asthmatic for a like reason. Law continues: "The disease is one of pet dogs, kept in towns, deprived of exercise, of fresh air, and of the opportunity to relieve the bowels at will; and gorged with highly spiced meats and sweets at least three times a day. Sluggishness and obesity are marked characteristics of the dog when first attacked, though in the advanced stages the violence of the paroxysms and their frequent recurrence may have induced extreme emaciation." And Mr. Dan Hamilton, M.R.C.V.S., informs me that eczema in dogs is caused in precisely the same way. Similarly, Law connects asthma in horses directly with the feeding and want of exercise. Veterinary surgeons have often told me that it is possible to induce asthma in almost any colt by improper feeding. For the following instance I am indebted to Mr. Begg, veterinary surgeon to Lanark County Council. The case is too long to give except in summary; but the details all support the views stated:

A six-year-old van horse, which had cost £46, was seen in January, 1903, on account of asthma, was put in a loose-box, fed on soft oleaginous rations and damped fodder. By the end of March he was reduced to a skeleton—weak, staggering, and

brought almost to his knees by paroxysms of coughing; value £3. He was turned out to grass in April, improved at once, and by July there were no signs of asthma. He was returned to work, but during the following winter, probably owing to the confinement and altered feeding, he developed a less severe attack of asthma. This passed off, and in July he was sold for £40, and was hale and working five years later.

It is likely that many diseases in animals, such as pleurisy and tuberculosis, are lumped together as asthma; but it is trifling with the subject to dismiss asthma in birds as aspergillosis. If the disturbed metabolism arising from over-feeding and under-exercise is the cause of asthma, the bird in which we should most expect it is the canary; and canary breeders are quite familiar with what they rightly call asthma. A bird has naturally a more active respiration and metabolism than man; but a caged canary can get no exercise, and has little to do but feed.

A canary breeder got an asthmatic canary from a neighbour to destroy. It had been fed *ad lib.* on all sorts of seed and other food. He himself had been cured of asthma many years ago, and instead of destroying the bird, thought he would try a plan similar to that I had laid down for him. Instead of filling up its dish with a day's food, he gave it small regular feeds of plain canary seed, and thus cured it. I have had many asthmatic canaries examined by pathologists, who have found tuberculosis in none.

Asthma is probably a disease of civilization. While man had to hunt and kill what he ate, asthma was not likely to arise. An asthmatic visited a Red Indian location in Canada. An old squaw said to him: "You breathe heavy." "Yes; I have asthma." "You English eat too much." "What would you have me do?" She held up two fingers: "Two days no eating." Which was exactly the advice I had given, and which elicited the story. Inquiry among medical friends in various parts of the globe has elicited a number of interesting facts which, however, require sifting, which cannot be detailed here, but of which the following is a summary:

In Central Africa, still uncivilized, there is no overcrowding, the diet is mixed; tuberculosis, asthma, and enlarged tonsils and adenoids are unknown. In China and Manchuria, while the day is largely spent in the open air, overcrowding is common at night; the diet is largely carbohydrate (rice), but no milk is drunk; teeth are often filthy; asthma and enlarged tonsils and adenoids are rare, tuberculosis of glands and lung common. In Siam the life is rather more indoors, the diet more mixed, and milk is drunk and betel chewing common; teeth are filthy; asthma and enlarged tonsils and adenoids not uncommon; tuberculosis greatly increasing. In India conditions vary in different parts and among the different races, but carbohydrate forms a large proportion of the diet, and milk is drunk; tuberculosis is common, asthma also in some parts, but enlarged tonsils and adenoids do not seem to be common.

The inference seems to be that asthma tends to appear in proportion as the life is spent indoors, as there is overcrowding, as carbohydrate forms the staple diet, and as milk is drunk. This about milk is interesting, because my experience proves beyond doubt that asthma cannot be cured when much milk, raw or boiled, is taken. I have cured two cases of hay fever in adolescents who took large quantities of milk by simply cutting it down.

Asthma, then, being in many if not in most cases a toxæmia due to want of adaptation of food, not only to the needs, but to the biochemical peculiarities of the individual, its essential treatment is obvious, simple, but involving much painstaking trouble in every case. I have dealt with this elsewhere at length.

Some think that attention to the nose has been over-emphasized, but, for reasons given above, this is doubtful. After that the main thing is to procure elimination of toxins and prevention of further toxæmia; or, to use Guelpa's phrase, "Désintoxication et renouvellement des tissus." Long before the appearance of Guelpa's book the fact of week-end asthma and the voluntary fasting of the asthmatic during his paroxysm till it ceased had taught me the value of fasting, and it was practised with success. And Hyde Salter sixty years ago rightly took pains with the individual diet. But not enough attention has been paid to cookery. With an unusually irritable *vagus* it is important that the digestive organs be taxed as little as possible.

THE value of the estate of Sir Leander Starr Jameson has been sworn at £45,082 18s. 10d., the net personality being £42,747.

A SURVEY OF 1,064 OPERATIONS FOR THE COMPLETE REMOVAL OF TONSILS AND ADENOIDS,

WITH A CONSIDERATION OF THE POSSIBILITY OF A REDUCTION IN THE NUMBER OF OPERATIONS,

BY

EDWARD D. D. DAVIS, F.R.C.S. ENG.,
ASSISTANT SURGEON, NOSE, THROAT, AND EAR DEPARTMENT,
CHARING CROSS HOSPITAL.

THE consideration of the large number of operations performed for the removal of the faucial tonsils and adenoid growth, or hypertrophied pharyngeal tonsil, makes one pause to wonder whether all these operations are really necessary. It must be remembered that the tonsils, both faucial and pharyngeal, are normal structures present for some definite purpose in every child.

When the tonsils and adenoid growths are septic, diseased, or hypertrophied, so that their presence produces definite and harmful symptoms, every one will agree that they should be efficiently removed, but the predisposing and exciting causes of infection and enlargement of these structures should be reviewed before deciding that operation is necessary. The experience gained by a close study of 1,064 cases, in addition to those seen during seven years in charge of a London County Council school clinic for affections of the nose, throat, and ear, may throw some light on their etiology. The cases have been carefully examined before operation, they have been seen two or three hours after operation, again one week afterwards, and, finally, six months later.

Etiology.

The age at which children require operation varies, but I have not seen one case of congenital enlargement of the tonsils or adenoid tissue, and agree with Lambert Lack¹ in doubting its occurrence. The youngest case was aged 16 months, and there were only five under 2 years. Children operated on under 2 often have recurrence, and the best results are obtained when the operation is performed between the ages of 3 and 6.

The strumous or scrofulous child is a common victim of "tonsils and adenoids" and has a typical appearance with enlargement of the lymphatic glands, phlyctenules of the cornea, excoriation of the nostrils by nasal discharge; such children lack immunity to catarrhal affections and to tuberculosis. This type of child runs in families, several members of the same family are strumous, and there is a family history of tuberculosis, suggesting that the condition is due to a tuberculous predisposition, in addition to bad surroundings in a poor home in a large city. The removal of tonsils and adenoids in such a child, followed by a visit to the seaside with good surroundings and food, leads to a vast improvement in health.

The most marked cases of enlarged tonsils and adenoids amongst children occur in those in whom there is evidence of dirty, badly ventilated, unhealthy homes with their cheaper carbohydrate diet. In these cases parasites, particularly *Pediculus capitis*, are invariably present, and the fact that they cause enlargement of the superficial cervical glands makes one consider what influence pediculi have on the lymphatic tissue of the pharynx. To operate on these children and to send them back to their squalid surroundings is to court failure. The removal of tonsils and adenoids in these cases is almost as illogical as scouring the hair of patients to remove pediculi and immediately allowing them to use the same dirty, infected pillow and clothes. The percentage of well-cared-for children with good surroundings seen in private practice who require operation is much smaller than in the case of poorer children seen at hospital.

Cameron² has shown that excessive carbohydrate diet among the poorer classes, which is the rule owing to its cheapness, produces a catarrhal tendency. Again, there is evidence to indicate that infection—for example, the tubercle bacillus—is conveyed to the tonsils by contaminated water, milk, and food.

The effect of locality is difficult to estimate, and at one time it was found that a large number of the cases came from the low-lying districts of the river banks, and from a valley town where there were watercress beds and a stream running through its centre, but statistics of this