elevation of the pelvis is attained (Fig. 3). The weight of the pelvis is transferred to the pelvic band sufficiently to enable the middle portions of the mattress to be withdrawn without disturbing the pelvis. The sacrum may then be washed or the bedpan introduced without movement of the pelvis, and, consequently, without disturbing its relations with the femur. The elevator is so arranged that it may be wrought by the nurse or by the patient. The latter may assist himself while lying supine by elevating or lowering his pelvis, and so adjust the level to suit his convenience (Fig. 4).

Do not overdo the elevation of the pelvis, as an over-enthusiastic nurse might do; just elevate sufficiently to suit.

Such pelvic elevators may be used for many other purposes: For elevating the pelvis during the application of bandages or surgical dressings; washing and cleansing the patient; for the prevention of bedsores and for the dressing of them when they have occurred; for fracture dislocation of spine, paraplegia, and for bedridden persons who have few attendants, especially as many of the patients could thereby assist themselves by working the elevator to any desired extent. The pain or aching in the back so often complained of during the first days after operation, when patients are compelled to lie supine, can be greatly relieved by the pelvic elevator, which can be adjusted and readjusted, raised and lowered as often as required.

It may also be employed as an elevator for any

FIG. 4.—Pelvic elevator sustaining weight of patient: centre piece of mattress withdrawn.

other part of the body, and, with the addition of extra slings, may act as a support for an arm or a leg. When not in use it may be employed as a cage for keeping the weight of the bedclothes from a fractured limb, or as a fracture cradle.

The divided mattress, as illustrated, was introduced by me in the early seventies for patients who had femoral osteotomies performed on them on both limbs at one time, and in whom it was necessary to main-

tain the limbs rigidly in proper relation with the pelvis until ossification was completed. Since then it has been in use for patients with fractures of the femur, and for other injuries and ailments. It has been found of much convenience to the nurse and comfort to the patient.

The pelvic elevator in one form or other has been used by me for a considerable period, but the newer, lighter forms here depicted have been introduced of recent years. One of these, made of wood in forty-eight hours by the mechanic at Mount Stuart Hos-

pital, Bute, has served its purpose very well.

Desiring, however, to get an efficient and cheap mechanism, stout iron rod was used, the superstructure being made of a single rod bent as illustrated, with a joining plate at the top, where the arches meet, through which the elevating screw goes.

Mr. Hilliard carried out the instruction, and, as shown in Fig. 2, he has introduced a modification which occurred to him, and which allows of the collapse of the mechanism for carriage to a distance.

Further Observations

ON

THE HAEMAGGLUTININ REACTION,

WITH SPECIAL REFERENCE TO THE OPERATION OF
BLOOD TRANSFUSION AND TO NEPHRITIS.

BY

C. J. BOND, C.M.G., F.R.C.S., COLONEL A.M.S.,

HONORARY CONSULTING SURGEON, NORTHERN COMMAND; HONORARY
CONSULTING SURGEON TO THE LEICESTER ROYAL INFIRMARY;
MEMBER OF THE MEDICAL RESEARCH COMMITTEE.

In the *BRITISH MEDICAL JOURNAL* of December 8th, 1917, I drew attention to the use of the haemagglutinin reaction as a method of testing the toxicity of different antiseptic reagents to human blood serum.

A further study of the haemagglutinin content of the blood serum in a series of 250 patients when tested with my own washed red corpuscles and those of the sheep has shown:

I.

That different human serums vary very much in their haemagglutinin content when tested with the same standard red cells. This grading of haemagglutinative capacity is independent of the specific power of different serums to agglutinate the red cells of different individuals according as they fall in one or other of the established groups. Individuals whose serums belong to the same agglutinative group differ in haemagglutinative capacity. Some have a very low and some a high capacity for agglutinating the same washed red cells. The result of the serum test has been recorded on the same plan in all cases; thus +++ indicates a high, ++ a moderate,

and + a slight degree of haemagglutinin effect, while -- indicates the absence of any reaction. In all cases the macroscopic method was used (see *BRITISH MEDICAL JOURNAL*, December 8th, 1917).

This means, of course, that incompatibility of donorship in the operation of blood transfusion is also a graded character, and it is therefore desirable, when testing incompatibility, to express not only the presence but the degree of agglutinative reaction. I have also previously shown that 1 part of non-agglutinative to 4 parts of agglutinative serum is sufficient to neutralize haemagglutination *in vitro*, but before applying this fact to the human subject it is desirable to ascertain by experiment on animals whether the same holds good in the living body. But this grading of agglutinative capacity is also important from the biological point of view.

It is supposed that the familial incidence of a haemagglutinative capacity of blood serum in parents and children follows Mendelian laws. It is, however, evident that a character—for example, haemagglutinin content—which varies from zero to a high positivity is much more difficult to handle as a method of testing inheritance than a sharply defined character of more constant value.

II.

But not only does the specific haemagglutinative capacity vary in degree in different serums, it also varies from time to time in the serum of the same patient. This fact is even more important than the first. I mentioned in my earlier communication that there were reasons for thinking that a non-agglutinative serum may become agglutinative in a patient who has passed successfully through a systemic infection. As the result of a study of 140 patients whose serums have been tested against my own red cells before

and after recovery from wound infection, I have records of 13 cases in which, on a subsequent test, some change was detected in the haemagglutinin content of the blood serum. In all cases the change was either from a negative to a positive reaction, or from a low to a high degree of haemagglutinin content. This generally coincided with a healthier condition of the wounds and with improvement in the general health of the patients.

Nephritis.

I have also tested the haemagglutinative capacity of the blood serum against my own red cells and those of the sheep in 50 cases of nephritis. Of these a second examination has been made, after an interval, in 25 cases, and a change in haemagglutinative capacity was observed in 15. In 7 cases a negative was converted into a positive reaction, and in 8 from a lower to a higher haemagglutinin content as far as my own corpuscles were concerned. With the sheep's corpuscles 5 cases showed a change from - to +, and one from a lower to a higher degree of agglutinative capacity. In three cases only was the change in the other direction. In the first of these a ++ was converted into a - reaction with my corpuscles, the action with the sheep's corpuscles remaining the same—namely, negative. In the second the reaction with my corpuscles remained constant, namely, ++, but dropped from a positive to a negative reaction with sheep's cells. In the third case the human cell reaction fell from ++ to +, but with the sheep's cells the reaction rose from + to ++. The two blood examinations were made in nearly all cases at about a month's interval, and most of the patients were improving generally, with a diminution in the amount of albumin in the urine.

The interesting problem arises in these albuminuric cases whether any leakage of haemagglutinins goes on from the kidney along with the leakage of serum-albumin and globulin.

A somewhat extensive examination of the action of these albumin-containing urines on washed red cells failed to show any distinct evidence of haemagglutinative effect. Certain changes, however, occur in the red cells which I regard as initial stages in agglutination, but the reaction does not become complete owing possibly to physical or biochemical factors in the urine. A control experiment in which a known volume of agglutinating serum was diluted with urine in different proportions showed that distinct evidence of agglutination could be obtained in serums diluted up to 1 part in 10 with urine, even after standing for twelve hours. Different samples of healthy urine also vary in the degree to which they destroy the agglutinative capacity of a given serum. This variation in the urine of different individuals to degrade the blood serum is undergoing further investigation, and may possibly provide a test of urinary toxicity.

Patients suffering from albuminuria consequent on nephritis show, on the whole, a lower haemagglutinin content of blood serum than is found in a corresponding group of partly healthy persons. A series of fifty cases of soldiers suffering from nephritis and a series of fifty male patients whose blood was examined for the Wassermann reaction, in many negatively, gave the following results. The cases are arranged in series showing the haemagglutinative strength:

	+++	++	+	-	Total.
Nephritis cases ...	3	16	20	11	50
Wassermann cases...	13	12	22	3	50

If this reduced haemagglutinin content in nephritis cases is considered in conjunction with the rise in haemagglutinin content which takes place as the albuminuria diminishes and the patients recover, we are led to the conclusion that a leakage not only of serum-albumin and globulin but also of haemagglutinin does take place from the kidneys in nephritis. This is important, because if the blood serum is drained of haemagglutinin it may also be drained of other substances of the antibody group, and this loss may be associated with the lessened resistance to infection which is known to exist in these patients.

Conclusion.

In a considerable number of cases under conditions which require further analysis, but which seem to be associated with recovery from albuminuria in the nephritis cases, and successful resistance to general infection in the wound cases, a definite alteration has been found to occur in the blood serum, for the most part in the direction of an increased haemagglutinin content. The bearing of this fact on the problem of blood transfusion now becomes apparent. If the blood serum of a recipient may be at one time compatible with the red cells of a given donor, and at another time may agglutinate those cells, the donor's serum may also, as the result of recovery from infection or other illness, change in the same way.

Cases have been recorded by Major Robertson, Major Lee, and others, in which transfusion of blood from the same donor to the same recipient had no harmful effects on one occasion but was followed by death on a second. I venture to suggest that changes in the haemagglutinin, and probably in the haemolysin, content of the blood serum of the recipient or donor or both occurred in the interval between the two transfusions, which were responsible for the unfortunate result.

GENERAL ANALGESIA BY ORAL ADMINISTRATION.*

BY

CAPTAIN JAMES T. GWATHMEY, M.R.C., U.S. ARMY,

AND

CAPTAIN HOWARD T. KARSNER, M.R.C., U.S. ARMY.

THIS report is in the nature of a preliminary communication based on animal experiments and a sufficient number of clinical cases to support the conclusions of the animal work. It will be seen that general analgesia in which there is loss of sensation, with or without loss of consciousness, can be established for otherwise painful dressings and for short operations. The desirability of such a procedure becomes apparent immediately on entering a military hospital, and it has also a definite place in civil surgery.

Many war wounds are accompanied by fractures of bones, and the importance of keeping such patients quiet is universally recognized. If the dressing of such a wound is accompanied by severe pain, it is customary at this base to produce nitrous oxide-oxygen or light ether anaesthesia for at least the first few dressings, necessitating in most cases, because of limitation of apparatus, removal of the patient to the operating room. This usually means pain and the danger of misplacing the bone fragments before and after dressings, as well as loss of time to surgeons, nurses, and orderlies. General analgesia, produced simply and quietly without taking the patient from his bed, is the logical solution of this difficulty, and the technique evolved as the result of this study makes it possible to administer the analgesic by mouth in perfect safety. It is, of course, applicable to practically all forms of painful dressings, and is being developed to embrace short surgical operations, such as resection of a rib, removal of foreign bodies, revision of wounds. In such operations, however, it may be necessary to supplement the analgesia with novocain for the skin incisions, a hypodermic injection of morphine, or even by light inhalation anaesthesia.

Local conditions made it seem advisable to select rabbits as the animals for the preliminary work. The animals were tied to a board and the various substances given by stomach tube, after which the animals were immediately released and placed under observation.

Quinine and Urea Hydrochloride (Nikalgin):

Weight of Animal.	Amount.	Result.
2200 gni. ...	4 c.cm. }	... No systemic effect.
2105 " ...	8 c.cm. }	
2250 " ...	16 c.cm. }	
1795 " ...	30 c.cm. }	
2215 " ...	60 c.cm. }	{ Unable to stand after 10 minutes and died in 8 to 17 hours.

* Aided by grants from the American Red Cross.