A Clinical Lecture ON THE BAD HABIT OF VAGINAL DOUCHING.

GIVEN AT THE MANCHESTER ROYAL INFIRMARY.

BY

W. E. FOTHERGILL, M.A., B.Sc., M.D.

THE inventor of the vaginal douche-can was no friend to womankind, for the result of his ingenuity has, on the whole, been a curse rather than a blessing. Just as there was a pessary age, characterized by the haphazard use of pessaries in the treatment of the minor ailments of women, so there has been a long period of indiscriminate douching which has not yet reached its belated end. Soon after the discovery of micro-organisms as a cause of disease, the newborn infection idea began to put out of joint the nose of its little elder brother the displacement theory, and it was surmised that infection must be to blame for most of the minor pelvic troubles. Less was heard about versions and flexions, and those blessed words "endometritis" and "ovaritis" came into fashion as diagnoses. Less ingenuity was bestowed upon the adaptation of pessaries, and much trouble began to be expended on the disinfection of the vagina by douching, regardless of the fact that it was seldom infected. Not only was the douche used for the cure of supposed infection, but also for its prevention. For some time it was even the custom to swill the vagina with antiseptic lotions before, during, and after labour. The results were so deplorable that this practice was soon dropped. I well remember being told by the late Professor Sir Alexander Russell Simpson, in the year 1895, to tear down from the walls of the Edinburgh Maternity Hospital the instructions for routine douching which had been in force for a period during which the hospital had to be closed more than once on account of epidemics of puerperal pelvic infection.

For a time douching remained under the control of the medical profession; but then the general public took the bit in its teeth and began buying douche-cans on its own account. It has gone on doing so ever since, and the pernicious but apparently seductive habit of antiseptic douching is now generally begun on the advice of an elderly relative. A patient, now aged 43, began douching herself with alum water at the age of 19, on the advice of her stepmother. She had the "whites" at the time, and the alum naturally made the complaint worse. She married and went on douching, but changed from alum to Condyl's fluid on the suggestion of her mother-in-law. For the last fifteen years she has douched herself night and morning; and after trying various other irritants, is now using a lysol substitute which makes her smart badly. The vulva, vagina and cervix are now "red raw" and streaming with discharge. She has had menorrhagia and dysmenorrhoea for several years, and recently has suffered continuous pelvic discomfort.

A healthy young lady mentioned to her mother-in-law that she had a slight excess of normal vaginal discharge just before and after the menstrual periods. She was provided by this relative with a douche-can, which she used every day for two years, the "whites" becoming worse and worse all the time. The discharge became constant and slimy in character, the menstrual flow was increased in duration and quantity. The patient began to have congestive dysmenorrhoea, and later there was dull aching pain throughout the intermenstrual intervals. When the douching was stopped the symptoms gradually disappeared, the last first and the first last. Cases like these are not exceptional; they crop up every day both in hospital and in private practice. Indeed, every patient who complains of minor pelvic ailments should be asked whether she douches or not; and, if she says "Yes," should be told to stop it. Nurses often douche without any instructions; sometimes because they think it is the correct thing to do, and at other times because their patients seem to expect it. This must also be stopped.

ANTISEPTIC AND MEDICATED DOUCHES.

In considering the action of douches we must remember that the vaginal surface has no glands, and is not a mucous

membrane, but is covered by stratified squamous epithelium, which, though very strong and resistant to infection, is, so to speak, not quite waterproof. Thus the vaginal secretion is simply serum which exudes from the surface of the squamous epithelium, and which acquires its normal creamy appearance by admixture with leucocytes and epithelial debris. Sliminess of the vaginal discharge is due to an excess of mucus which comes from the lining of the uterus and not from the vaginal wall. Again, it must be remembered that the normal vaginal secretion is definitely acid in reaction, and that this acidity inhibits the life of most septic and pathogenic organisms. Thus the normal content of the vagina is a barrier to ascending infections, and it is generally sufficient to protect the uterus. For instance, after the experimental introduction of streptococci, the healthy vagina has been shown to be free from these organisms in the course of a few hours. In fact, most vaginal infections quickly cure themselves if left alone, and many cases cure themselves in spite of a certain amount of vaginal douching, which, in these cases, secures the credit that should by right belong to the unaided vagina.

But douching does actual harm in more ways than one: for it washes away the normal acid secretion; it kills the acid-producing bacteria which are the normal inhabitants of the vagina; it kills the superficial layers of cells, and irritates the subjacent layers. Further, it promotes hyperaemia of the parts, and thus increases the quantity of the secretion, favours menorrhagia, congestive dysmenorrhoea, and intermenstrual congestive pain. This explains the fact that the woman who douches much never gets rid of her vaginal discharge, and generally has other complaints in addition.

The above remarks refer to chronic douching; are there any excuses for what might be called acute or subacute douching? Take septic infection of the uterine cavity after abortion or labour. In these cases vaginal douching clearly does not touch the spot at all, and intrauterine douching is positively dangerous, as it sometimes causes fatal peritonitis by forcing septic material and noxious antiseptics through the tubes into the peritoneal cavity. The proper way to treat uterine sepsis is to dry-clean the uterine cavity thoroughly, using a general anaesthetic if necessary; to apply concentrated antiseptics with no excess of fluid; and thereafter to leave the uterus and vagina severely alone. Take gonorrhoeal infection of the vulva. If it is treated by douching, the infection is very likely to be carried up through the vagina to the cervix, whence the gonococcus has a clear run over the mucous lining of both uterus and tubes. Again, when the cervical canal is once infected, douching is futile, and proper disinfection under anaesthesia is the only method likely to be effective. In fact, there is no "home treatment" for uterine gonorrhoea in women. Yet there are still some who supply their patients with the inevitable douche-can and order copious and frequent libations of Condyl's fluid, while in one clinic women are actually advised to wash out their own bladders!

Though it appears that antiseptic and medicated douches might well be dispensed with altogether, there are exceptions to the rule. Thus, in the palliative treatment of cancer of the uterus, disinfectant and deodorant solutions must be used. Again, when preparing a prolapsed and ulcerated vagina or cervix for operation, the nurse will give one or two mildly antiseptic douches every day for a week or ten days, during which the patient is kept in bed and the vagina lightly packed. After vaginal operations, also, a few antiseptic douches are useful on those rare occasions in which the incisions become septic. In such cases the indication is definite. But often there is no indication at all. The patient says, "Doctor, I have a vaginal discharge." The doctor makes no examination, but says, "Get a douche-can," and orders alum, iodine, lysol, mercuric perchloride, boracic acid, zinc sulphate—anything, in short, that occurs to him. The patient begins douching, the doctor forgets to stop her, and she goes on with it for years. The idea that a vaginal discharge must be infective is very widespread and persistent, though it may be far from correct. When a woman complains of a discharge it is most frequently "the whites," namely, excess of the normal creamy vaginal discharge. This is common before and after menstrual periods in perfectly healthy persons. Its persistence throughout the intermenstrual period is often an expression of a low general condition;

but some women have more vaginal secretion than enough throughout life, and quite apart from variations in health. Any form of local treatment aggravates "the whites," but the condition often disappears with improvement in health brought about either by general treatment or by change of air and scene. When the vaginal discharge is slimy, the uterus is secreting too much mucus, and this is increased by douching or any other local treatment of the vagina. If any local treatment is required it is curettage of the uterus; but this generally fails unless the general muscular and vascular tone is improved at the same time.

HOT DOUCHES.

Hot douching is often ordered in cases of menorrhagia and other kinds of uterine bleeding. It is true that the application of water at a temperature of about 120° F. contracts the blood vessels and leaves them in a state of contraction which lasts for from half to three-quarters of an hour. But water which is hot enough to produce this effect is uncomfortably hot for the skin round the vulva and anus as it flows out of the vagina. It is possible, given a strong-minded nurse and a weak-minded patient, that really hot douches may sometimes be given for several days together. But in practice, when the hot douche is ordered, it is given warm; and water at 100° F. increases pelvic hyperaemia and so increases uterine bleeding. Thus it happens that when hot douching is ordered for menorrhagia, the patient really receives treatment by warm douching, and the bleeding gets worse instead of better. The hot douche, therefore, should probably be kept for use in obstetric haemorrhage, in which it is doubtless very useful on occasions.

WARM DOUCHES.

Under certain circumstances it is highly desirable to increase pelvic hyperaemia, and for this purpose the prolonged warm douche is useful if given under proper supervision and for a limited period. The typical case for this treatment is a patient who is recovering from pelvic cellulitis following infection of a torn cervix. There is a more or less massive deposit of inflammatory exudate in the pelvic cellular tissue. Its absorption must be favoured in order to avoid permanent fixation of the pelvic organs. The same holds true also in reference to inflammatory deposits in other pelvic tissues and organs. Thus, in the subacute phases of pelvic infection, vaginal douching is seen at its best. A slow stream of normal saline solution at a temperature of 100° F. to, say, 105° F., kept up for ten or twelve minutes once or twice a day for two or three weeks, forms a part of the proper treatment of these cases. But it must be done with the patient lying on her back, for when a douche is given with the patient sitting up the flow of water never separates the vaginal walls, but simply runs out beside the douche nozzle without even washing the vaginal surface. To act as a poultice or fomentation the water must balloon the vagina, and this it does only when the patient is on her back.

CONCLUSION AND SUMMARY.

You may have noticed that some people cannot learn at all, that some can learn only by their own experience, and that some are sufficiently astute to profit by the experience of others. To those of you who care to try the results of my experience I offer the following practical hints:

1. Do not allow patients to douche themselves at all; but, when you have a definite reason for douching, let a nurse do it.
2. Do not use antiseptic douches unless there are germs to be killed in the vagina.
3. Remember that vulvitis is common but vaginitis is rare.
4. When you wish to apply antiseptic lotions to the vulva tell the patient not to douche herself with the lotion but to put it in a large bowl and sit in it.
5. Do not order hot douches (120° F.) in cases of menorrhagia and metrorrhagia.
6. Do not douche for leucorrhoea either creamy or slimy.
7. Do not douche for aches and pains in the pelvis which are not due to pelvic infection.
8. Use warm douches (100° to 105° F.) as you would use poultices or hot fomentations, in cases of pelvic

infection which are neither too acute nor too chronic to be improved by the production of temporary local hyperaemia.

THE TREATMENT OF FURUNCULOSIS AND OTHER DEEP-SEATED COCCOGENIC INFECTIONS BY COLLOSOL MANGANESE.

By SIR MALCOLM MORRIS, K.C.V.O.

THE routine treatment of boils and other deep-seated cocco-genic affections, such as ecthymatous impetigo and some cases of acne, is often most disappointing. In the treatment of furunculosis some authorities lay stress upon constitutional treatment; and in certain cases, as where the process is associated with diabetes, this is obviously important. Others rely rather upon the local application of antiseptic agents, with the view of absorbing the boils. But both forms of treatment frequently fail. In many instances boils to boils succeed, and when at last the dismal procession ends, this often appears to be less the result of treatment than because the disease has run its natural course. This is true of vaccine-therapy, as well as of the earlier methods. Sometimes the inoculations appear to arrest the disease; in other cases they signally fail to do so.

In these circumstances it seemed to me desirable to give a trial to the treatment by collosol manganese instituted at the General Hospital at Etaples by Mr. J. E. R. McDonagh.¹ The results in all my cases are so remarkable as, in my judgement, to demand publication.

It should be premised that at first the collosol manganese (Crookes) was used in the form of a single solution. Afterwards it was employed in the improved form of two solutions, which had to be mixed in the syringe; of this, much smaller doses (about one-half) suffice.

CASE I.

The patient, a medical man, came to me on November 24th, 1917, with large, deep-seated boils on the hairy part of the face, which he believed to be an infection from a carbuncle in a diabetic patient, from whom he had contracted a septic whitlow. Culture showed that the micro-organism was the *Staphylococcus pyogenes aureus*, and an autogenous vaccine was prepared, with which he was inoculated on November 30th. By this time a boil on the right cheek had enlarged to the size of a Tangerine orange; there was much oedema of the face and eyelids and great pain. On December 6th pus was discharged and the pain relieved; but during the next seven weeks, boils continued to appear at intervals, usually in groups of four or five, on one or both sides of the face. By January 14th eleven injections of vaccine had been given, antiseptic lotions and ointments being also applied. On January 31st the patient consulted me again and I advised the injection of collosol manganese. The first injection was given the next day. "Improvement," to quote from the patient's own notes, "began on the third day afterwards. Subsequent boils, which continued to appear for a further fourteen days, were not painful, and many of them aborted without pus formation. In all, four injections were given—1.5 c.cm. on January 31st, 2.5 c.cm. on February 5th, 3 c.cm. on February 12th, and 1 c.cm. on February 28th; the first and second in the gluteus, the third and fourth in the flank. In the first three the old preparation was used; in the fourth the new. By March 5th I was able to report 'All clear.' From November 30th to January 31st, while the vaccine was being given, thirty-three carbuncular boils appeared; after the collosol manganese was begun there were only seven, the last appearing on February 20th, three weeks after this treatment was begun." The patient's general health also greatly improved; he began to feel better within a week of the first manganese injection.

CASE II.

The patient was an army captain who was subject to acne. When the first collosol manganese injection (0.5 c.cm. of the new preparation) was given, on March 13th, he had been in hospital four months with facial boils and acute impetigo contagiosa of the same region. During that time he had had vaccine injections, and various antiseptics had been applied, without result. Another manganese injection (0.5 c.cm.) was given on March 16th, and a third (1 c.cm.) on March 19th. By this time all the lesions had cleared up.

CASE III.

I saw the patient, a lady, for the first time on February 16th. She had extensive follicular impetigo of the scalp, which began in the preceding October, and had gradually become more severe. When she came to me she complained of great pain from small multiple deep abscesses. There was a good