iliac fossa pointed to it being appendicular, but the history that the patient gave of bile draining from the upper wound for several months suggested that it either had something to do with the gall bladder or the second part of the duodenum. From the huge dilatation of the common bile duct it was concluded that there must be some blockage of the common bile duct, and this could only have arisen from an impacted gall stone or from a perforated duodenal ulcer. The most likely explanation of the condition was thought to be that there had been a perforated duodenal ulcer the cicatrix of which had constricted the common bile duct. The local inflammation around the gall bladder had at the same time involved and obliterated the cystic duct, so that the gall bladder had not yielded to the pressure of the secreted bile.

A letter from Dr. Sattler of Bremen subsequently confirmed these surmises. He stated that he operated upon the patient on June 12th, 1912, for a perforated duodenal ulcer with general peritonitis. The surroundings of the ulcer were very necrotic and there were many adhesions to the stomach and gall bladder; the peritoneal cavity was full of pus and bile, so that a counter-opening for drainage was made in the right iliac fossa.

Thus it was evident we were dealing with a case of constriction of the common bile duct following a perforated ulcer of the duodenum. As there was no bile entering the intestine it was decided to endeavour to construct a new channel for it by making an opening between the dilated common bile duct and the small intestine.

Second Operation.

On January 23rd, 1913, a vertical incision was made through the outer margin of the right rectus muscle. The sinus was dissected out and traced to the dilatation of the common bile duct. The latter had contracted down markedly since the previous operation. The sinus was cut away and the opening in the bile duct so made was clamped. The bile duct was freed and the clamp brought as far forward as possible, but owing to the density of the adhesions and the contraction of the bile duct since the previous operation very little relaxation could be obtained.

A loop of jejunum was then brought up through the transverse mesocolon and an anastomosis was made between this loop and the dilated bile duct by two rows of silk sutures. This was rendered difficult by the fixity and the depth of the "duct" and the friability of its walls. The sutures were unevenly placed and there was considerable tension on them. When completed the opening would admit two fingers. Owing to the fixity of the bile duct it was impossible to draw back the loop of jejunum through the aperture in the mesocolon; hence it was found necessary to perform a lateral anastomosis between the afferent-efferent portions of the loop on the under surface of the transverse mesocolon in order to prevent intestinal obstruction.

A cigarette drain was put in the region of the choledoch-enterostomy, and the abdominal wall was closed by through-and-through silkworm gut sutures.

The patient stood the operation extremely well, owing, no doubt, to his greatly improved general condition.

After-History.

Two days after the operation, on the removal of the cigarette drain, bile was noticed coming along the track, but as the faeces contained bile it was obvious that some bile was finding its way into the intestine.

The condition of the patient at this time gave rise to anxiety. He was vomiting freely, the temperature was 101° F., and the pulse 120. A drainage tube was again put in, and shortly afterwards, besides bile, the discharge contained digestive juices. The skin around the sinus rapidly became excoriated, but this was controlled by bismuth and paraffin. The patient became emaciated, the faeces were pale and fatty, and the appetite poor.

After six days the amount of bile and digestive juices escaping gradually diminished and the appetite rapidly improved. The stools became darker and he began to put on weight.

At the end of a fortnight the discharge ceased, the sinus rapidly closed, and the faeces became normal in colour and content.

On March 3rd, when the patient was discharged, he looked extremely well. The abdominal wall was quite strong save at the right margin of the transverse incision where the drainage tubes were at the first operation.

The size of the hernial opening and the difficulty of dealing with it suggest the inadvisability of transverse division of the recti in suppurative conditions which preclude adequate suture.

Although a sinus obviously formed at the site of the choledoch-enterostomy owing to the difficulty of adequately suturing the anastomosis, it healed rapidly and did not interfere with the efficiency of the new passage.

An account of constriction of the bile ducts following

the cicatricial contraction of a duodenal ulcer is given by Moynihan in *Duodenal Ulcer* (1910). He describes two conditions:

1. When the ulcer was situated near Vater's ampulla, and constricted both the common bile duct and the pancreatic ducts.

2. When the common bile duct alone may be involved as it runs behind the first and second parts of the duodenum.

Details of eleven cases of the first class are given, and of five cases of the second class, of which one was his own. No case is recorded in which the constriction followed a perforation of a duodenal ulcer. The cases recorded are all of the very chronic type of ulcer, in which there is marked induration around the base.

The case described here falls in the second group of cases, in which the common bile duct alone was involved. Since the ulcer perforated into the peritoneal cavity, and not into the retroperitoneal tissue, the ulcer must have been situated in the free part of the duodenum—that is, in the first or at the junction of the first and second parts.

The fact that bile was present in large quantities in the peritoneal cavity suggests that the ulcer had perforated through the bile duct in addition to the posterior wall of the duodenum. Whether the bile duct was caught in the sutures which were used to close the ulcer, or whether the constriction followed the cicatrization of the ulcer, is not clear. The fact, however, suggests that in these cases great care should be exercised in suturing the perforation lest the bile duct be involved.

In this case it was impossible to perform an anastomosis between the bile duct and the duodenum owing to the numerous adhesions around the latter. The comparative ease with which the anastomosis can be performed by bringing a loop of jejunum through the transverse mesocolon, and the perfectly satisfactory result ultimately obtained, points to choledoch-jejunostomy being the operation of preference in these cases.

My best thanks are due to Mr. E. Rock Carling for permission to publish this case, and also for the kind assistance he has given me in reporting it.

A CASE OF HAEMATIDROSIS.

BY

CHARLES T. SCOTT, M.B.CANTAB.,

MARKET HARBOUROUGH.

THE following case was brought to my notice by Dr. H. Adcock, of Middleton, Northants, in conjunction with whom these notes are written.

Madeline —, aged 11 years, a bright intelligent girl of rather nervous temperament, commenced to have abnormal perspirations about four months ago, following a "chill on the liver"; since then they have been gradually growing more and more frequent and severe.

The sweats occur upon the forehead only, from the eyebrows to the roots of the hair, and as far as the outer margins of the orbits laterally; they consist of either clear watery fluid or white froth or bright pink fluid. At the present time they recur at intervals of from a minute to an hour, but during waking hours there is rarely a lapse of more than five to ten minutes, and often there are as many as eight "sweats" in ten minutes.

The "frothy" are the most common variety, accounting for 60 of a counted consecutive hundred; the red variety formed 34 per cent. and the watery 6 per cent. They are most frequent soon after waking in the morning and towards bedtime. The quantity secreted at each "sweat" is from 10 minims to 1 drachm; the secretion invariably appears rapidly, so that in a second the forehead is bathed in sweat, and occasionally it is so rapid that it can be seen as it issues from the forehead in a fine jet and can be heard upon the bedclothes. There is a distinct aura, which takes the form of a faint "sick feeling" in the epigastrium, more severe when a coloured sweat is heralded, so that the child will say, "A pink one is coming now," or "Now there is a frothy one," and her prognosis is invariably correct.

There are no sweats during sleep, but she is frequently awakened by one coming on; as a rule she sleeps for some hours at night with no sweats, but her rest is always disturbed by waking for sweats.

During the last few weeks the salivary glands have been affected in a similar manner to the sweat glands of the forehead, and the same aura may herald a sweat or a "mouthful," which latter may also be pink or white.

For the last fortnight she has had attacks of tonic spasm of the muscles of the arms, so that the hands may become rigid in the "accoucheur's position," or she may involuntarily grasp some object, or the hands may be raised and tightly pressed against the back of the head, causing pain in the back and sides, suggesting hysteria. She has suffered from almost continuous headache, frontal and occipital, since the sweating began. She has had no vomiting.

From a scraping of the skin of the forehead staphylococci and streptococci were grown in small numbers only. Histological examination of the sweat shows red and white blood corpuscles in all three types of sweat, but far more in the "pink" sweat than in the others. Bacteriological examination shows staphylococci, a few streptococci, and a Gram-negative bacillus, apparently of the Friedländer group.

The patient was a premature child (eight months); her mother was badly frightened shortly before the birth by being stabbed in the head by a burglar. She has always been addicted to walking and talking in her sleep, but has had no serious illness. A few years ago tenotomy was performed upon both eyes for strabismus. She has not shown any signs of menstruation. There is no tendency to haemophilia in the patient or her family.

Just a week before the sweats commenced she had a very severe shock. Being particularly nervous of air raids, her parents tried to keep her out of their way; one day a gas explosion occurred in the next house while the child was in bed; she was frightened by the noise.

Examination reveals no abnormal signs in the nervous system, heart, lungs, or digestive system. The urine is copious, up to 6 pints daily, is acid, sp. gr. 1020, and contains no albumin or sugar. The reflexes are very brisk and the fundi healthy.

The case would appear to be one of an unusual type of functional disturbance consequent upon the shock of the explosion. It has not hitherto proved amenable to treatment.

INTERMITTENT HAEMATEMESIS FOLLOWING ON AN INJURY TO THE CHEST.

BY

CAPTAIN A. H. D. SMITH, M.B., CH.B. EDIN., M.C.,
ROYAL ARMY MEDICAL CORPS.

THE interesting point about the following case is that there were no symptoms of any sort until the patient was injured in 1915. The haemorrhages have all taken place after excessive exertion. Apart from the discomfort and fright caused by the haemorrhages he is in no way handicapped by the condition.

History.

Private K., aged 39, consulted me on March 4th, 1918, on account of haematemesis. In October, 1915, when a cavalryman, he was kicked in the precordial region by a horse. He attended hospital for eight weeks as an out-patient. As far as he knows the ribs were not broken, but he was in strapping for the whole eight weeks. He was said to be suffering from a severe contusion of the chest; there was no vomiting or spitting of blood after the injury, and beyond pain there were no symptoms of any sort; previous to the injury he was quite well, and had been training for service overseas since June, 1915.

Before enlistment he was in extensive business as a book-maker. Both his parents are alive and well; there is no history of heart disease, lung disease, or apoplexy in the family. He has three children alive and well; his wife is healthy, and there have been no miscarriages.

The patient's history is as follows: He has always enjoyed excellent health; there is no specific history. For many years he was a heavy spirit drinker, on occasions drinking as much as one and a half bottles of whisky a day, and he averaged three-quarters of a bottle of whisky daily for ten years. Some three years ago he consulted a medical man, who told him he had a diseased condition of the liver, and advised him to give up spirit drinking; since then he has been for all practical purposes a total abstainer.

Since the injury he has suffered from intermittent precordial pain; about every six months he has had attacks of haematemesis. The attacks come on suddenly without warning, usually during violent exertion. About half a teacupful of blood comes away with a gush; there is no pain at the time

but for a short time after the vomiting he has difficulty in breathing. In the intervals between the attacks he is quite well, but since the accident he has suffered from insomnia.

On February 27th, 1918, while walking up a slope against a very strong wind, he found difficulty in breathing, and blood suddenly rushed into his mouth. Again, on March 3rd, while walking sharply uphill, the same thing happened, only in this case the blood was clotted. He was short of breath for some time after the attack, but was able to walk four miles home and carry on his work next day.

Physical Examination.

Respiratory system: Nothing abnormal found; no evidence of old fracture of ribs.

Circulatory system: Nothing abnormal; no thickening of vessels; no varicose veins.

Nervous system: No objective symptoms.

Abdomen: Much tenderness on palpation over liver; this organ (by percussion) decreased in size; spleen also tender, but not as tender as liver; no history of jaundice; nothing abnormal found otherwise.

Genito-urinary system: No abnormal condition discovered; urine normal.

Haematopoietic system: Nothing abnormal.

Facilities for a more elaborate examination were absent at the time.

Differential Diagnosis.

The haematemesis has, of course, to be differentiated from haemoptysis. In this case there is no history of cough, or of pulmonary or heart disease. Apart from the injury in 1915 there is no reference to any chest trouble. The patient states that the blood is in his mouth before he has any warning, and that on March 3rd the blood was clotted.

The patient can give no information as to the presence of food in the vomit. There is no history of melaena. The absence of pain in the intervals helps to differentiate from gastric, duodenal, and oesophageal ulcers (peptic). The absence of wasting helps to differentiate from malignant disease. The absence of anaemia is a strong point in excluding ulceration of any sort.

There is no haemophilia or abnormal blood condition; there is no affection of throat or nose which might account for the haemorrhage. The history of excessive spirit drinking spread over many years (confirmed by condition of liver and spleen found on examination) points to the condition being secondary to the hepatic (or splenic) condition.

Treatment.

The patient has been advised to abstain from spirit drinking and lead a normal, active, healthy life, but to avoid all undue exertions, such as rushing upstairs and walking quickly up hills. He has been advised to take his meals slowly and not bolt his food, and to abstain from drinking very warm fluids. No drug treatment has been given.

I think the probable cause of the recurrent haemorrhages is a varicose condition of the veins at the lower end of the oesophagus. The absence of any arterio-sclerosis excludes a small aneurysm. The veins may have been injured by the trauma to the chest, and so an extra strain has been thrown on veins already weakened by cirrhosis of the liver and spleen.

I beg to acknowledge the kindness of Surgeon-General Julian, D.D.M.S. Western Command, for permission to publish this case.

THE PHAGOCYTIC RESPONSE TO THE INTRODUCTION OF BACTERIA INTO CLEAN WOUNDS.

BY W. JAMES WILSON, M.D., D.Sc.,

BREVET MAJOR R.A.M.C.(T.F.).

(From the Laboratory of — General Hospital, B.E.F., France.)

In the course of an extensive experience of the bacterial flora of war wounds I have been struck by the small numbers of *B. coli* which are present as the wound tends to clean. Even wounds of the buttock which are repeatedly contaminated with faeces contain after a few days comparatively few *B. coli* although they are swarming with enterococci.

At first I imagined that the explanation of the selective effect of the wound on faecal flora might be that the wound fluids offered an unsuitable medium for the growth of *B. coli*. No doubt this is an important factor, but the following few experiments show that the immunity is an active one, and that the phagocyte plays an effective part in its production.

Experiment I.—Fate of Living Culture of *B. coli* Introduced into a Clean Wound.

The patient (R.) had a wound (6 cm. by 5 cm.) of right forearm just below the elbow-joint. On November 23rd, 1917, the wound was granulating, and a swab taken on that date gave a growth