TABLE I.—Mortality of Nits of *Pediculus humanus* due to Immersion in Watery Solutions of Liquor Cresoli Saponatus Fortis.

Temperature and Period of Immersion.	Strength of Solution.	No. of Eggs.	No. Hatched.	No. Died while Hatching.	Percentage Mortality.
	Percent.				
Temp. 60-65° F.	2.0	94	3	7	97
5 minutes	1.5	51	26	9	49
	1.0	69	37	10	46
	0.5	46	41	2	11
	0.2	79	64	2	19
20 minutes	2.0	106	Nil	—	100
	1.5	82	1	—	99
	1.0	93	5	4	95
	0.5	68	24	10	65
20 minutes	0.2	37	30	3	19
	2.0	68	Nil	—	100
45 minutes	1.5	92	Nil	—	100
2 hours	1.0	64	Nil	—	100
2½ hours	0.5	99	Nil	—	100
22 hours	0.2	90	Nil	—	100
Temp. 48° F.					
20 minutes	2.0	95	Nil	—	100
	1.5	104	Nil	—	100
Control*		85	66	1	23
Temp. 32° F.					
20 minutes	2.0	137	1	1	99.3
	1.5	123	Nil	—	100
Control*		51	46		10

* The controls were exposed to the temperature, but not immersed.

TABLE II.—Mortality of Nits of *Pediculus humanus* due to Immersion in Watery Solutions of "Lysol" (equal parts of Crude Carbolic Acid and Soft Soap).

Temperature and Period of Immersion.	Strength of Solution.	No. of Eggs.	No. Hatched.	No. Died while Hatching.	Percentage Mortality.
	Percent.				
Temp. 60-65° F.	2.0	120	Nil	Nil	100
5 minutes	1.5	121	Nil	1	100
	1.0	82	2	6	98
	0.5	79	52	7	34
	0.2	89	70	2	21
20 minutes	2.0	89	Nil	Nil	100
	1.5	82	Nil	Nil	100
	1.0	72	Nil	1	100
	0.5	103	46	8	55
20 minutes	0.2	94	60	8	34
Temp. 48° F.					
20 minutes	2.0	96	Nil	—	100
	1.5	102	Nil	1	100
Temp. 32° F.					
20 minutes	2.0	127	1	—	99.2
	1.5	99	7	2	93

in the latter case is not great, and may possibly be largely, if not entirely, a matter of chance, dependent upon uncertainty of action at low temperatures.

An interesting feature of the series is the regular rise in the percentage of lice which die while hatching as the action of the solutions become less fatal up to a mid point, when it again declines to a minimum in proportion as the hatching ceases to be checked owing to the weakness of the solutions. It is a matter of question whether death during escape from the egg is due to the action of the fluids on the eggshell, or to a want of robustness in the larvae.

The evidence as a whole seems to establish the fact that steeping for twenty minutes in a 2 per cent. solution, either lysol or the cresol soap, is quite effective provided the temperature is not below 50° F.

A PLACARD is to be posted in the offices of town and country clerks in the State of New York stating that no marriage licence will be issued until both parties make the following statement on oath: "I have not to my knowledge been infected with any venereal disease, or if I have been so infected within five years, I have had a laboratory test made within that period which shows that I am now free from infection from any such disease."

SOME SIMPLE HOT AIR APPLIANCES FOR FIELD UNITS.

By LIEUT.-COLONEL C. MAX PAGE, R.A.M.C., S.R.

THE rapid transference of the wounded from such organizations as regimental aid posts and advanced dressing stations to the active surgical centre at the casualty clearing station is one of the principal problems which confronts medical units in the forward areas. During this period of transport the main difficulty with which the medical officer has to contend is shock; and once the wounds have been protected and any fractures satisfactorily splinted, his efforts are directed to keeping up or improving the general condition of the patients. The maintenance of body heat plays a most important part in this respect, and is a matter which, if neglected at this stage, may prejudice the whole later course of the case by leaving the man unfit for the essential operation at the casualty clearing station. Various schemes to this end are already in use, but I think it is worth while to describe some apparatus which has proved of practical value in dealing with the problem. More elaborate and capacious plant can be designed on the same principles under stationary conditions, when time and labour are of less account. That described below is essentially portable, and can be made up by any practical man from material ready to hand.

The first need is for some means by which a constant supply of dry and warm blankets and pyjamas can be maintained, and for this purpose a blanket drying chamber is used.

Blanket Drying Chamber.

The apparatus consists of a frame 32 in. square built of 3 in. by 2 in. wood. Two sides are tenoned and the others morticed (Fig. 1); on one face of the two morticed sections

FIG. 1.—Plan of blanket rack, 32 in. square. The main frame is made of 3 in. by 2 in. wood. The central rails form a continuous length of wire or cord wound through the slots in the side members. The outer rail is made of wire or standard splinting.

are cut, at regular intervals, twelve slots half through the wood and wide enough to take ordinary telephone wire. The frame is carried as a bundle of sticks. For use they are fitted together to form a square frame. This is kept firm by lacing with a continuous cord or wire around the slots to form the rack, as shown in the drawing. The wood should be so cut that there is a ½ in. projection of the tenoned and morticed ends; a groove is cut in the middle of these projections and another wire passed round the whole frame through them to form a surrounding rail; if preferred, this rail can be made of aluminium or steel splinting (from fracture box).

To form a drying chamber the frame is suspended by cords in a horizontal position (or supported on a light stand) about 5 ft. from the ground. Blankets folded to fit are hung over the surrounding rail and fixed with a few pins. These blankets are so arranged as to form four complete sides of a box (with blanket walls). The floor will form the bottom, and another folded blanket is placed over the rack, after objects to be dried have been hung on the wires, to form the top.

Two Beatrice stoves, with a simple tin baffle plate to protect the flame, are put on the floor inside the box as the

source of heat. In use no obvious air entry or exit should be left. Under average conditions (with two double-wick stoves) the temperature inside the blanket chamber is raised to about 120° F. in twenty minutes; at this stage, when the apparatus is in use for ordinary drying, the roofing blanket is pushed a little on one side to allow of ventilation, and the contents will then rapidly dry off. If the contents are very wet the temperature in the chamber should be raised higher before free ventilation is allowed.

Simple suspension of the frame, as described above, is the most practical arrangement for forward area work; under stationary conditions it may be supported on a skeleton stand.

Use as a Hot Air Sterilizer.

The same apparatus can be used for sterilizing small batches of clothing when a steam disinfectant is not available. The chamber should then be formed by at least a double fold of blanket all round, and three Beatrice stoves used. With this arrangement the temperature in the chamber can be raised to about 200° F. in twenty minutes. An exposure to this temperature for half an hour will kill lice eggs and the scabies parasite, as the garments, turned inside out, or blankets, being suspended, have their surfaces well exposed. This method does not damage the leather of breeches.

Hot Air Generator.

In an advanced dressing station the first method for raising the wounded man's body temperature is the hot air bath. For this purpose a hot air generator, as shown in the figure, is convenient (Fig. 2). It is made from a standard biscuit tin, or a four-gallon petrol tin; the latter is the stronger. The side of the tin is cut away except for a strip about an inch wide at the top, and a door may be put in the gap so made. The top is cut out and a domed cap fitted in its place. Into one side of the domed cap a short length of tin pipe is set, and a further series of lengths of tin pipe,

FIG. 2.—Heat generator made from biscuit box or four-gallon petrol tin.

prepared so as to telescope together, form the delivery pipe; the right angle bend is formed by an elbow of the ordinary pattern French stove piping. Solder, though useful for finishing off the dome, should not be relied on for the fixation of any part. If available, sheet asbestos may be applied to the dome and elbow. The addition of handles to the sides of the box is convenient. A double-wick Beatrice stove is placed in the box as a source of heat. The Primus pattern of stove is less reliable, and is not altogether safe for use in the closed apparatus described.

To give a bath, a properly blanketed stretcher on trestles is used, and an air chamber formed over the trunk of the patient by means of a simple cradle (Fig. 3), the whole

FIG. 3.—Skeleton cradle for hot-air baths. Made from aluminium splinting.

being covered by two or three blankets. The cradle frame may be made of skeleton splinting from the field fracture box or of stout wire. This arrangement allows dressings or operations to be carried out on the extremities while the heat is being applied. The box containing the stove is placed under the middle of the stretcher, and the piping arranged to deliver hot air just above the stretcher level. The box and contents are thus well out of the way, and only the pipe elbow projects from beyond the stretcher pole. With this apparatus a hot-air bath of from 130° F. to 300° F. can be given. For the wounded man a fifteen minutes' bath at about 130° F. is, I think, most satisfactory. The high temperature bath should be reserved for the treatment of urinary cases or myalgia under stationary conditions.

Memoranda:

MEDICAL, SURGICAL, OBSTETRICAL.

DISORDER OF THE RIGHT EXTERNAL SEMICIRCULAR CANAL.

THE two following cases, which present many points of similarity, are of interest:

1. A young woman who had previously suffered from three attacks of retrobulbar optic neuritis, after partaking at breakfast on January 15th, 1918, of rolled ham, which was noticed to be of unusual colour and from which fat was absent, found herself on the way to business falling towards a friend on her right. She struggled against this symptom and giddiness during three days. Then vomiting set in. When seen, there was constant lateral nystagmus. The patient could not stand unsupported, and there was vomiting of cerebral type. There was no deafness to speech test, and no tinnitus. The reflexes of the lower limbs were normal. Twelve days later the nystagmus could be barely elicited; the patient could walk and stand. The condition was that of one after nine days in bed.

2. A man aged 53, after partaking at breakfast on January 26th of ham which was noted to be watery, went to business. Some three hours later he suddenly collapsed on to the floor of his office. He was removed home. His gait as he was helped was to the uninitiated that of a drunken man, and many have been the jokes at his expense. Seen five hours later, he was retching and vomiting mucus like an alcoholic in the early morning; he is not alcoholic. Nystagmus, lateral, was most marked. He could not stand or walk unaided. There were no subjective aural sensations, nor was there any deafness to speech. Now, eighteen days after the onset, there remains only slight incapacity for mental activity. During a week there was slight hissing tinnitus, and for a few days there was a distinct tendency to fall to the right.

In neither case was there paralysis of cranial motor nerves, nor drowsiness, nor any definite ptosis.

H. D. O'SULLIVAN, B.A., M.B., B.C. Cantab.

Burton-on-Trent.

THROUGH-AND-THROUGH RAMROD WOUND OF HEAD.

ON November 9th a girl, aged 8, was brought to me with the ramrod of a gun fixed firmly in her head. It appears that a gun had been left on a shelf, loaded with the gunpowder charge only, and the ramrod left in it. The child in pulling something off the shelf had caused the gun to fall forward, when it went off. The ramrod had entered the cranium one inch above the top of the forehead, half an inch to the left of the middle line, and was protruding for some four or five inches from a point half an inch to the right of the same line and two and a half inches above the occiput. On admission the child was conscious, but the pulse was scarcely perceptible. There was no bruising of the skin, nor any haemorrhage, and the ramrod was so firmly fixed in the skull that the point had to be sharply tapped with a mallet in order to loosen it sufficiently to allow of its withdrawal. For