H. T. Butlin. I performed the operation on March 3rd, 1902. The patient made a good recovery, was afterwards married and has had four children. I saw her yesterday when she was in good health and had not suffered from any recurrence of haematemesis.

CASE VI.

A boy of 15 years with a cyst between the mucous and muscular layers of the pyloric end of the stomach. I performed pylorotomy on July 17th, 1906, and reported the case in the *Lancet*, August 18th, 1906.

Perhaps the most disappointing cases are the gastrostomies for the relief of stricture of the oesophagus, which is usually due to malignant disease. These cases are left until the last moment. The patients have experienced much suffering, and as a result of insufficient food have lost a large amount of flesh and vitality. When I have been able to operate upon them early I have succeeded in relieving their distress and prolonging their lives for as much as six months. I cannot claim to have obtained for them a longer respite. The modified Witzel's operation which I perform does not allow any regurgitation of food.

Contrast this experience with the medical treatment of these cases. Those who suffer from pyloric obstruction remain invalids. They continue to exist upon patent foods, they swallow large quantities of medicine, and they use up many hours of their doctor's time in a vain endeavour to obtain relief. Some of them indulge in stomach lavage. They lose weight, amounting in some cases I have seen to five or six stone, more than half their normal weight, and they ultimately succumb, after suffering untold misery. They are found in every consulting room, they go from doctor to doctor. Disappointment drives them to quack medicines, and ultimately, often when it is too late, they drift into the hands of the surgeon.

I do not suggest that all these cases should be operated upon without medical treatment, but I do claim that my results justify you in urging these people to consult a surgeon when you find that medical treatment fails, and I would emphasize the necessity of following this course before the condition is so far advanced as to render surgical intervention unduly dangerous, if not hopeless.

A CASE OF VALVULAR DISEASE OF THE HEART, SHOWING UNUSUAL POWERS OF PHYSICAL ENDURANCE.

BY

ROBERT D. RUDOLF, C.B.E., M.D. EDIN., F.R.C.P.,
CONSULTING PHYSICIAN TO THE CANADIAN FORCES
IN ENGLAND.

EVERY medical officer has come across soldiers who, in spite of evident organic disease of the valves of the heart, have been able to carry on at the front, and have only finally been invalidated for some other condition; but the following case is an example of such unusual physical power that it seems worthy of being put on record:

Lieut. A. B., aged 25, infantry officer, was invalidated to England on account of slight "gassing," and two months later was seen at a convalescent hospital, still complaining of depression, especially in the mornings, of weakness, and of shortness of breath on climbing hills.

History.

He stated that he had never been ill except when, at the age of 13, he suffered from an attack of rheumatic fever, and his mother had made him "go carefully for twelve months afterwards." He had been an athlete all his life, and won many prizes in this capacity at school. At his university he was captain of both cricket and football teams. During the last four years he had developed specially the 100 yards race. He had been in the army for about three years, and had never been sick or laid up, except as follows later.

Last October, while in France, he won the 100 yards championship of his division, and also the broad jump, doing the former in ten seconds, and attaining 22 ft. 6 in. in the latter. He was presented by the G.O.C. with a gold medal as the champion athlete of the whole corps. During last winter he was captain of his battalion football team, and played in every game during the season.

In November last he was blinded for two hours by the near explosion of a shell, and since then his right eye has not been as good as formerly, but he carried on all military duties and continued, as already stated, to lead his football team until the spring, when he was slightly gassed. He was then invalidated to England, though unwilling to leave France.

Condition on Examination.

The patient was fairly fit in appearance, but stated that he was often depressed in the morning. If he exerted himself much he suffered from palpitation and shortness of breath. All these symptoms only dated from the gassing. The pulse was moderate in rate and quite regular, but of water-hammer type; no distinct capillary pulse could be made out. The liver was not enlarged, and there was no oedema. On percussion of the heart the right border was found to be one inch beyond the edge of the sternum. The apex beat was felt heaving powerfully in the sixth intercostal space in the nipple line. On auscultation a loud double aortic murmur was heard and also a very evident double mitral one.

The condition was evidently one of aortic incompetence, and possibly stenosis (although the systolic murmur here might have been due to dilatation of the root of the aorta), and of mitral leakage and probable stenosis, although the presystolic murmur might have been of the Flint type. The heart was certainly considerably hypertrophied.

Dr. Lewis, in his paper in the *BRITISH MEDICAL JOURNAL* of December 14th, p. 647, on Medical Reports on Soldiers discharged from the Army for the conditions known as "D.A.H." and "V.D.H.," says that "In some patients with aortic disease exercise tolerance is perfect. Many soldiers have fought in the front line with this lesion and without mishap." He continues: "But these are cases in which the lesion, as judged by the state of the pulse, is slight and in which there is no material enlargement or other complicating factor."

Every one has seen cases of aortic regurgitation of this "slight type," who have stood military service for long, and I have notes of a number who have endured even unusual strain without a breakdown, but the above case is out of this class. The patient would undoubtedly come under Dr. Lewis's class of "developed aortic disease"; yet he not only stood the stress of active service in the front line for years and was at last only invalidated on account of gassing, but actually outrivalled all his fellows as an athlete. His reserve heart power must have been greater than that of most men having sound hearts.

The case emphasizes the correctness of the view that has so often been urged in recent years by Sir James Mackenzie and others, that everything depends upon the state of the heart muscle. This patient had evidently an exceptionally strong and sound myocardium.

The reason why most cases of chronic V.D.H. break down is not so much on account of the presence of the valvular lesion as because the infection which caused the damage of the valve at the same time invaded the heart muscle more or less. In the case here recorded the rheumatic invasion at the age of 13 must have been unusually limited to the endocardium.

Nevertheless, valvular lesions, whether causing stenosis or incompetence, are not only always *loci minoris resistentiae* against further infection, but must also, by their disturbing influence upon the smooth onflow of the blood, throw extra work on the heart muscle, and I for one do not go so far as some in the modern tendency to ignore such lesions. It is not my intention here, however, to go further into this debatable ground, but rather merely to put on record what appears to be a case of very exceptional endurance of a hampered heart.

NASAL DIPHTHERIA AND THE EPIDEMIC OF INFLUENZA.

BY

E. MARTEN PAYNE, M.B. ABERD.,
BLACKBURN.

It seems to have escaped the notice of those who have published notes on the prevailing epidemic of influenza that a large number of patients supposed to be suffering from that disease are in reality suffering from nasal diphtheria.

Feeling convinced that the heavy mortality during the epidemic could not be due to influenza pure and simple, I began to examine certain cases for nasal diphtheria, and found that many cases from which I took nasal swabs gave positive results. In many a provisional diagnosis of influenza had been made, and they were apparently convalescent when the swabs were taken. Others, at the onset, showed signs which indicated the trouble, and in others again the swabs were taken almost at random