layer of that membrane to deal with what infection remained after removal of the primary focus. I had always found it necessary, however, to drain the abdominal wall when there had been soiling of the wound edges. By filling the wound with the emulsion after closing the peritoneum, I have found it safe completely to close the wound, and thus avoid the potential incisional hernia which a drainage tube leaves. The following two cases furnish fairly severe tests of the efficacy of the emulsion in recent wounds.

CASE II.

T. L., a farmer, aged 38, with gunshot wound of elbow. The lower third of the humerus, the upper end of the radius, and the surrounding soft tissues were destroyed. The lower end of the remaining fragment of the humerus, the upper end of the lower fragment of the radius, and the upper extremity of the ulna were exposed in a wound extending from four and a half inches above the elbow to three inches below the joint, and the floor of the wound was formed by the brachial artery. There was no destruction of nerve trunks. The comminuted bone, and as much as possible of the damaged muscle, was removed. Owing to extensive destruction of skin it was impossible to close the wound, but the edges were drawn together at the upper end of the wound as far as possible without tension, and the cavity was lightly packed with gauze wrung out of flavine. The wound was dressed every three days. Owing to its size the wound has been somewhat slow in healing, and now at the end of nine weeks there remains a superficial ulcer about the size of a florin, and at no time during the period of healing has there been any evidence of infection—and this in spite of the fact that the wound was badly contaminated with manured soil, clothing, and the wad of the cartridge. The scar tissue is small in amount, and is soft and elastic. The limb is flail. After improving the nutrition of the limb by massage and electrical treatment, I intend to ankylose the elbow by means of a bone graft, and I have little doubt that a limb which at first looked fit only for amputation will be converted into a strong and very useful one.

CASE III.

O. J., a woman aged 22, had a compound comminuted fracture of the fibula due to direct violence. There was much bruising and laceration of surrounding tissues, and the anterior tibial nerve was divided just above the annular ligament. The patient was in a filthy condition, and dirty clothing was embedded in the wound. The patient's skin was cleaned with petrol, and sterilized with iodine. As much as possible of the soiled and damaged tissues was excised; the damaged nerve was sutured with fine catgut. After all bleeding had been controlled, the emulsion was applied to the surface, and the wound was closed with salmon gut. In seven days the six-inch wound was healed except for half an inch at its upper end. There was no evidence of infection in this half-inch, which had healed by the twelfth day.

The fracture of the fibula followed a normal course. In three months protopathic (but not epicritic) sensation has returned in the area of distribution of the anterior tibial nerve.

The only disadvantage that this emulsion has is its price, which works out at about five shillings for eighteen ounces. Having in mind the way in which it expedites the patient's recovery, and limits the disability resulting from his injury or disease, the saving of man power, particularly in industrial accidents, is such that the relative cost of the preparation is negligible.

TREATMENT OF ACUTE GONORRHOEA: MESSAGE-PACK METHOD.

By J. P. FOGARTY, M.C., MAJOR A.A.M.C.,
S.M.O. OVERSEAS TRAINING BRIGADE, A.I.F.

It is now an Australian A.M.C. order that the message-pack method of early treatment of gonorrhoea shall be used to the exclusion of all others.

As the originator of this method, and the first to point out to the A.D.M.S., A.I.F. dépôts in the United Kingdom its superiority over irrigation and sealing up processes, I wish to submit the following detailed account of my efforts and the results obtained.

My original object in devising the method was to put into practice a theory that the speediest and the best curative results would be obtained by dressing and draining the infected portion of the urethra, by the application of an antigonococcal chemical to the infected surface, and by preventing the spread of the mischief.

To attain these objects the basis of the treatment consists in packing the anterior portion of the urethra with a material soaked in silver solution. The pack not only absorbs the discharge and drains the infected part, but, by

mechanically dilating the urethra, it enhances the prospects of the chemical reaching the site of infection, and prevents apposition between infected and non-infected surfaces.

For packing I have found soft gauze far superior to cotton-wool. It is more absorbent, less irritating, and more easily passed into and withdrawn from the urethra; it has no tendency to break, and shreds are not detached from it.

In the large majority of my cases argyrol has been used for preference, but during periods of shortage silver vitelline has been tried with equally successful if not quite as speedy results.

From a curative point of view an argyrol solution stronger than 5 per cent. is not necessary, and 7 per cent. and 10 per cent. show a tendency to cause soreness, necessitating less frequent packing, thus depriving the case of the benefit of the mechanical advantages of the pack. I have found haemorrhage even with 10 per cent. a rare occurrence, and not in one single instance with 5 per cent.

Method.

The patient having passed water, and the glans penis and prepuce having been well washed with 1 in 8,000 solution of potassium permanganate, a probe is introduced into the urethra to determine its general direction and whether there is any narrowing or other obstruction.

A strip of soft open woven gauze is cut half to three-quarters of an inch in width and about seven inches in length. The width of the strip should vary with the size of the meatus, and with experience and practice can be estimated accurately. Having rolled the strip into a cylinder and well soaked it in 5 per cent. argyrol, one end is folded over the extremity of a probe and passed up the urethra until resisted at the bend. Thus the whole anterior urethra will be packed. The excess of the pack may be cut off at the meatus.

The procedure is facilitated by opening the meatus with the tips of the thumb and index finger of the hand holding the penis, and by balancing the free end of the strip on the little finger of the hand holding the probe. If the penis be slightly tilted upwards and the strip well balanced the loss of argyrol by dripping will be obviated. There is no danger of injury to the urethra if a moderately large probe with a good bulbous extremity be used.

Having packed the anterior portion of the urethra, the penis is then gently massaged for five minutes, all the movements being made towards the root of the penis.

The pack is then withdrawn, the meatus, in so doing, being gently compressed between the finger and thumb in order to leave behind in the urethra as much argyrol as possible.

A second pack is then introduced in the same manner as the first and retained for four hours, or, if the patient does not require to pass water, even longer. Whilst the pack is being retained for four or more hours there is no tendency for the discharge to trickle outwards along its side, and on withdrawal there is no evidence, such as a flow of pus, to suggest an accumulation of discharge behind the pack. On the other hand, on withdrawal, the pack will have lost most of its argyrol colour, and pus, especially in cases with copious discharge, may be easily squeezed from it.

The number of packs given in the day's treatment will be guided by the condition of the case, but as a general rule the pack is indicated with a purulent discharge or with acute subjective symptoms, such as scalding, even before the purulent stage is reached. The majority of my cases have responded to two packs a day (morning and evening), but when necessary, as in a number of cases, I have found three packs in one day easily tolerable.

The pack is rarely necessary after the fourth morning of treatment. As soon as the discharge, observed before passing water in the morning, ceases or becomes non-purulent (clear watery) the pack should be discontinued and treatment confined to gentle irrigation with 1 in 5,000 potassium permanganate three times a day.

Results.

Between the dates March 1st and December 31st, 1918, I have applied the treatment to 742 cases of acute gonorrhoea. Of these the number "not cured" and evacuated to a venereal hospital on or before the eighth day of

treatment according to orders was only 13, comprised as follows:

Uncomplicated failure	...	7
Complicated (bubo 1, epididymitis 5)	...	6

Up to early September, that is, with 498 cases, cure was determined by the absence of subjective symptoms combined with a dry urethra or merely a clear watery secretion observed on massage and expression in the early morning before the patient had passed water.

Since that date the decision on physical signs as to cure of the remaining 244 cases has been supported by the microscope. Unfortunately, owing to the difficulty in obtaining counterstain, only the smears relative to 184 of the cases were stained by Gram's method. Smears of all these 184 cases were taken before water was passed on the mornings after the first, the third, and the fifth day of treatment, with the following results:

Number in which Gonococci were Found.		
In smear taken on morning after the first day	...	148
In smear taken on morning after the third day	...	23
In smear taken on morning after the fifth day	...	1
(This case was considered a failure, and is included in the 7 uncomplicated failures mentioned above as having been evacuated to venereal hospital.)		

Further, with this series of cases confirmed by the microscope, advantage was taken of an order to the effect that men cured of gonorrhoea would be retained in the dépôt for an observation period of nine days after cure. All 183 cases were examined and smears taken on the third, the sixth, and the ninth days of this period, and in no single instance was a return of the physical signs observed or were gonococci found.

Relapse.

Unfortunately I have been unable to trace my cases beyond twenty-four hours after their arrival at the dépôts to which they have been marched out. Nevertheless, it is worthy of note that, whilst it is an A.A.M.C. order that all men will be medically examined within twenty-four hours of marching into a dépôt and disabilities found reported, I have received official intimation of only nine of my cured cases having relapsed. Most of my cases proceeded to France, and the question of a fresh infection would have to be considered.

Advantages of the Method.

Apart from the encouraging results, only seven uncomplicated failures in 742 cases (183 out of 184 proved by at least four negative smears in each case), the method has in addition the following advantages:

1. The ease and simplicity with which it can be carried out.
2. It is comfortable and soothing to the patient, alleviating the sensation of wanting to pass water, and there is much less tendency to painful erections which characterize the condition under irrigation methods.
3. It reduces irrigation and syringing during the acute purulent stage to a minimum, and thus lessens the risk of driving the infection into the posterior urethra or further.
4. It does away with the leakage of the discharge at the meatus and thus obviates the possibility of troublesome balanitis. This condition did not occur once in my series of cases.
5. On withdrawal a definite pattern on the pack or the adherence of mucus or shreds will often indicate local areas of disease.
6. It has an economic advantage over other methods, much less silver solution being required.

I am of opinion that by specially designed speculum or introducer, infection of the posterior urethra could be treated to advantage in similar manner.

ACCORDING to a bulletin published by the Health Department of Chicago, among the ten principal cities of the United States, Pittsburgh had the lowest death-rate from tuberculosis for the year 1917, the rate being 147.05 per 100,000 of population. Chicago stands second with a death-rate of 148.37 per 100,000. Next come Detroit with 160.66, Boston with 170.87, New York with 176.75, Cleveland with 174.7, Los Angeles with 199.42, St. Louis with 202.95, and Baltimore with 236.61.

Memoranda:

MEDICAL, SURGICAL, OBSTETRICAL.

TREATMENT OF INFLUENZA AND INFLUENZAL PNEUMONIA.

In view of the recrudescence of influenza, and especially of cases complicated with pneumonia and bronchopneumonia, I beg to suggest methods of treatment which I found satisfactory in a large majority of cases.

In cases of influenza, even if severe, I prescribe the following:

R. Sodii salicylas (B. W. and Co.)	...	gr. vij-x
Sp. ammon. aromat.	...	m x
Sp. chloroformi	...	m x
Tr. belladon.	...	m v-vij

Every four hours.

The belladonna has an excellent effect in removing the frontal headache. If there is an irritable cough, I add to the dose 5 minims of antimonial wine. Under this treatment cases usually pursue a normal course. The state of the bowels must of course be attended to, and the patient kept in bed for forty-eight hours after the temperature becomes normal.

If pneumonia or bronchopneumonia sets in, the latter so often double, my internal treatment is 5 to 7 grains of ammonium carbonate with 3 to 5 minims of tincture of strophanthus. If symptoms arise indicating fibrillary action of the right auricle, I add 5 minims of tincture of digitalis to each dose. I place most reliance, however, on the external treatment, which consists in the free application of turpentine. It is mixed with hot water, and flannels soaked in it are applied round the chest walls.

The solution must be strong enough to produce a glow and a tingling effect with some redness of the skin. The turpentine obtainable now is of poor quality; the quantity mixed with the hot water, therefore, must be increased.

Directly after applying the turpentine a well-fitting pneumonia jacket made of Gamgee tissue is put on. The turpentine fomentations must be repeated sufficiently often to keep up a glow on the skin.

If a good nurse is not available, the patient could be rubbed with warm turpentine, and the pneumonia jacket afterwards put on. Under this treatment the patients experience a great sense of relief; respiration becomes less laboured; the delirium so often present, especially in double pneumonia, subsides, and the patient sleeps better. Recovery often takes place in very critical cases. Turpentine capsules given internally do good, but patients often revolt against them after a few have been swallowed.

Bournemouth.

A. HUMPHREY DAVY, M.D., M.Ch.

Reports of Societies.

PERIODICITY OF MEASLES.

At a meeting of the Section of Epidemiology and State Medicine of the Royal Society of Medicine on February 14th, the President, Lieut.-Colonel E. W. GOODALL, R.A.M.C., in the chair, Dr. BROWNLEE gave an account of the results of an investigation into the periodicity of the epidemics of measles in the large towns.

The Method of the Periodogram.

He began with a brief description of the method of the periodogram analysis as applied to the discovery of periodicities in series of statistics. In general, if one periodicity only existed, ordinary inspection associated with a process of trial and error was quite sufficient to determine the periods accurately. As an example of this, sunspots were quoted. In certain towns, such as Aberdeen and Paisley, the only period for measles existing for very considerable stretches of years was quite obviously two-yearly, but when several periods existed, especially if one period were more important than a second, it was often impossible accurately to determine the second by mere inspection, while if three or more periods existed, inspection was useless. The method of the periodogram was essentially a method which examined all periods. The first person seriously to consider periodicity in disease was