each twenty-four hours. Delirium usually sets in much earlier than in lobar pneumonia, but it is very rarely violent or maniacal, even in alcoholics, but is of a muttering type, as in typhoid.

The pulse is by far the best guide as to progress in these cases. It falls in volume and increases in rate to a remarkable degree, almost as soon as the patient becomes ill. The rate rises to about 120, and in many cases that rate was maintained almost to the very end, with the pressure steadily falling. The respiratory rate is more variable, ranging from 40 to 60, or higher. The breathing differs from that in pneumonia. The breaths are deeper, and are not cut short in the way commonly seen in lobar pneumonia. The principal differences from ordinary pneumonia observed on auscultation are that practically no areas of complete absence of breath sounds are found, and the fact that all parts of both lungs are affected. Over the sternum and inner margins of the lungs loud bronchitic rales are heard, and further out, and towards both bases, coarse crepitations everywhere, with occasional patches of loud pleuritic rubbing. The sounds indicate a severe general bronchopneumonia with pleurisy. Percussion reveals areas of dullness in several parts of both lungs, or there may be a board-like dullness over a large area on both sides. The intercostal spaces are drawn in with inspiration. Deep and rapid breathing of the air-hunger type is frequently seen, and is of the gravest significance. The great quantity of sputum gives trouble very early in delirium, owing to the absence of efforts to clear the throat while coughing goes on. This leads to an accumulation of sputum in the throat which is very troublesome. The blood in these cases showed marked leucocytosis, especially of polymorphonuclear leucocytes.

Most of my patients were malarial subjects, and about 50 per cent. showed parasites in the blood, either active reproductive forms or crescents, but the course of the disease did not seem to be affected, the mortality being equal in those with parasites and those without.

Post mortem the lungs were either entirely or in great part hepatized. The pleura was covered with pus exudate, and there were many soft adhesions and free pus in the pleural cavity; the liver showed signs of early fatty degeneration, but the other viscera were not markedly affected. There did not appear to be any characteristic change in the spleen, though enlargement due to this specific disease might have been present and not distinguishable, owing to the malarial enlargement present.

The course of the disease was generally short. The mortality was very high, and no treatment appeared to be of the slightest avail.

I tried the injection of saline intravenously and intracellularly, and by the rectum, and quinine by various routes. I also tried novarsenobenzol intravenously, and exhibited salicylates and expectorants, besides giving the usual stimulant dietary, but the course was unaffected. The condition seemed to be one of severe toxæmia from the very commencement.

As with plague, typhus, and many other of the gravest epidemic ailments, prophylaxis should be given the first and greatest attention.

THE OCCURRENCE OF MORGAN'S BACILLUS IN CHRONIC DISCHARGING WOUNDS.

BY

HAROLD E. WHITTINGHAM, M.B., CH.B. GLASG.,

MAJOR R.A.F., M.S.,

BACTERIOLOGIST, R.A.F., CENTRAL HOSPITAL, HAMPSTEAD.

THE following observations were made on wound infections occurring in a Poor Law infirmary. Here it was noticed that certain deep wounds, which were moderately clean on admission, gradually went from bad to worse in spite of careful treatment. These cases ultimately developed chronic discharging sinuses, the discharge having a peculiar odour very like that of a bacillary dysentery stool. Several of these patients were obviously suffering from a toxæmia; blood cultures performed were negative.

Cultures made from wound discharges of six of these cases revealed the presence in all of an organism identical with Morgan's No. 1 bacillus.¹ This organism was the predominant one in most instances, but other organisms, such as streptococci and staphylococcus, were numerous.

Autogenous vaccines of Morgan's bacillus, given in two of these cases, greatly improved the wound and general toxic condition of the patient.

Where did this infection with Morgan's bacillus occur? Was it the cause of this chronic wound infection?

The points to decide were: (1) What organisms were present in the wounds on admission to the hospital? (2) What organisms gained entrance to the wounds during residence in the hospital? (3) The source of any infection developing whilst in hospital.

Technique.

Immediately on admission to hospital swabbings were taken from wounds of twenty cases. Cultures were made aerobically in broth, on agar, blood agar and MacConkey's bile-salt-lactose-neutral-red-peptone-agar, and incubated at 37° C., and on gelatine plates at room temperature for seventy-two hours. Anaerobic cultures were made in glucose broth incubated at 37° C. for four days. At weekly intervals, for as many weeks as the patient was detained in hospital, the above procedure was repeated. The stools of these cases were examined on two consecutive days for organisms of the dysentery group.

Results Obtained.

From the swabs taken on admission various micro-organisms were found, often a mixed flora, chiefly *Streptococcus viridans* and *Staphylococcus aureus* and *albus*, in one case *B. pyocyaneus*, and in another *B. perfringens*. In none of the cases could Morgan's bacillus be isolated. Examination of pus from these wounds whilst in residence in hospital revealed the presence of Morgan's No. 1 bacillus in five out of the twenty cases—that is, 25 per cent. These organisms made their appearance in two weeks after admission in four of the cases, and in the third week in the remaining one. This was the only modification found of the admission flora except in one instance, where *Staphylococcus pyogenes aureus* was superimposed.

Morgan's bacillus soon became the predominant organism in the five infected cases. In this connexion two agar platings might be mentioned. One plate developed 205 Morgan's bacillus and 26 streptococcus colonies, the other plate 226 colonies of Morgan's bacillus and 51 of *Staphylococcus pyogenes albus*, after incubation for forty-eight hours at 37° C. The characters of this Morgan-like bacillus¹ were fairly constant in all the cases observed—namely, a short Gram-negative bacillus, non-motile or feebly motile, fermenting glucose with the production of acid and a little gas, although in some there was no gas formed; slight acid production in maltose. Lactose, saccharose, mannite, and dulcitol were unchanged. Litmus milk was rendered slowly alkaline from the second day without previous acidity. Indol production was distinct in all except one case. This latter organism corresponded closely with Eyre's No. 9.²

As regards the source of infection with this bacillus, all the cases which developed this condition occurred in two wards widely separated and under different medical officers. The sterilized dressings used in the wards were examined on two occasions, with negative results. Stools of all cases proved to be negative to organisms of the dysentery group.

Agglutination reaction of the bacilli isolated with the patient's own serum yielded negative results, except in one case, where there was a positive reaction to a dilution of 1 in 40 in half an hour at 37° C.

Treatment.

Wounds infected with Morgan's No. 1 bacillus responded slowly, if at all, to ordinary surgical drainage and antiseptic dressing. In four cases treated by autogenous vaccine there was marked improvement of the local and general condition. The septic type of fever and odour began to disappear usually about three to four days after the first inoculation. Four to five inoculations were found to be sufficient. The dosage of vaccines employed was:

First dose	...	100 million	Morgan's No. 1 bacillus.
Second dose	...	200	" "
Third dose	...	300	" "
Fourth dose	...	400	" "
Fifth dose	...	500	" "
Sixth dose	...	600	" "

In two cases there was a marked local and general reaction after the first inoculation, but when this passed

off there was a distinct improvement in the condition of the patient.

Conclusions.

That an infection by Morgan's No. 1 bacillus was superimposed on the wounds in this Poor Law hospital is evident from the absence of that organism in the twenty cases dealt with on admission, and its presence in five of them at varying periods after admission. The pathogenic action of this organism in wounds is evidenced by the toxic condition of the patient and the chronicity of the suppuration as compared with similar wounds from which this bacillus was not recovered; also from the fact that under treatment with an autogenous vaccine prepared from this virus the general and local conditions markedly improved. The odour perceptibly diminished from the wound in three days, and within ten days a dull and apathetic bed patient would be up and cheerily smoking, with his wound comparatively "clean."

The source of infection in the hospital is uncertain. The question of non-sterile dressings was considered but threw no light on the subject; bacteriological tests all proved negative. That the source was general is apparent from the fact that cases occurred in surgical wards widely separated, and attended by different medical officers and members of the staff. This point eliminates "carrier" infection. Owing to my leaving the institution I was prevented from concluding these investigations. But as the dysentery group of the bacilli, including Morgan's No. 1 bacillus, has been repeatedly found associated with cases of asylum or institutional dysentery in this country, the suggestion arises that a Poor Law infirmary is a likely habitat of these organisms, and the infection may have occurred from the dust of the place.

Although these notes are incomplete their publication may call forth similar observations from others. Should this be the case institutions of the kind should not be used as general hospitals for troops, at least until they have been thoroughly cleansed and disinfected throughout.

REFERENCES.

¹ H. D. Morgan, BRITISH MEDICAL JOURNAL, 1906, vol. i, p. 908; *ibid.*, 1907, vol. ii, p. 16. ² J. W. H. Eyre and E. P. Minett, BRITISH MEDICAL JOURNAL, 1909, vol. ii, p. 1227.

Memoranda:

MEDICAL, SURGICAL, OBSTETRICAL.

THE NASAL PLUG AS A GUARD AGAINST INFLUENZA.

THE wearing of masks in railway carriages and crowded public rooms has been advocated as a means of protection against influenza. There is, however, a natural disinclination among most people to make themselves conspicuous by wearing masks, and rather than do this they prefer to take their risk of infection.

This objection can be met by the use of the nasal plug, which, when properly employed, is just as efficacious as the mask. All that is necessary is to carry about a piece of cotton-wool and, as occasion requires, pull from this two small plugs and insert one into each nostril.

It is important that these plugs should be made properly. Each should be big enough to fill the nostril completely, and the texture loose enough to allow of easy respiration through it. This is ensured by taking quite a small piece of cotton-wool and pulling it out so that it becomes really *fluffy* and about the size of the end of the thumb. One of these plugs is then lightly packed into each nostril so that it is just out of sight. While in place air should be inspired through the nostrils only, never through the mouth. Expiration may take place either through the nostrils or mouth. This procedure requires the minimum of attention and speech is not interfered with as this normally takes place only during expiration. If the plugs are made too thick, then, besides interfering with respiration, they render the speech stuffy as though one has a cold in the head.

They are best removed, on coming out into the fresh air, by blowing them into a handkerchief. They can then be taken home and burnt.

The use of an antiseptic wool such as cyanide is an extra precaution, but ordinary cotton-wool is more easily

procured and answers the main purpose which is mechanical.

It should be impressed upon the public that—

1. Infection takes place by the droplet method—small particles of mucus laden with microbes which may be coughed by an infected person to a distance of three or four feet.

2. It is when within this range that the danger of infection occurs.

3. This danger is increased when the atmosphere is confined, as the air becomes more heavily laden.

4. To prevent any of these droplets from being breathed in some form of screen or filter is necessary.

H. C. LUCKY, Captain R.A.M.C.,
Bacteriologist, No. 6 Stationary Hospital, B.E.F.

MUSK IN INFLUENZA.

THE present serious epidemic of influenza, with its heavy mortality both from the disease itself and from subsequent pneumonia, reminds one forcibly of the epidemic of 1890. It also recalls to my mind a method of treatment I adopted. This was the employment of musk as a powerful heart stimulant. I used it in the form of a pill, dose 1 grain, in combination with an equal quantity of quinine, given every four hours. I cannot tell whether it was *post hoc* or *propter hoc*, but shortly after the commencement of the treatment I noticed a distinct improvement in the course of the disease, especially in the mortality from pneumonia. The drug certainly did no harm, and it would probably be worth trying in the hope that some good may come from it. The drug is certainly costly; I am told it costs 6d. a grain at the present time.

Bournemouth.

ARTHUR RANSOME, M.D., F.R.C.P.

LOSS OF HAIR DUE TO LIGHTNING.

DURING a thunderstorm in October, 1918, a house was struck by lightning and much damage done. Three children, a girl aged 9 years and two boys aged 7 and 5 years, were in the same bed on the first floor. According to the mother's statement, the children were all lying with their heads up at the same end of the bed, and all were asleep on their right side, as is their usual custom. The children were naturally frightened by the thunder and lightning, but did not suffer any harm. A fortnight afterwards the hair on the left side of their heads from the middle line downwards began to come out, and in a few days the scalp on the left side was quite denuded of hair. The right side of their heads was protected from the lightning by the pillow. At the present time (three months later) the hair is slowly growing again, and all the children are in good health. The lightning seems to have had a direct effect on the hair follicles like the x rays.

HUGH T. ASHBY, B.A., M.D., M.R.C.P. Lond.,
Visiting Physician to the Manchester Children's Hospital
and to the Salford Royal Hospital.

THE ASSOCIATION OF HERPES ZOSTER AND VARICELLA.

THE interesting article by Dr. R. Cranston Low (January 25th, p. 91) recalls a case that occurred in my practice some time ago.

Mrs. P., aged 40, was delivered of her fourth child in April, 1917, the delivery being normal. Three weeks afterwards severe pains set in in the region of the right hip, followed after a week's interval by a typical eruption of herpes zoster affecting the distribution of the twelfth dorsal and first lumbar nerves. A week afterwards several vesicles appeared on the left side of the neck. Phenacetin was administered, and afterwards morphine for the relief of the pain, which was excessive, but no other drugs were used. My friend Dr. W. H. Maidlow saw the case in