Diphtheria.

Of the 307 cases of aortic regurgitation 22, that is, 7.2 per cent., gave a history of diphtheria.

TABLE X.—Showing the Distribution of Cases with a History of Diphtheria and a Comparison between them and the Total Aortic Regurgitants with regard to their Rheumatic History.

	No. of Cases.	With Rh. Fever.	With Rh., G.R., or Ch.	Without Rh. History
Total Ao.R. ...	307	162=52.8%	73=23.8%	72=23.4%
Ao.R. and diphtheria ...	22	15=68.2%	5=22.7%	2= 9.1%

A comparatively large proportion of these cases were pure aortic lesions, namely, 13 as against 9 with mixed lesions, the ratio here being 1.44 instead of the ratio 0.76 shown by the total number of aortic cases with a history of rheumatic fever. Of the 7 cases without a history of rheumatic fever, 4 were pure aortic lesions and 3 mixed—a ratio of 1.3.

These figures cannot be said to lend any support to the idea that diphtheria is a cause of aortic regurgitation, but the large proportion of pure aortic cases in this group is interesting, though it may well be accidental.

Pneumonia.

The proportion giving a history of pneumonia is not large, 20 cases in all, that is, about 5.9 per cent. of the total 307 cases. Twelve of these gave histories of acute rheumatism, 2 of some rheumatic affection, and 6 no rheumatic history. Here again, as the table shows, there is no evidence from the figures that pneumonia is in itself a cause of aortic regurgitation.

TABLE XI.—Pneumonia.

	No. of Cases.	With Rh. Fever.	With Rh., G.P., or Ch.	Without Rh. History
Total Ao.R. ...	307	162=52.8%	73=23.8%	72=23.4%
Ao.R. and pneumonia ...	20	12=60.0%	8=40.0%	6=30.0%

Since an attack of pneumonia is apt to affect the muscle of the ventricles, it is not surprising that these cases of aortic regurgitation with a history of pneumonia show a larger proportion of mixed lesions than is shown by the 162 cases with a history of acute rheumatism; the ratio of pure aortic to mixed lesions being 0.43 where pneumonia had occurred, as against 0.76. The cases without a history of rheumatic fever comprised 2 cases of pure aortic lesions and 4 of mixed, a ratio of 0.5.

The analysis of these cases is against pneumonia being a cause of aortic regurgitation.

Influenza.

As judged by these cases, influenza is but of little importance in the production of aortic regurgitation. Of the 307 cases, 142 (46.3 per cent.) gave a history of one or more attacks of influenza, but in 82 of these there was also a history of rheumatic fever, and in 31 a history of chorea, "rheumatism," or growing pains—that is, in 79.6 per cent. there was a possible rheumatic history.

TABLE XII.—Influenza.

	No. of Cases.	With Rh. Fever.	With Rh., G.P., or Ch.	Without Rh. History
Total Ao.R. ...	307	162=52.8%	73=23.8%	72=23.4%
Ao.R. and influenza ...	142	82=57.7%	31=21.8%	29=20.5%

The ratio of pure aortic to mixed lesions was 0.73, which is very similar to that of the total number of aortic regurgitants with a history of rheumatic fever. Of the 60 cases without a history of rheumatic fever, 21 were pure aortic cases and 39 were instances of mixed lesions, the ratio being 0.5, against 0.76 for the whole of the cases of aortic regurgitation with a history of rheumatic fever.

(To be continued.)

A PLEA FOR THE HOMOGENOUS NERVE GRAFT.

By E. WELBY FISHER, F.R.C.S.,

TEMPORARY ASSISTANT SURGEON, ST. GEORGE'S HOSPITAL.

In the majority of cases of complete nerve division that have resulted from gunshot injury the nerve ends are capable of approximation for suture when operation is performed for this purpose. But there are quite large numbers in which approximation is not possible owing to the destruction of a considerable segment of nerve trunk, together with extensive tissue loss. Many such a case has been operated upon, and the surgeon who has been confronted with the problem of how to make good the loss has failed to proceed further or has resorted to one of various illogical expedients, without, in my experience, any good result.

As surgical specialist to the Pensioners' Resurvey Boards at the Duke of York's Head Quarters I have had ample opportunity of observing these results, and I have seen no instance of recovered nerve function where bridging with foreign material has been done, or where the nerve ends have been connected by a wrapping of fascia, Cargile membrane, or vein. If Van Lair's experiments are to be relied upon, it is inconceivable how anatomical union can by such means occur.

He found, as a result of nerve trunk excisions in animals, that where a space of 4 cm. was left between the nerve ends no recovery occurred; with an excision of 2 cm., recovery did not occur for thirty months; with a 1 cm. excision, recovery of function was delayed to fourteen months; whereas in the case of approximation and suture, recovery occurred in eight months.

These experimental operations were aseptically performed so that there would be absence of scar tissue and the circumneural areolar and fatty tissue left would offer but little obstruction to growth, if any, furnishing rather an admirable conducting sheath to support and direct the growing nerve fibres in a correct course were this all that was required.

But it is clear that something more than a conduit is necessary. It is recognized that the neurilemma elements are essential to sustain and nourish nerve fibres in their growth. Where interrupting scar is present, as it invariably is in gunshot injuries, it is doubtful if the smallest breach is passed. Wherever a nerve has been wounded and divided, completely or in part, palpation will reveal the presence of a neuroma, which means a mechanical and functional block, and my experience is that until this block is removed and direct approximation of healthy elements obtains, function is not restored, in spite of massage, electricity, and other futile means wastefully expended over a period of years. It is not reasonable to expect growing fibres, after being contorted in scar, to traverse such a bewildering obstacle and reach their destination. Thus excision of a lateral scar the result of a lateral notch is to perform a recognized operation.

In view of the large number of pensioners still languishing with disabilities that are the direct result of sectional nerve lesions it is fortunately proving by accumulating evidence that lapse of time should be no deterrent against suture. The longest period that has elapsed between section and suture of the cases I have observed that have been followed by recovery is two years and seven months. This was a case of complete ulnar nerve division.

The results in general of nerve suture as they appear at the Resurvey Boards are remarkably good, and the degree of restoration of function is found on palpation to be in inverse ratio to the development of a neuromatous swelling where junction has been effected. Conformity at the site of union approximates the normal when coaptation is exact and restoration of function is complete; whereas if operation results in failure, marked neuromatous swelling occurs, which is a help in deciding for reoperation without undue delay. "What is anatomically correct will not be physiologically wrong" is a phrase that in nerve surgery may be taken as a guide if operative results are to give satisfaction. It is not possible to make certain of establishing perfect histological union, but the oval contour and striation of a nerve does permit of coaptation that approximates morphological perfection, so that torsion can be avoided and nerve fibres in their growth are likely on the whole to

find their paths correctly. The sectional area of a nerve has the appearance and manner of a cribriform connective tissue plate, the apertures of which are occupied at the proximal end by severed normal nerve fibres and at the distal end by protoplasmic neurilemma cords of equal number. If nerve fibres in their growth pass into other than their rightful channels, so-called paradoxical regeneration will occur that in mixed nerves in particular is followed by disconcerted and incomplete recovery; but when coaptation of nerve ends is made morphologically exact, such paradoxical regeneration is unlikely, for sensory and motor nerve fibres are not intimately admixed but can be traced throughout their course in separable strands.

A nerve graft fulfils all histological and anatomical requirements. Successfully introduced, it may be regarded as deriving vascularity and nutrition from capillary connexions that must at once form, and as undergoing the same Wallerian degenerative change of its nerve fibres, in common with all medullated nerve fibres severed from their cells. The loose connective tissue which surrounds and is introduced with a graft affords protection and mobility, and is a natural safeguard against adhesion that is so commonly responsible for local tenderness, contracture, irritative phenomena, and pain.

If a nerve graft is to afford a normal channel for every separate axis cylinder growing from above it must have a sectional area at least as great as or greater than that of the nerve trunk itself. Any provision short of this will be inadequate, and will result either in a syndrome of dissociation, a recovery at most of partial degree, or in failure of any recovery owing to the block of a faulty and defective union. Such unsatisfactory results are to be seen by the utilization of autogenous grafts for the median, ulnar, and musculo-spiral nerves when their continuity is re-established by the interposition of segments of cutaneous trunks of small and disproportionate calibre, like the radial and internal cutaneous nerves taken from the same subject. Union by suture of the epineural sheath, so insisted upon by many operators, is in this connexion an essential desideratum in the prevention of block, by eliminating the ingress of scar tissue and the emergence of nerve fibres, which become governed and directed on a normal course.

Homogenous nerve grafts are obtainable from amputation stumps which are being operated upon for painful neuromata, and it is quite feasible that certain men would be donors to their comrades who remain disabled from extensive nerve loss. In such case the time of operation is a matter of arrangement and choice. Opportunity is also afforded when amputation is performed if disease does not contraindicate, but such is only of chance occurrence, and rarely available opportunely. It is possible that nerve segments preserved in a cooling chamber might be employed with success as grafts, but, though I have had them ready in case of need, no occasion has arisen to prove their use, nor have I seen their use reported.

There are considerations other than anatomical which have been dealt with above that contribute to success in nerve reconstruction, but taking for granted a recognized and approved surgical technique, it is the anatomical factor which, in my opinion, counts above all others for or against a good result, and the importance of which as fundamental requires emphasizing. And the anatomical factor as a matter for attentive consideration is not confined to the nerve. The other tissues in the wound area that have suffered injury and loss are replaced by scar tissue, which is the product not of differential regeneration but of repair, in which the cellular elements of the universal connective tissues alone take a proliferating part. The breach of the surface and underlying tissues becomes indiscriminately occupied and mended by an inelastic material, binding parts together which should possess free mobility, dragging and distorting them from their normal positions and course, and fixing them to the overlying surface or subjacent bone.

I made it a practice in my military hospital work to excise all adherent scars with the object of reconstituting disturbed anatomy and restoring the mobility and normal function of the various structures. The aching of scars in cold damp weather is real, from loss of the insulating subcutaneous fatty covering; the pain of scars is true, and can be localized by careful palpation to fine neuromatous

terminations of divided subcutaneous nerves. Contraction and extension of muscles, free action of tendons, and consequently joint movement, are compromised by adhesion to skin and underlying bone. Nerves by implication evince not only disordered conduction, but distressing tenderness at the affected region both on manipulation and pull, so that attitudinization and protective contracture maintain deformity that resists correction from fear of pain. Many so-called functional cases have in these conditions an underlying cause, a sufficient aggravant in susceptible subjects being an adherent scar by the mere annoyance of its continued drag and pull. All these evils can be eliminated by excision of the scar, which, when an implicated nerve is approached, should be the route chosen rather than another elsewhere.

The foregoing remarks will, I hope, suitably preface the report of a case which I operated upon at St. George's Hospital, a case of extensive loss of the median nerve in which the intercalation of an homogenous graft fifteen months after injury was followed by a result so successful and rapid that I enter a plea for its general adoption and its trial at all events before resorting to tendon transplantation.

J. H. P., aged 32, was wounded on June 7th, 1917, the disability recorded being "gunshot wound of the right forearm with fracture of the radius." When seen by me in August, 1918, there were anterior and posterior long scars in the middle line above the wrist, adherent to and binding the underlying tendons and preventing their free action, especially as affecting extension of the fingers. Beneath the upper end of the anterior scar, which was 4 inches long by $\frac{1}{2}$ inch broad, a large fixed and tender neuromatous swelling was palpable in the course of the median nerve, giving rise on manipulation to electrical painful sensations referred to the area of cutaneous distribution of the median nerve in the hand, in which there was complete median paralysis and anaesthesia, with sores and other trophic changes. There was shortening of the radius by fracture and sequestration, but union was firm and its alignment good. An undue prominence of the head of the ulna was a consequence, with a disabling outward deviation of the hand.

He was admitted to St. George's Hospital, where I operated upon him on August 30th, 1918. In the first place the posterior scar was excised and the extensor tendons involved were freed. The anterior scar was then excised and the damaged implicated tendons were rectified and freed. All scar tissue as it involved the median nerve was removed, leaving an interval of $\frac{1}{4}$ inches between the cut ends, the lower being flush with the upper margin of the anterior annular ligament.

Whilst this operation was in progress Mr. Fedden, in an adjacent theatre, was amputating a lower limb through the thigh. The case (female) was one of chronic ulceration, but he was able to reassure me that the Wassermann test was negative. The skin surface was smeared with iodine, and I hastily cut down upon and removed a long portion of posterior tibial nerve, which, being of approximate calibre to the median nerve and abundantly surrounded with a loose fatty covering, appeared ideal for the purpose of a graft. This I dropped into a dish of methylated spirit and inserted in the wound, which was found bloodless on removal of its packing. The introduced nerve was cut to suitable length and sutured with fine catgut in position in accord with the anatomical principles described, a stay suture passed through the body of the nerve and graft first securing the ends, and small interrupted sutures circumferentially placed giving continuity of the sheaths. The wound surface was drenched with methylated spirit, dried, and given the slightest smear of bipp as an inhibitory agent against the development of sepsis. Fascia, skin, and subcutaneous tissue were united by catgut suture, no drainage being employed. Skin approximation was found impossible until linear incisions parallel to the wound on the radial and ulnar borders had first been made, the incisions passing only through the skin, leaving undivided the subcutaneous layer to preserve mobility when epithelialization should subsequently occur. The wounds healed by first intention, and the breaches made in the skin closed in a few weeks' time.

On October 14th, 1918, I performed a second operation, excising three-quarters of an inch of the ulna just above the attachment of the pronator quadratus. Coaptation was found difficult, and from the bone removed a disc the thickness of a shilling was therefore sawn and inserted to make the fit exact. Apposition was effected by a metal plate, the union being made secure and firm by screwing away from the line of junction. The deformity of the hand was thereby corrected. Healing was primary.

The subsequent progress of the case was most satisfactory. The man soon expressed a feeling that the hand was changed, that it felt alive. Tinel's sign was obtainable in the palm within three months' time. At the beginning of February, 1919, five months after the introduction of the nerve graft, an examination was made and the following notes recorded:

Union of the ulna firm; hand in straight alignment with the wrist. Extension of little and ring fingers beyond the straight line prevented by flexor tendon contracture. Hand grip good.

The hand was in a state of good nutrition, all signs of trophic

change having disappeared. There was absence of discomfort, tenderness, and pain.

Sensation to pin-prick wellnigh complete, being present over the whole median area with the exception of the tip of the index finger and the two terminal phalanges of the middle finger. Epicritic sensation present over the whole median area with the exception of the terminal phalanges of the thumb, the whole of the index finger, and the middle and distal phalanges of the middle finger. Thenar muscular power largely recovered, opposition of the thumb being present to considerable degree. Slight faradic response of abductor pollicis observable. Tip of thumb appposable to that of index and middle, not quite to that of ring and little fingers.

The hand at this time was restored to considerable practical use, and the man had resumed his occupation, which was that of a skilled electrician.

• HYPERPYREXIAL HEATSTROKE:

A BRIEF NOTE ON ITS ETIOLOGY AND PREVENTION.

BY

K. G. HEARNE, M.B., CH.B. EDIN.

In a recently written thesis on heatstroke—a disease which I studied in Mesopotamia during the exceptionally hot summer of 1917—the etiology, prevention, and treatment of the condition has been discussed more or less fully. It is impossible here to do more than touch lightly upon a few of the chief points which have a practical bearing on the causation and prevention of the disease. The object of this note, therefore, is to suggest the adoption, in heatstroke countries generally, of the following principles of prevention, which have proved completely successful in Mesopotamia after a prolonged and very thorough test under varying conditions. The remarks on etiology are based on a close study of some 240 cases, many of which occurred in the huts of a large base hospital among patients who were already in hospital with some other complaint, and in consequence an excellent opportunity of observing the disease in all stages of its development was afforded.

Etiology.

As regards etiology, and notwithstanding the diverse theories that are put forth in present-day textbooks on the subject, my observations in Mesopotamia point most conclusively to the belief that heatstroke is entirely due to suppression of sweating, which may be present from one to forty-eight hours before the attack.

With sweating suppressed, the bodily temperature tends to adjust itself, in accordance with physical laws, to the temperature of the atmosphere, which may be 115° to 120° at the time, and even to rise higher, for, as Sutton¹ and Graham and Poulton² have pointed out, a rise in bodily temperature produces increased respiratory and nitrogenous changes, resulting in a further increase of internal bodily heat. The processes continue until, when a temperature of about 108° or more is reached, sudden unconsciousness, delirium, and convulsions appear—the result of the physical action of heated blood on the specialized brain cells. The clinical picture of the fully developed attack of heatstroke is thus formed.

I have myself repeatedly observed the pre-existence of suppression—in one case for as long as eighteen hours before the attack—while in numerous other instances I have obtained absolutely reliable histories of it from patients on their recovery.

The cessation of perspiration is probably due to exhaustion of the sweating mechanism during several previous days of intensely severe weather, and there is strong reason to believe that the defect is localized in the sweat glands themselves. Furthermore, large doses of diaphoretics are powerless to produce sweating when once suppression has set in.

Prevention.

By the early recognition of suppressed sweating the prevention of heatstroke becomes an extremely simple matter. The practice which I adopted among hospital patients under my charge in Mesopotamia was to inspect them all periodically—every hour on extremely hot days—and to detect in this way those who required preventive treatment; any who were discovered to have a temperature

of 103°, with complete suppression, were stripped and covered with a wet sheet. This measure simply amounts to an artificial means of producing the same physical effects as sweating. In cases in which the temperature had already risen dangerously high before discovery a portable electric fan directing a current of air over the surface of the wet sheet was placed beside the bed; the evaporative process was greatly assisted by it.

In mild cases, or where the treatment requires to be maintained for several days until the natural sweating power returns, a convenient modification of the method, and one more comfortable for the patient, is to raise the wet sheet on a couple of bed cradles, thus forming a kind of cool chamber in which the naked patient lies protected from the surrounding hot atmosphere of the hut. With this device the additional aid of a fan playing a current of air over the outside surface is usually necessary in order to produce an efficiently cool atmosphere below the cradles on very hot days.

Needless to say, the sheet requires to be kept wet, and much time and labour is saved by using for this purpose some form of spray, such as the Holder-Harrinan pneumatic disinfecting spray, filled with cold water.

If it were possible to remove the patient to a ward which was artificially cooled to a suitable temperature the individual cooling devices would, of course, be unnecessary. The success of these preventive measures can be judged from the fact that in my hut (sixty beds constantly occupied) at the hospital, nine cases of heatstroke with four deaths occurred within the first few days of the 1917 heat wave, but, directly the preventive system was commenced, the outbreak stopped; only two more cases occurred during the remainder of the heatstroke weather, which lasted continuously for two months. The average number of cases occurring in this district (Basra) during the whole period was nine a day. Neither of the two patients who developed the condition in my hut after prevention had been commenced had been placed under preventive treatment.

It will thus be seen that hyperpyrexial heatstroke is an absolutely preventable disease, the leading signs of onset being a hot dry skin, which is typically harsh to touch, and a raised bodily temperature. There is usually also frequency of micturition. These signs are present from one to forty-eight hours before the attack, consequently ample warning is given. The position is simply this: that the patient, having lost his natural means of heat dissipation, requires his bodily temperature to be artificially regulated for him during extremely hot weather until he again begins to sweat, or until the heat wave abates.

If Europeans in heatstroke countries were made acquainted with the premonitory signs and principles of prevention, I see no reason why heatstroke, in a dry climate at least, should occur. I have had no experience of prevention in extremely humid climates such as Calcutta, where the atmospheric moisture may, possibly, hamper the evaporative process, but have no doubt that such a difficulty, if present, could be easily met by employing some other method of cooling the patient.

Even among troops on the march the principle of prevention is applicable. Water, and something that will act as a fan, are practically always obtainable, thus permitting an efficient form of first aid being applied to any man who exhibits the warning signs.

Among sleeping communities—for example, patients in hospital, or soldiers sleeping in barracks on very hot afternoons during heatstroke weather—periodic inspections are necessary in order to detect developing cases. These inspections can easily be carried out by an intelligent orderly, or staff of orderlies. The patient's general condition can be estimated roughly by touching the skin, and only occasionally is it necessary to take a thermometer reading, while those who are freely sweating can, practically speaking, be entirely disregarded.

In the after-treatment following upon an actual attack of heatstroke the tendency to relapses of hyperpyrexia are treated on exactly the same lines as for prevention. The patient frequently does not regain his powers of sweating for days after the attack.

REFERENCES.

- ¹ Sutton: *Journ. of Path. and Bact.*, 1909, vol. xiii, p. 62. ² Graham and Poulton. *Quart. Journ. Med.*, 1912, vol. vi, p. 82.