On January 2nd, 1919, Captain E. J. Coombe took over the case. The patient lay in a lethargic state, curled up; the head was retracted; very wasted. When awakened was fairly intelligent, but he fell asleep immediately he was left alone. "No convulsions; no 'cephalic cry.'" Intense pain on moving head. Slight headache at times. Pupils widely dilated. Paralysis of lower limbs, incontinence of urine and faeces. Ophthalmoscopic examination (artificial dilatation of pupils not necessary): Optic disc pale; blood vessels engorged. Diagnosis: Encephalitis lethargica.

On January 3rd lumbar puncture was performed, and 20 to 30 c.cm. fluid withdrawn; it was under great pressure, milky, and somewhat opalescent. Ten grains of urotropin in 10 c.cm. of physiological salt solution were injected into the spinal canal. The patient improved somewhat after the lumbar puncture, and on January 5th was less lethargic.

January 6th. Report on cerebro-spinal fluid: Milky; 20 c.cm. centrifugized gave a greenish deposit $\frac{1}{2}$ in. in depth. Polynuclear leucocytosis of 1,000. Stained films (Gram and Leishman) showed a minute coccus Gram-positive, chiefly extracellular, but occurring occasionally intracellularly; usually diplococci and having apparently no capsule. Seen also in short chains of 3 to 5 cocci. All attempts at growth on agar, blood serum, Loeffler's serum, and haemoglobin agar failed. Anaerobic cultures could not be tried as the laboratory has not the equipment.

On January 9th it was noted that the patient slept continuously and was unable to speak when awakened. Lumbar puncture: fluid under pressure less than before; 15 to 20 c.cm. withdrawn and 10 c.cm. saline solution containing 10 grains urotropin in solution injected. The patient collapsed with respiratory failure. Ether was injected and artificial respiration performed for thirty minutes before respiration was restored. Urotropin gr. 10 was ordered thrice a day. The condition remained almost unchanged. On January 12th the cerebro-spinal fluid from lumbar puncture was almost clear; a minute deposit on centrifugization. Stained films showed few pus cells present and no cocci.

On January 15th the lethargic condition was less pronounced; he was able to say "Yes," and recognized people when awake. The paralysis and incontinence had not improved. From this date the patient gradually sank, the bronchitis increased, and he died on January 29th, 1919. Permission for autopsy was refused.

CASE II.

Pte. J. M., aged 16 years, one month's service, was admitted to Fargo Military Hospital, Salisbury Plain, on October 27th, 1918. Temperature 103°, pulse 104, respirations 32. He complained of headache, pains in the abdomen and limbs, and frequent cough.

October 28th. Coarse râles at bases of both lungs. Symptoms of early consolidation at the right base. His condition on October 30th was unchanged, and 5 c.cm. polyvalent vaccine was given.

On November 10th he complained of pains in head and joints, and became very collapsed and pulseless. Hypodermic injection of strychnine (gr. $\frac{1}{2}$) given. Vomited brown fluid. Marked retraction of head and rigidity of neck (*tache cérébrale*). Kernig's sign present; 0.5 c.cm. vaccine given. Ice-bag to head. There was slight improvement on November 12th. He was conscious, had no headache, and no vomiting. Kernig's sign still present. Rigidity of neck persisted; complained of intense headache.

November 15th. Sputum purulent; cough loose; moist râles all over both lungs. Headache less severe.

November 18th. Headache improving. Answers questions intelligently. Moist râles all over both lungs.

November 19th. Kernig's sign still present. Dullness at right base and left upper and lower lobes. On November 21st 10 c.cm. antistreptococcal serum were given.

December 3rd. He has periodic attacks of severe occipital headache, with occasional attacks of vomiting. Has become very emaciated. The temperature varies between 101° and 104°.

December 5th. Vomiting of "cerebral type." Persistent severe headache. "Cephalic cry." Knee-jerks sluggish. No ankle clonus.

January 2nd, 1919. Temperature 101°, pulse 108, respirations 24. Patient emaciated. Pupils dilated, reacting sluggishly to light. Head retractions. Complaints of frequent headache. Slightly drowsy, but intelligent. Intense pains on moving neck. Incontinence of urine and faeces. Occasional "cephalic cry." Ophthalmoscopic examination: Pale disc; engorged vessels; pupils dilated sufficiently for examination. Calomel gr. $\frac{1}{2}$ thrice a day was ordered.

January 3rd. Lumbar puncture; fluid under pressure; 20 c.cm. withdrawn and 10 c.cm. saline solution containing 10 gr. urotropin injected. The cerebro-spinal fluid was colourless. The fluid was examined on January 6th. It was clear, and contained a few polynuclear leucocytes, but no lymphocytes. Stained Gram and Leishman. A few minute cocci were seen, Gram-positive, generally extracellular, sometimes intracellular; no capsule observed; usually diplococcus but sometimes in short chains, 3 to 5 cocci. The patient improved after lumbar puncture. Temperature 99°, pulse 104, respirations 24. Headaches less frequent. Paralysis and incontinence as before. Appetite good.

January 9th. Lumbar puncture; fluid under pressure and milky; 10 c.cm. saline solution containing 10 gr. urotropin injected. Urotropin gr. 10 thrice a day was ordered. On January 12th he was better, and wanted to get up, had no headaches and less pain on moving the head. Paralysis and incontinence

remained. Stained films from the cerebro-spinal fluid showed many pus cells and no cocci. On January 16th the condition was unchanged; but on January 21st he was very lethargic, inarticulate, and very emaciated. Incontinence of urine and faeces persisted, and he had difficulty in swallowing fluids.

February 3rd. Died. Permission for autopsy was refused.

The symptoms in both patients exhibited a remarkable similarity.

The persistent headache and head retraction suggested tuberculous meningitis, but there was no paralysis of cranial nerves, and the differential blood counts showed no lymphocytosis. One common symptom was the widely dilated pupils, at first reacting sluggishly to light, but later there was no sign of contraction, even during examination of the fundi. The appearance of the fundi in both cases was remarkable; instead of the usual signs of the onset of optic neuritis the discs in both were somewhat pale, with clearly defined edges and marked congestion of the retinal veins. The patients, when not suffering from severe headache, were so lethargic that they hardly seemed conscious that the fundus was being examined, and on one occasion Pte. M. was found to be soundly asleep at the conclusion of the examination.

Pte. L. showed, in the earlier stages of the disease, more pronounced symptoms of cerebral irritation than Pte. M. He lay curled up on his left side with his head beneath the bedclothes, but in the later stages both lay on their backs, sleeping quietly. At no time was stertorous breathing observed in either case, nor did either patient suffer from convulsions. Paraplegia and incontinence of urine and faeces supervened at an early stage in both cases and persisted until death.

Lumbar puncture gave temporary relief from headache in both cases, but, as already noted, Pte. L. had respiratory failure on one occasion after removal of 15 to 20 c.cm. of fluid and injection of 10 c.cm. normal saline solution containing 10 grains urotropin. This treatment was tried as a last resort in the hope that it would destroy any infective organism. The fact that urotropin is inert in neutral or alkaline media makes it improbable that it would have any action on micro-organisms suspended in the cerebro-spinal fluid.

The onset of encephalitis lethargica in two typical cases of influenza suggests that the minute Gram-positive coccus isolated by Captain J. A. Wilson in cases of influenza and the apparently identical organism found by one of us (J. R. C.) in both of these cases, is the probable cause of encephalitis lethargica, as well as of the other diseases enumerated in the preliminary report. Subsequent lumbar punctures showed no trace of this coccus, or the cerebro-spinal fluid would have been sent to a larger laboratory for more detailed investigation.

It is recognized that in the absence of *post-mortem* observations and of any result from the attempts to obtain cultures of the coccus this report is incomplete, but the unusual clinical symptoms which appear to differentiate this disease from various forms of meningitis and from other types of encephalitis may assist in the diagnosis of other cases.

REFERENCES.

- ¹ Von Economo, C.: *Die Encephalitis Lethargica*. Leipzig: Franz Deuticke, 1918. ² BRITISH MEDICAL JOURNAL, February 1st, 1919, p. 127.

HYPNOSIS, SUGGESTION, AND DISSOCIATION.

BY

WILLIAM BROWN, M.A., M.D. OXON., D.Sc. LOND.,

LATE MAJOR R.A.M.C.,

READER IN PSYCHOLOGY IN THE UNIVERSITY OF LONDON,
KING'S COLLEGE.

THERE is an overwhelming consensus of opinion at the present day that the one satisfactory method of treating the various forms of functional nervous disorder is that of mental analysis and re-education. The physician endeavours to unravel the tangled emotional skein of the patient's past life, as well as the emotional circumstances which ushered in the disease in its manifest form. He reveals mental conflicts and misunderstandings of the past, and mal-adaptations to the present, which he then helps the patient to correct and to solve. He demonstrates the causal connexion of these mental antecedents with the physical or mental symptoms from which the patient is suffering. He thus helps the patient to understand

himself, practically as well as theoretically. When this understanding is complete and is acted upon by the patient, the latter is cured.

To sum up this method, I would suggest the term "autognosis," or self-knowledge. It embraces the two conceptions of analysis and re-education, and is free from the implications which the term psycho-analysis carries at the present day.

The question then arises whether the old-time methods of suggestion and hypnosis can find any place within the autognostic method. The principal use to which hypnosis was put before the war was as a means of reinforcing suggestion. Having hypnotized the patient, the physician proceeded to suggest away his various functional symptoms, and, to make doubly sure, often gave post-hypnotic suggestions to produce other general and specific effects. This method is undesirable for several reasons. In the first place, it treats the symptoms and not the cause. The same or other symptoms may therefore appear later on. Secondly, the patient is passive and abnormally suggestible. This increased suggestibility persists after the treatment, and is a source of mental weakness to the patient. Thirdly, repeated hypnosis, which is necessary in the more severe cases, tends to produce automatism and somnambulism. If the hypnosis is so deep that the patient loses touch with his waking consciousness and awakes as from a dream, these defects are serious and are only to be accepted as a choice between two evils—for example, in cases of alcoholism of long standing, or in other drug habits that have resisted all ordinary methods of treatment. In light hypnosis, where the patient has reached the arbitrarily defined stage at which he is unable to open his eyes when the operator tells him that he cannot, but is otherwise fully conscious of his surroundings, the drawbacks are correspondingly less.

But hypnosis can be used with a different purpose from that of reinforcing suggestion. All the severe cases of "shell shock" of the hysterical type (that is, showing functional disturbance, or loss, of sensory or motor powers) which I saw near the firing line in France suffered from loss of memory. I used light hypnosis to clear up this amnesia, and, as might be expected, I invariably succeeded. At first I used the method in the ordinary way, as an aid to the employment of suggestion, and suggested away the various other symptoms—mutism, paralysis, contracture, tremor, etc. But I soon discovered that if, after hypnotizing the patient, I suggested the recall of the lost memories with hallucinatory vividness, and reinstated the accompanying emotion of fear as an actual present experience, the voice returned of itself and the paralysis, etc., disappeared or diminished, without any suggestions directed towards these symptoms. The extent to which the symptoms disappeared varied in proportion with the extent to which I succeeded in reinstating the original emotional experience.

To illustrate by an example:

A patient, Pte. J. A., aged 41 years, was admitted to my neurological wards on the Somme suffering from deafness, mutism, and right-sided paralysis. His right arm was held stiffly across the chest, and he walked with the right leg stiff and the right foot turned outwards. There was also slight paralysis of the right side of the face. The tendon reflexes were exaggerated, but the plantar reflex was flexor and there was no ankle clonus. The abdominal reflexes were present, and equal on the two sides. The tongue was steady and protruded in the middle line.

The patient's history was as follows: He was partially buried by a bomb that went over his left shoulder and burst just behind him, causing him to lose all control of himself. He was then sent to a field ambulance, where he remained for four days and then returned to duty. On arrival at his unit he was noticed to be still tremulous, and was sent to see his regimental medical officer. Just as he reached the aid post a shell exploded near him and rendered him unconscious. On regaining consciousness he could neither hear, speak, nor walk. It should be added that, about six weeks before this, he had received a blow on the right arm from another man's shovel while trench digging. Since then his right arm had been tremulous whenever he was under shell fire.

I got the patient to lie down on a couch, and then showed him written instructions to close his eyes and think of sleep. He was instructed to relax all his muscles, breathe slowly and deeply, and think only of sleep. After a few seconds his breathing and facial expression showed me that he was asleep. I then made a sudden loud noise by banging two books together. His eyelids flickered and I found that he could hear.

I sent him to sleep again, and gave him the suggestion that the moment I put my hand on his forehead he would again feel

the explosion of the shell which knocked him down near the aid post, and would seem to be going through the same experience again, exactly as it happened. The moment my hand touched his forehead he became very tremulous, breathed rapidly, and then shouted out: "Feeny, Feeny, kill them! . . . Feeny, Feeny, where are you? It's getting too hot here. I can't put up with it any longer. . . . Feeny, come here and get me out, I can't move my legs. I can't stand up. [Very rapid breathing.] I can't walk to the dug-out. Take hold of my arms, then I shall manage to get along. . . . I don't want to lie on that wire bed. I want to get out of this. We have had everything from a rifle bullet to a Jack Johnson over here to-day, and three parts of us are knocked out." While the patient was talking like this, his right arm and leg showed signs of movement. Before waking him I reminded him where he was, who was speaking to him, and the principal facts about his admission to hospital, thus reassociating his hypnotic dream with his present experience. I also told him that on waking he would remember everything that he had just been going through. I then woke him up, and found that he could speak (although with a stammer), and could walk about normally. His right arm was still tremulous, and showed marked extensor paralysis.

Some days later I again sent him to sleep, and suggested that he would again live through his experience of being hit on the right arm by his comrade's shovel. He showed signs of pain, seized hold of his right upper arm with his left hand, writhed, and shouted "Oh my arm! . . . put it in my pocket." He then put his hand in his pocket. His hand was trembling. Thereupon I sent him more deeply to sleep. He then withdrew his hand from his pocket; it was quite steady.

In the case of this patient, then, the revival of memories of his two accidents, in a state of light hypnosis, with such vividness that he again felt the same emotions and seemed to be living again through the actual experiences, restored his voice and to some extent the motor power in his right arm and leg, and abolished the tremor of his right arm. All this occurred without the use of specific suggestions. But I had to help myself out later on with specific suggestions, as the case was more than usually difficult, nor could I keep him long enough to complete his cure.

During a period of sixteen months 121 cases of mutism passed through my neurological wards. Every one of them spoke when made to live again through their terrifying experiences. Some of them stammered, but these were a small minority. I did not need to give a specific suggestion that they would be able to speak. Cardio-vascular and secretory changes (tachycardia, hyperidrosis, etc.), dependent upon disturbance of the sympathetic nervous system, showed alteration under the same method of treatment. Curiously enough, paralysis of the voluntary muscles (apart from those concerned with the voice) diminished less readily than other symptoms. On the other hand, voluntary paralysis cleared up, sometimes quickly, sometimes slowly, under the influence of suggestion and persuasion, without the need of hypnosis.

These are facts. What conclusion can be drawn from them? It seems to me that we have a twofold dissociation in these cases. In the first place the shock of the shell explosion has produced a dissociation from the patient's personal consciousness of certain psycho-physical functions together with the memories linked with them immediately after the shock. But a second dissociation has also occurred, namely, that between the psychical and physical counterparts of the emotional reaction of fear. The physical counterpart then persists instead of being evanescent. The first dissociation has to do with the sympathetic. Revival of emotion with hallucinatory vividness soon after the original shock, abolishes the second dissociation as well as the first, and so brings the physical manifestation of the emotion again under the sway of the conscious personality.

In cases seen in England and Scotland I found that lost memories were not so easily recalled, and that the reinstatement of emotion with hallucinatory vividness was very difficult to effect. However, in the few cases in which this did occur, the symptoms showed definite and sudden improvement. Chronic cases pass through a period of later mental development, in which different mental processes, memories and motives, recent and remote, interact and produce the well-known fixation of symptoms. We are thus brought back to the method of autognosis, in which memories are revived, in reverse order, by patient questioning and heart-to-heart talks. But it is the experience acquired in the field, with early and what I would call relatively "pure" cases, that

enables us to understand the therapeutic value of the revival of memories; and in light hypnosis we have a helpful method, if judiciously and sparingly used, to speed up the process or make it sufficiently complete. It is only in a small minority of the cases that this is needed or advisable.

TRANSVENOUS ANEURYSMORRHAPHY.

DESCRIPTION OF THE OPERATION.

BY

WILLIAM PEARSON, M.D., B.Ch., F.R.C.S.I.,

SURGEON, ADELAIDE HOSPITAL, DUBLIN; LATE MAJOR R.A.M.C.
(TEMPORARY), AND A DISTRICT CONSULTANT SURGEON.

I wish to draw attention to an operative procedure which I have employed in the treatment of aneurysmal varix with most gratifying results, and which I have termed "transvenous aneurysmorrhaphy." It appears to meet the needs of a certain number of cases in an admirable manner.

I will first give an account of the case, which presents some points of peculiar interest, and will then offer some observations in regard to the operation itself.

CASE.

Pte. H. M., aged 22, was wounded in the right thigh by a rifle bullet on October 1st, 1916. On October 10th he came under the care of Captain P. Cheal, R.A.M.C., who found a small healed entrance wound on the antero-internal aspect of the thigh 4 in. above the knee-joint, with an exit wound at the same level on the posterior surface not quite healed. Considerable persistent swelling developed in the leg and foot, and was considered to be due to venous thrombosis. There were no signs of aneurysm. On January 5th, 1917, massage was instituted, after which a very marked vibratory thrill was observed for the first time in the neighbourhood of the wound. On January 6th I saw the case in consultation with Captain Cheal. The foot and leg were dusky and showed slight general oedema, and there was definite dilatation of the superficial veins from the knee downwards. In the upper part of the popliteal space there was slight swelling, with systolic pulsation, and other signs of arterio-venous aneurysm were present. There was faint pulsation in the anterior and posterior tibial arteries. I regarded the case as one of varicose aneurysm, and advised operation.

Description of Operation.

The operation was performed on January 15th, 1917. The popliteal vessels were exposed in the upper part of their course where the artery and vein were firmly adherent to one another without any intervening sac.

FIG. 1.—Aneurysmal varix of popliteal vessels, with a sacculated venous aneurysm on the opposite wall of the vein.

Here the vein was greatly dilated and the artery below was considerably smaller than usual. When freeing the side of the vein remote from its connexion with the artery a saccular venous aneurysm was exposed (see Fig. 1) and was removed, disclosing the arterio-venous stoma through the lumen of the vein. It was of elongated elliptical form, $\frac{3}{4}$ in. in length, and lying in the long axis of the vessel. The opening in the vein was enlarged very slightly upwards and downwards, and through it the aneurysmal opening was sutured (see Fig. 2) with fine silk sterilized in liquid paraffin. The vein was then closed in a similar manner.

On removal of the tourniquet there was no bleeding and the circulation through the vessels seemed perfect. Operation was completed by suture of the fascial tissues and the skin.

After operation good pulsation was felt in the dorsalis pedis and posterior tibial arteries. On the following day there was no trace of oedema in the limb, the colour was normal, and the dilatation of the superficial veins had disappeared. Healing was uneventful, the circulation remained good, and the patient was allowed to walk four weeks after operation. He was kept under observation for two months longer and was then discharged to light duty, the circulation being normal.

It is remarkable that until massage was instituted, three months after he was wounded, the patient exhibited no signs of arterio-venous aneurysm, although there were evidences of venous obstruction present from the beginning. I have previously¹ drawn attention to the rapid development of traumatic aneurysm following massage in certain cases of gunshot injuries, and it would appear that massage was the exciting cause here also, though naturally the vascular injury dated from the time of the wound, and

it is probable that in the absence of massage active movements would have produced the same result. A reasonable explanation of the signs previously present is that the arterio-venous opening and the lumen of the vein were occluded by a clot which was disturbed by massage, but in that case it is difficult to understand why no evidences of embolism appeared, and at operation the lumen of the vein appeared perfectly smooth and normal. Consequently I find it hard to account for the train of events in this case. Another point of interest is the presence of a distinct venous aneurysm which accounted for the swelling and pulsation at the upper end of the popliteal space, and led to the diagnosis of varicose aneurysm rather than of aneurysmal varix.

A few words are necessary in regard to the technique of this operative procedure:

Apart from the necessity for rigid asepsis, scrupulous care must be exercised to avoid injury to the delicate intima of the vein. Consequently any blood which may obscure the view on opening the vessel must be washed away by gentle irrigation with saline solution, which is then carefully mopped up by a gauze swab or aspirated by a syringe. On no account must a swab be introduced into the vein for the purpose of wiping off blood; such a measure will favour subsequent venous thrombosis. The vessel wall should be smeared with sterilized vaseline, as recommended by Carrel.

The method of introducing the suture to close the arterio-venous communication is important: The needle is first passed through a small portion of the fibrous tissue which binds the vessels together at one end of the communication, on its external aspect (that is, entirely outside the vessels and between the two), and does not penetrate the intima of either—it thus resembles the commencement of the outer row of sutures in a gastro-jejunostomy. This stitch is immediately tied and the needle is then passed obliquely from without inwards, emerging on the inner surface of the vein close up to the extremity of the stoma (see Fig. 2). The extreme edges of this are then brought together by fine continuous sutures through intima and media until the opposite end

FIG. 2.—The venous aneurysm has been removed and the opening in the vein slightly enlarged, its edges being retracted by fine traction stitches. The arterio-venous stoma is being closed by a fine continuous suture. Note that no knot appears on the interior of the vessels (see text). In practice the stitches do not lie so close together as figured, and when drawn taut they do not present so much on the surface.

is reached, when the needle is again passed out from the lumen of the vein obliquely through its wall and the overlying tissue which binds it to the artery, so as to emerge between the two, when it is secured in a similar manner to the other end. Thus it will be seen that there is no knot projecting in the lumen of either vessel, and if the stitches are placed accurately and drawn taut throughout they do not present in the lumen of either vessel any more than in the case of ordinary through-and-through stitches which produce eversion of the edges in vascular suture. The closure of the incision in the vein presents no difficulty, being effected by the ordinary method advocated by Carrel.

While many cases of aneurysmal varix cause little or no inconvenience to the patient and do not require operation, others demand treatment by reason of circulatory disturbances. In some of these, when conservative and reconstructive measures are desirable, the procedure I