INTRAVENOUS INJECTIONS OF ANTIMONIUM TARTARATUM IN KALA-AZAR.*

BY

GEORGE C. LOW, M.A., M.D., C.M., M.R.C.P.,

TEMPORARY MAJOR I.M.S.,

PHYSICIAN, ALBERT DOCK HOSPITAL, LONDON SCHOOL OF TROPICAL MEDICINE; LECTURER, LONDON SCHOOL OF TROPICAL MEDICINE AND KING'S COLLEGE (UNIVERSITY OF LONDON).

IN the *Transactions* of the Society of Tropical Medicine and Hygiene for December, 1916 (vol. x, No. 2, pp. 37-42), I briefly described the history of the introduction of intravenous injections of tartar emetic in trypanosomiasis, leishmaniasis, and other tropical diseases. It is unnecessary, therefore, to enter upon this again, apart from saying that the beneficial effects of the drug have been well proved since that date, especially in leishmaniasis.

The case I now report is of special interest, first, because the infection was acquired almost certainly in Baghdad, where this disease does not apparently exist indigenously; secondly, because it is the first case I know of which has been treated in England by this method; and, thirdly, because an unfortunate attack of influenza, by carrying the patient off just at the termination of his treatment, gave the opportunity of definitely determining whether or not a complete sterilization of his *Leishmania donovani* had taken place.

History of the Case.

The patient was born in England, and was never out of it till 1916. Joining the army in the end of that year, he served a month in France and then was transferred to India. He arrived there in March, 1917, and was stationed first at Bombay then at Karachi. While there he went for a trip to Bangalore and back, but, apart from that, had visited no other parts of India. After this he went to Mesopotamia, arriving in Baghdad on July 30th, 1917, where he stayed until invalided in June, 1918, to Bombay and then home to England.

Previous Illnesses.—Influenzal cold, 1913; inflammation of antrum and otitis after it; no pus; never venereal disease.

Present Illness.—The patient, as just stated, arrived in Baghdad on July 30th, 1917. He enjoyed good health until the following March, when, on the 25th of that month, he felt feverish and took his temperature, which was found to be raised. After this night sweats began to develop, so on April 19th, 1918, he went into hospital. His temperature was then found to be 102° F.; spleen enlarged. Blood films: no malaria or relapsing fever; three blood cultures for the enterica group all negative. No undulant fever reaction. The illness went on, and quinine had no effect upon the temperature. Leucocytes then 10,000. At this time *Entamoeba histolytica* cysts and *Lamblia intestinalis* cysts were found in the stools. For this he had two courses of emetine, one by hypodermic injection, the other a combined one of emetine hydrochloride hypodermically and emetine bismuth iodide by the mouth. Two negative examinations for *histolytica* cysts were recorded after this. At the end of May, 1918, the liver was punctured but nothing was found. As the pyrexia and other symptoms continued the patient was sent to Basra on June 16th, 1918, then to Bombay, and finally, on August 29th, 1918, home to England.

On arrival he was admitted into St. Thomas's Hospital, under the care of Captain A. E. Russell, R.A.M.C., on September 29th, 1918. I saw him with Captain Russell on October 16th, and on examining smears of spleen juice that had been taken by splenic puncture found *Leishmania donovani* parasites. The patient's blood count at that time was as follows: Red cells 3,300,000; white cells 1,500; haemoglobin 60 per cent. He was running a remittent temperature, had a large spleen and all the other clinical signs were very typical of kala-azar.

Tartar emetic, intravenously, in small doses, was then begun. I saw him again on November 10th, and found that the temperature was still keeping high, and the patient was depressed and looking ill. On November 19th the patient was transferred to the Albert Dock Hospital and passed under my charge. I at once pressed the antimony, and after the first big dose of 2 grains the temperature fell for good and never rose again, with the exception of a rigor, once or twice, after some of the injections. (I am greatly indebted to Captain Russell for allowing me to use his notes and for giving me permission to publish the early part of the case.)

The following table gives the details of the treatment at a glance.

Table showing Progress of Case and Drug Employed.

Date.	Antimony Tartrate.	Date.	Antimony Tartrate.
1918.		1918.	
Oct. 18 ...	gr. 1 intravenously	Dec. 23 ...	gr. 2 intravenously
" 22 ...	gr. 1	" 27 ...	gr. 2
" 25 ...	gr. 1	" 30 ...	gr. 2
" 29 ...	gr. 1		
Nov. 1 ...	gr. 1	1919.	
" 5 ...	gr. 1	Jan. 2 ...	gr. 2
" 12 ...	gr. 1½	" 6 ...	gr. 2
" 15 ...	gr. 1½	" 9 ...	gr. 2
" 21 ...	gr. 2	" 13 ...	gr. 2
" 25 ...	gr. 2	" 16 ...	gr. 2
" 28 ...	gr. 2	" 20 ...	gr. 2
Dec. 2 ...	gr. 2	" 23 ...	gr. 2
" 5 ...	gr. 2	" 27 ...	gr. 2
" 9 ...	gr. 2½	" 30 ...	gr. 2
" 12 ...	gr. 2½	Feb. 3 ...	gr. 2
" 16 ...	gr. 2½	" 6 ...	gr. 2
" 19 ...	gr. 2½	" 10 ...	gr. 2

Notes to Table.

November 12th, 1918. Blood count: Red cells 3,900,000; white cells 1,480; haemoglobin 85 per cent. Down to this date at St. Thomas's Hospital.

November 19th. Transferred to Albert Dock Hospital, London School of Tropical Medicine.

December 6th. Blood count: Red cells 4,450,000; white cells 4,500; haemoglobin 90 per cent. The temperature fell to normal after the first 2 gr. dose and remained so.

December 27th. Gaining weight; spleen gone down very much.

January 6th, 1919. White cells 4,800; spleen still palpable, though much reduced in size.

January 23rd. No sign of disease left now.

January 30th. White cells 4,000; Up and about; spleen just palpable under costal margin.

February 13th. Influenza.

February 17th. Died of influenza pneumonia at 11.30 p.m. Amount of antimony given: At St. Thomas's 7½ gr.; at Albert Dock Hospital 5½ gr.; Total 6½ gr.

By February 10th, 1919, just before the patient developed influenza, he was apparently in perfect health, and all signs of the disease had disappeared. He was up and about, eating well, going out and to all intents and purposes clinically cured.

The attack of influenza was severe. He felt tired on the night of February 12th. By next morning he had developed a temperature of 103°, but no lung symptoms. On the 14th, 15th, and 16th the temperature was still high, but the strength was well maintained. On the morning of the 17th a patch of pleurisy developed on the right side, just above the liver, and by the afternoon definite pneumonia had set in. This spread with alarming rapidity, and the patient sank and died at 11.30 p.m. from the severe toxæmia.

An autopsy was allowed and this was carried out on February 19th, at 10 a.m., thirty-four and a half hours after death.

Autopsy Notes.

External Appearances.—Post-mortem staining already well advanced, black liquid exuding from nose and mouth.

Thorax.—No fluid in pericardium, no pericarditis.

Heart: Contained ante-mortem clot in right ventricle. Organ ordinary size; no lesions of valves; muscle fairly hard, not specially friable; shows cloudy swelling. Aorta, no atheroma.

Lungs and pleural cavities: A few ounces of blood-stained fluid in right pleural cavity, none in left; no old pleural adhesions. Right lung: Recent pleurisy with lymph over lower lobe. Whole of this lobe solid with confluent bronchopneumonic patches, also oedematous. Upper and middle lobes congested, no pneumonia. Left lung: Commencing bronchopneumonia in lower lobe; congestion in upper. No old tubercle in either lungs.

Abdomen.—No peritonitis.

Liver: Weight 6 lb. 5 oz. No perihepatitis. Organ in a state of what appears to be an early multilobular cirrhosis, markedly fatty also; yellow colour; not much blood in organ.

Spleen: Weight 1 lb. 9 oz.; enlargement chiefly upwards. Firm to the feel; pulp firm; darkish colour, no perisplenitis. Kidneys: Cloudy swelling and fatty changes; no nephritis. (Albumin was never present in the urine during life.)

Stomach: Contained some black grumous fluid; mucous membrane, patchy congestion at places.

Small intestine: Nothing definite to note.

Large intestine: A series of small pigmented scars in caecum, chiefly round opening of appendix; look like healed ulcers, very small—2-4 mm. only across. (Were those amoebic ulcers which gave rise to the *Entamoeba histolytica* cysts found in the faeces in Baghdad?) No large ulcers in colon. No sign of kala-azar ones. Some small deeply congested areas near rectum. Mesenteric glands congested and slightly enlarged.

Groin glands palpable, slightly enlarged.

Bone marrow very dry, normal colour.

* Published by permission of Sir R. Havelock Charles, G.C.V.O., Medical Adviser to the Secretary of State for India.

Smears were taken from the liver, spleen, bone marrow, mesenteric lymph glands, inguinal lymph glands, kidneys, lungs, and heart. Though a very exhaustive search was made in the many smears taken from these different organs, nothing resembling *Leishmania donovani* was seen.

In sections of the same organs a similar absence of parasites was noted, this proving definitely that the specific parasite had really disappeared. Sometimes an infection with *L. donovani* is cleared out after any severe septic change, and as the patient had suffered from a severe toxæmia from his influenzal pneumonia just before death, the possibility of this having something to do with the disappearance of the parasites, though unlikely, must not, of course, be forgotten. From the disappearance of all clinical signs early in the treatment after the large doses were employed, it may be surmised that the sterilization took place about that time or a little later, and at the time the influenza developed there is little, if any, doubt that he was quite free from all infection.

The autopsy in this case, in addition to yielding valuable information as to the destruction of the specific parasite of kala-azar, is also of importance in enabling one to determine what effect, if any, large doses of antimony intravenously have upon the different viscera. Can one produce a state of chronic poisoning by too large and too prolonged doses?

Breil and Priestley¹ report a case of poisoning in an aboriginal Australian, 22 years of age, suffering from ulcerating granuloma.

Treatment in this case was begun with doses of 0.08 gram of tartar emetic in normal saline, administered intravenously, increasing after two days to 0.1 gram, and after a further two days to 0.12 gram; injections were given on alternate days, until altogether 1.74 grams had been given.

The day following the last injection the patient complained of slight nausea and vomiting and looked ill. On the third day after he became delirious, had a rise of temperature, and vomited dark blood. Death took place sixty-four hours after the final injection. Diarrhoea did not occur, nor was blood and albumin noted in the urine.

At the autopsy the liver was found to be of normal size; microscopically a widespread fatty degeneration was present, with here and there acute focal necroses. The kidneys showed lesions corresponding to an acute interstitial nephritis with calcareous deposits in the medullary substance. The heart muscle showed slight fatty changes; the stomach was oedematous and congested, but did not show any ulceration. No chemical analysis was made.

Analysing this case, the kidneys would appear to have been at fault, and the excretion of the drug was probably hindered on account of this. The total antimony given was not large, only 1.74 grams. The symptoms were those of an acute rather than a chronic poisoning.

While engaged in writing up the details and consulting the literature of my case an opportunity of studying a similar case, I note, has occurred to Archibald and Innes in Egypt.²

These observers report a case of bilharziosis treated by intravenous injections of tartar emetic, death taking place at the end of a course of 33 grains from influenzal pneumonia. In this case the liver weighed 3½ lb., and was of a pale yellow colour; microscopically the liver cells were granular and vacuolated, and areas of necrosis were seen (there is no mention of any cirrhosis being present). The kidneys showed cloudy swelling and fatty degeneration and infiltration.

In discussing the question of poisonous effects of antimony the authors cite Knowles as recording 5 deaths out of 20 cases of kala-azar treated by this method. No details are given, however, as to whether these were due directly to the antimony or not. Archibald and Innes's case evidently died, like my case, from influenzal pneumonia, and the question of whether the antimony had anything to do with the fatal issue is *sub judice*.

In the case described above it will be remembered that the liver showed a well marked cirrhosis, was double the normal size (6 lb. 5 oz.), and to the naked eye was yellow and distinctly fatty. Was this condition, then, due to the kala-azar (cirrhosis of the liver has been described in this disease), or was the fatty change produced by the toxæmia of the influenzal pneumonia, or was the cirrhosis and fatty change due to the large doses of antimony given from time to time?

In order to determine this, careful microscopic examinations of sections of the liver were made and studied by Dr. Newham, myself, and Major Foulerton, R.A.M.C., who

kindly took some of the tissue for analysis. These sections show considerable fatty infiltration of the parenchymatous cells of the liver, and also considerable fibrous changes of a somewhat irregular type. As Major Foulerton is now working at the poisonous effects of trinitrotoluene, tetrachlorethane, and the salvarsan group on the liver and other viscera, I have handed this aspect of the case over to him and he will report upon the matter in detail later.

Apart from the liver and kidneys, the other viscera showed no changes of importance. The stomach was perhaps slightly congested; it showed no ulceration. The small intestines showed no appreciable change, while all that was visible in the large intestine were some small pigmented spots suggestive of healed ulcers around the ileo-caecal valve and a few congested areas near the rectum. The former may have been old amoebic ulcers in the early stage, and may have been the source of the *E. histolytica* infection detected and successfully treated in Baghdad. Apart from these the large gut showed no sign of any visible lesions. No *E. histolytica* (free or cystic forms) were found in the stools when the patient was under my care. He had a moderate *E. coli* infection, and on one occasion four coccidia (Isospora) were found in the stool. No trace of where these came from could be detected on a close examination of the gut, even with a hand lens.

It is thus evident that antimony given in large doses intravenously over long periods of time may produce fatty changes in the viscera, and that in other cases toxic symptoms of an acute nature may appear. Recently Christopherson³ has drawn attention to the fact that tachycardia has been met with in cases in Egypt undergoing antimony treatment for bilharzial disease. I have a case of that disease under my care now, and several times after a 2-grain intravenous injection of tartar emetic the patient has complained of a feeling of constriction over the cardiac area. As he has also developed some indigestion, I have stopped further injections for the present.

As regards kala-azar, what one requires is a test to determine when the leishmanial parasites have disappeared. Whenever one can be certain of this, then the injections should cease, as they can then do no further good, while, on the other hand, they may do considerable harm.

Rogers relies on the absence of fever for a considerable time, steady gain in weight, and diminution in the size of the spleen as indicating recovery and cure, and until some better test can be evolved we may, more or less, accept this.

Kala-azar is a deadly disease, and it is justifiable to push the antimony even to a dangerous extent. The fatty changes that have been described in the few cases that have come to autopsy might quite well have disappeared if the patients had not been carried off by intercurrent disease, and it is very likely that all cases show these changes to a certain extent but recover from them when the drug is stopped. The degree of fatty change in the liver is much more extreme in ankylostomiasis, yet all these cases do not die, but apparently recover perfectly, once the parasites have been removed.

As regards bilharziosis, antimony intravenously is now on trial, and reports so far would seem to indicate that it is of value. Here again a test is required to enable one to say if the infection has disappeared or not, and so to obviate the danger of going on too long with the antimony. Archibald and Innes have suggested that the disappearance of the eosinophilia might be used for such a purpose. It remains to be seen whether this will disappear at once on the death of the adult worms. In the case of bilharzial disease mentioned above,⁴ which I have been treating with antimony injections, all blood and ova have disappeared now for some time, but the eosinophilia persists as high as when the patient was first seen. Possibly the infection is only dormant, and may spring to life again on the cessation of treatment.

CONCLUSIONS.

1. Antimonium tartaratum is evidently a specific against *Leishmania donovani*, and in the case under discussion sterilized the patient of his infection.

2. Antimony in large doses, given over prolonged periods of time, produces fatty changes in the liver and kidneys, which may seriously damage the resisting powers of the

patient, and may even cause death (Breinl and Priestley's case).

3. Antimony should not be given for longer periods than necessary, and therefore it is important that some definite test should be evolved which will indicate when the infection of kala-azar has disappeared and the patient is cured.

4. If antimony is finally proved to destroy the *Schistosoma haematobium*, then a similar test to determine when this destruction has taken place is required.

5. In all cases where antimonium tartaratum is given intravenously the patient should be confined to bed on the day of the injection and kept there till the day after, and should be carefully watched throughout the course, any indications of gastric or constitutional disturbances at once contraindicating further injections.

6. Not more than two injections a week should be given, and the salt should not be given in too concentrated a form.

REFERENCES.

- ¹ Breinl and Priestley: Notes on a case of Antimony Poisoning. *Journ. of Trop. Med. and Hyg.*, February 15th, 1918, vol. xxi, pp. 38-39.
² Archibald and Jones: *Journ. of Trop. Med. and Hyg.*, April 1st, 1919.
³ BRITISH MEDICAL JOURNAL, April 19th, 1919. *Journ. of Trop. Med. and Hyg.*, May 15th, 1919.

TENDON TRANSPLANTATION IN DROP-WRIST DUE TO NERVOUS INJURY.

BY

VERNON PENNELL, M.A., M.B., B.C.CANTAB.,

TEMPORARY CAPTAIN R.A.M.C.,

FELLOW OF PEMBROKE COLLEGE, CAMBRIDGE; SURGICAL SPECIALIST,
WAR HOSPITALS, EXETER.

WHILE realizing the great importance of resorting when possible to nerve suture or grafting in cases of drop-wrist, it is not my intention to decry such "ideal" treatment, but to bring forward arguments in favour of a more extensive appreciation of the advantages of tendon transplantation and anastomosis under conditions where the performance of suture, etc., is not possible, or even in some cases as an adjunct to this.

In the majority of cases in which the musculo-spiral nerve has been severed as a result of gunshot wounds, the humerus has been fractured, often suppuration has supervened, and secondary nerve suture cannot be carried out for some months, even under the most favourable circumstances. In spite of vigorous massage, movements, and electrical treatment, the extensor muscles waste, and the chances of a hand and fingers which shall function as formerly is indeed small.

Most men would rather have a useful hand now than an extremely problematical perfect hand in two or three years' time. Nor does the operation of tendon transplantation performed relatively early preclude or prejudice the later performance of nerve suture.

Nor does it appear unjustifiable to perform tendon transplantation about the wrist at the same time as nerve suture of the musculo-spiral in the arm, for, by the active movements at the wrist and finger joints which follow the completion of the tendon transplantation, troublesome adhesions in the small joints of the fingers and wrist are prevented, and the forearm extensor musculature is kept in better condition for the reception of the newly formed neurons than by the most skilled massage. The contention that if tendon transplantation is performed in addition to nerve suture it will in some cases become unnecessary later is more than balanced by the fact that, properly performed, it can do no harm after nerve regeneration has taken place, and the restoration of function to the paralysed muscles will in all cases be expedited.

The theory and practice of tendon transplantation stands or falls on its results. The questions to be answered are: Will the hand be benefited by transplantation? Will it prohibit or prejudice later nerve suture should this become possible? Will the residual group of robbed muscles be so weakened as to diminish their utility to vanishing point? In my opinion these questions can be satisfactorily answered, and a verdict given in favour of tendon transplantation when the following indications are present:

1. Severance with large loss of substance of the musculo-spiral nerve.
2. Severance of the musculo-spiral nerve with much bony injury and prolonged suppuration.

3. In cases where very rapid and complete wasting of extensor muscles has supervened.

4. In all cases of division of the posterior interosseus nerve.

The indication of the last group is based on the fact that the nerve is exceedingly small, runs a very circuitous course, is difficult to find, and the fact that the carpal extensors, extensor carpi radialis longior and ulnaris, are nearly always spared. The extensor carpi radialis longior and supinator longus being supplied by the musculo-spiral before its division into radial and posterior interosseus can be utilized as an extensor of the thumb and fingers.

The commonest type of posterior interosseus paralysis is that due to severance of the nerve just below the mid-point of the forearm, resulting in paralysis of the extensor ossis metacarpi pollicis, extensor brevis, and longus pollicis, and the whole or part of the common extensors of the fingers; more commonly this is partial, and the power of one or two fingers survives. These cases do exceedingly well with tendon anastomosis. The supinator longus, detached at its insertion, can be anastomosed with the extensor ossis metacarpi pollicis, and the extensor carpi radialis longior with the common digital extensors, or the extensor longus pollicis if the finger extensors are spared. In these (partial) cases it will be seen that whereas tendon anastomosis materially improves the hand within a few weeks, it does not, *a priori*, prejudice the further performance of posterior interosseus suture, were one skilful or rash enough to attempt it later.

The complete drop-wrist from musculo-spiral paralysis presents a more serious and difficult problem. This disabling condition is primarily due to the hyperflexion at the wrist-joint. Even in the normal hand the flexor muscles of the fingers cannot exert their maximum power unless there is a moderate degree of extension at the wrist-joint. When the normal hand is voluntarily placed in the attitude adopted by the drop-wrist of musculo-spiral paralysis, it is found that the fingers cannot be clenched—indeed, they cannot be fully flexed on the palm. Much more marked, then, is the condition in the flaccid paralysis of drop-wrist. In other words, the drop-wrist (or normal hand in this position) cannot make a "fist."

It behoves us, therefore, first to correct this hyperflexion of the wrist-joint and allow full play to the digital flexors. A patient with a wrist ankylosed in line with the forearm, with free finger movements, is in a better condition to make the fullest use of his fingers than one with a drop-wrist. I do not think this point is sufficiently appreciated by English surgeons. Continental surgeons have, however, recognized this, and carried out fixation of the wrist-joint by means of an artificial ligament manufactured out of fascia lata. The drop-wrist has not been cured—that is, return of power of extension—but a permanently slightly extended wrist has resulted, allowing full play to the flexors of fingers and thumb, which later can be abducted by its shorter abductor.

In these cases a fascial graft can be manufactured to act as a ligament permanently extending the wrist. It is attached to the third or fourth metacarpal distally and to the radius or ulna proximally, slightly nearer the upper extremity than the lower. If this is done, the flexor carpi radialis can be utilized for anastomosis with the extensor communis digitorum by passing its tendon through an aperture in the interosseus membrane, or, better still, the pronator quadratus, as there is less likelihood of the formation of adhesions with the tendon passing through muscle than interosseus membrane. The palmaris longus can easily be anastomosed with the extensor ossis metacarpi pollicis, and a very fair amount of abduction of the thumb results.

The method of transplanting the flexor carpi radialis and ulnaris to the insertions of the extensor carpi radialis longior and extensor carpi ulnaris works well, but leaves extensors of the fingers and those of the thumb without suitable "tendon donors," unless the final carpal flexor palmaris longus (not always present) is utilized, or a suture between the insertion of the pronator radii teres and the belly of the extensor communis digitorum effected, a not very satisfactory plan on *a priori* grounds as the resulting muscle pulls round a "corner."

The supporters of this method claim that after suitable education the flexor carpi radialis and ulnaris function as extensors of wrist or fingers.