patient, and may even cause death (Breinl and Priestley's case).

3. Antimony should not be given for longer periods than necessary, and therefore it is important that some definite test should be evolved which will indicate when the infection of kala-azar has disappeared and the patient is cured.

4. If antimony is finally proved to destroy the *Schistosoma haematobium*, then a similar test to determine when this destruction has taken place is required.

5. In all cases where antimonium tartaratum is given intravenously the patient should be confined to bed on the day of the injection and kept there till the day after, and should be carefully watched throughout the course, any indications of gastric or constitutional disturbances at once contraindicating further injections.

6. Not more than two injections a week should be given, and the salt should not be given in too concentrated a form.

REFERENCES.

- ¹ Breinl and Priestley: Notes on a case of Antimony Poisoning. *Journ. of Trop. Med. and Hyg.*, February 15th, 1918, vol. xxi, pp. 38-39.
² Archibald and Jones: *Journ. of Trop. Med. and Hyg.*, April 1st, 1919.
³ BRITISH MEDICAL JOURNAL, April 19th, 1919. *Journ. of Trop. Med. and Hyg.*, May 15th, 1919.

TENDON TRANSPLANTATION IN DROP-WRIST DUE TO NERVOUS INJURY.

BY

VERNON PENNELL, M.A., M.B., B.C.CANTAB.,

TEMPORARY CAPTAIN R.A.M.C.,

FELLOW OF PEMBROKE COLLEGE, CAMBRIDGE; SURGICAL SPECIALIST, WAR HOSPITALS, EXETER.

WHILE realizing the great importance of resorting when possible to nerve suture or grafting in cases of drop-wrist, it is not my intention to decry such "ideal" treatment, but to bring forward arguments in favour of a more extensive appreciation of the advantages of tendon transplantation and anastomosis under conditions where the performance of suture, etc., is not possible, or even in some cases as an adjunct to this.

In the majority of cases in which the musculo-spiral nerve has been severed as a result of gunshot wounds, the humerus has been fractured, often suppuration has supervened, and secondary nerve suture cannot be carried out for some months, even under the most favourable circumstances. In spite of vigorous massage, movements, and electrical treatment, the extensor muscles waste, and the chances of a hand and fingers which shall function as formerly is indeed small.

Most men would rather have a useful hand now than an extremely problematical perfect hand in two or three years' time. Nor does the operation of tendon transplantation performed relatively early preclude or prejudice the later performance of nerve suture.

Nor does it appear unjustifiable to perform tendon transplantation about the wrist at the same time as nerve suture of the musculo-spiral in the arm, for, by the active movements at the wrist and finger joints which follow the completion of the tendon transplantation, troublesome adhesions in the small joints of the fingers and wrist are prevented, and the forearm extensor musculature is kept in better condition for the reception of the newly formed neurons than by the most skilled massage. The contention that if tendon transplantation is performed in addition to nerve suture it will in some cases become unnecessary later is more than balanced by the fact that, properly performed, it can do no harm after nerve regeneration has taken place, and the restoration of function to the paralysed muscles will in all cases be expedited.

The theory and practice of tendon transplantation stands or falls on its results. The questions to be answered are: Will the hand be benefited by transplantation? Will it prohibit or prejudice later nerve suture should this become possible? Will the residual group of robbed muscles be so weakened as to diminish their utility to vanishing point? In my opinion these questions can be satisfactorily answered, and a verdict given in favour of tendon transplantation when the following indications are present:

1. Severance with large loss of substance of the musculo-spiral nerve.
2. Severance of the musculo-spiral nerve with much bony injury and prolonged suppuration.

3. In cases where very rapid and complete wasting of extensor muscles has supervened.

4. In all cases of division of the posterior interosseus nerve.

The indication of the last group is based on the fact that the nerve is exceedingly small, runs a very circuitous course, is difficult to find, and the fact that the carpal extensors, extensor carpi radialis longior and ulnaris, are nearly always spared. The extensor carpi radialis longior and supinator longus being supplied by the musculo-spiral before its division into radial and posterior interosseus can be utilized as an extensor of the thumb and fingers.

The commonest type of posterior interosseus paralysis is that due to severance of the nerve just below the mid-point of the forearm, resulting in paralysis of the extensor ossis metacarpi pollicis, extensor brevis, and longus pollicis, and the whole or part of the common extensors of the fingers; more commonly this is partial, and the power of one or two fingers survives. These cases do exceedingly well with tendon anastomosis. The supinator longus, detached at its insertion, can be anastomosed with the extensor ossis metacarpi pollicis, and the extensor carpi radialis longior with the common digital extensors, or the extensor longus pollicis if the finger extensors are spared. In these (partial) cases it will be seen that whereas tendon anastomosis materially improves the hand within a few weeks, it does not, *a priori*, prejudice the further performance of posterior interosseus suture, were one skilful or rash enough to attempt it later.

The complete drop-wrist from musculo-spiral paralysis presents a more serious and difficult problem. This disabling condition is primarily due to the hyperflexion at the wrist-joint. Even in the normal hand the flexor muscles of the fingers cannot exert their maximum power unless there is a moderate degree of extension at the wrist-joint. When the normal hand is voluntarily placed in the attitude adopted by the drop-wrist of musculo-spiral paralysis, it is found that the fingers cannot be clenched—indeed, they cannot be fully flexed on the palm. Much more marked, then, is the condition in the flaccid paralysis of drop-wrist. In other words, the drop-wrist (or normal hand in this position) cannot make a "fist."

It behoves us, therefore, first to correct this hyperflexion of the wrist-joint and allow full play to the digital flexors. A patient with a wrist ankylosed in line with the forearm, with free finger movements, is in a better condition to make the fullest use of his fingers than one with a drop-wrist. I do not think this point is sufficiently appreciated by English surgeons. Continental surgeons have, however, recognized this, and carried out fixation of the wrist-joint by means of an artificial ligament manufactured out of fascia lata. The drop-wrist has not been cured—that is, return of power of extension—but a permanently slightly extended wrist has resulted, allowing full play to the flexors of fingers and thumb, which later can be abducted by its shorter abductor.

In these cases a fascial graft can be manufactured to act as a ligament permanently extending the wrist. It is attached to the third or fourth metacarpal distally and to the radius or ulna proximally, slightly nearer the upper extremity than the lower. If this is done, the flexor carpi radialis can be utilized for anastomosis with the extensor communis digitorum by passing its tendon through an aperture in the interosseus membrane, or, better still, the pronator quadratus, as there is less likelihood of the formation of adhesions with the tendon passing through muscle than interosseus membrane. The palmaris longus can easily be anastomosed with the extensor ossis metacarpi pollicis, and a very fair amount of abduction of the thumb results.

The method of transplanting the flexor carpi radialis and ulnaris to the insertions of the extensor carpi radialis longior and extensor carpi ulnaris works well, but leaves extensors of the fingers and those of the thumb without suitable "tendon donors," unless the final carpal flexor palmaris longus (not always present) is utilized, or a suture between the insertion of the pronator radii teres and the belly of the extensor communis digitorum effected, a not very satisfactory plan on *a priori* grounds as the resulting muscle pulls round a "corner."

The supporters of this method claim that after suitable education the flexor carpi radialis and ulnaris function as extensors of wrist or fingers.

I am of opinion that it is not so, and that they never act as true extensors, in that they become independent of flexor impulses, and are dissociated from the rest of the muscles supplied by the median and ulnar nerves. True extension consists of a "pressor" impulse along the extensor tract, with inhibition of the flexor group of muscles. But, inasmuch as the flexor carpi radialis and ulnaris are powerful muscles and the flexors of the carpus have been diminished by their removal, they fix or slightly extend the carpus when a flexor impulse is transmitted via the median and ulnar nerves to the flexor musculature of the forearm, no matter what the ultimate insertion of its individual muscles may be (in this case the extensor surface of the metacarpus for two of them). It is this slight extension and fixation of the wrist which allows full play to the flexors of the fingers and thumb. Shortening of the tendons of the extensor carpi radialis longior and brevior acts in the same way, but is apt to stretch after a time, and is not so satisfactory as the fascial ligament mentioned above.

When the essential fact is fully realized that it is the acute flexion of the wrist which is so disabling in musculo-spiral paralysis, the cases of this complaint will rapidly become fewer. There should be no excuse for a drop-wrist to-day. If the wrist be fixed in slight extension by any of the above methods, the "uselessness" of the hand will largely disappear, even if nothing more is done to promote active extension of the fingers. The fingers, indeed, will largely settle this question for themselves, and will, to a certain extent, become "passively" extended by the "rebound" on the inhibition of their flexor impulse.

Notes on the Operation.

Asepsis.—This is of course essential; a tendon anastomosis which undergoes the present day epidemic of sepsis will not be a success.

Correct Alignment of the Anastomosed Tendons.—It is necessary to lay stress on the absolute necessity of the "donating" tendon being in a straight line from the belly of the muscle to the acquired insertion. If this is lost sight of a "lateral implantation" ensues, resulting in little or no benefit to the patient. Surgeons, for long too closely wedded to the anastomosis of two tendons through short incisions, have lost sight of the intervening "kink" round which the tendon works with a maximum of friction and a minimum of effectual pull. This is particularly noticeable in the transposing of flexor muscles to act as extensors by passing them through the interosseus membrane, or the anastomosis of the palmaris longus to the extensor ossis metacarpi pollicis. The general lie of the muscle and oblique opening through the interosseus membrane or pronator quadratus, with a straight line passing through the points formed by junction of belly and tendon, interosseus opening, and insertion into the bone, are the essentials to be aimed at. Muscles do not act well when pulling round an imperfect muscular or fascial corner.

Re-education of the Transplanted Muscles.—Flexor muscles probably never act as extensors when their insertion is transplanted with this end in view. They probably act by fixing the carpus or by overbalancing their recently divorced brother flexors, bringing about some degree of extension when a flexor impulse is initiated in the cerebral cortex, since they are more powerful muscles than the surviving flexors. In any case they allow the flexors of the fingers better play, and before this can be effected to the maximum of its power, considerable re-education is necessary. This takes the form of active and passive "purposive" movements, faradism to the bellies of the transplanted muscles, and ergographic contractions of the fingers, which by the "rebound" helps to stimulate the contraction of the newly acquired extensors.

Early Movements.—Slight passive extension of fingers and carpus should be begun quite early (after a week or ten days), but active movements should be delayed until six weeks after operation.

The method of anastomosis I have used has been that of buttonholing the receiving tendon, inserting the donating tendon through it, and carrying its distal end for some distance along the dorsal aspect of the receptor, and fastening with several fine linen thread sutures.

In conclusion it might be mentioned that the observations in this paper are based on some dozen cases. In almost every instance the patient voluntarily expressed

himself as well pleased with the result. I do not recall any case in which there was not at least a fair degree of improvement on the ante-operative paralytic condition.

NOTES ON CASES OF INFLUENZA IN A BASE HOSPITAL IN ITALY.

BY

J. O. WAKELIN BARRATT, M.D., D.Sc.,

CAPTAIN R.A.M.C.(T.),

PATHOLOGIST TO THE HOSPITAL.

DURING a period of ten weeks (September 20th, 1918, to November 30th, 1918) a severe epidemic of influenza occurred in the area served by the hospital. Among the cases admitted 43 deaths occurred; it is with these that the present paper deals.

In 15 of these 43 cases an examination of the sputum was made during the early stage of the illness. In 9 instances the sputum was so heavily contaminated from the mouth, being loaded with bacteria of mixed types, that it was not possible to pick out any which could be regarded as likely to have been present in the bronchial secretion. In 2 cases (13 per cent.) pneumococci were found, in 1 case (7 per cent.) streptococci, and in 4 cases (27 per cent.) influenza bacilli. By the term influenza bacilli is meant bacilli, sometimes free, sometimes enclosed within polymorphonuclear leucocytes (both forms frequently occurring together), having a length of 0.8μ to 1.5μ , and a breadth of 0.5μ to 1.0μ , not staining by Gram's method, and presenting some variation in length and breadth in different sputa and cultures, as also in the same stained preparation.

It is of interest to compare the above figures with those obtained in the routine examination of sputum from patients in the hospital during the period under observation. In all, 116 specimens of sputum were received; of these, 64 (55 per cent.) were too heavily contaminated in the mouth to be of value; 28 (24 per cent.) contained influenza bacilli, 2 containing also pneumococci and 1 streptococci; 10 (9 per cent.) contained *Diplococcus catarrhalis*, 7 (6 per cent.) pneumococci, 3 streptococci, 1 staphylococci, and 1 tubercle bacilli. In all cases the bacteria enumerated were present in great abundance. If the specimens which were heavily contaminated from the mouth are discarded, the above percentages are much higher: thus the specimens in which influenza bacilli were found form 57 per cent. in the first series and 54 per cent. in the second. The latter series represented all the cases in hospital; it was not confined to cases regarded as influenzal.

On *post-mortem* examination, pneumonia was present in every case except two (95 per cent.), affecting both lungs in 37 cases (86 per cent.), and complicated in 5 cases (11.5 per cent.) with pleural effusion, in 8 cases (18.5 per cent.) with empyema, and in 1 case with pneumothorax. Cloudy swelling of the kidneys, liver, and spleen, friability of the heart muscle, and enlargement of the mesenteric and mesocolic lymphatic glands were present in varying degree in different cases. Enlargement of the adrenals, with collection of a brownish liquid or of semi-liquid material between cortex and medulla, was met with. Slight enlargement of Peyer's patches, with or without enlargement of the solitary glands of the small intestine in the neighbourhood of the caecum, was also observed. In one case there was ulceration of the large intestine, and in two cases blood staining of the root of the aorta and of the pulmonary artery was marked.

Microscopical examination of the exudate from the affected lung showed influenza bacilli in 17 cases (38.5 per cent.), streptococci in 18 cases (42 per cent.), staphylococci in 7 cases (16 per cent.), pneumococci in 7 cases (16 per cent.), and *Diplococcus catarrhalis* in 2 cases (4.5 per cent.); in 7 cases (16 per cent.) no bacteria were found. Double infections were observed in 14 cases—namely, influenza bacilli + streptococci, 4 cases; influenza bacilli + pneumococci, 2 cases; influenza bacilli + *D. catarrhalis*, 2 cases; influenza bacilli + staphylococci, 1 case; and staphylococci + streptococci, 4 cases. In one case a triple infection of influenza + staphylococci + streptococci was observed.