hospital appeared immediately before a French medical board, which sent them on convalescence.

In no case did we observe any symptom of poisoning either by iodoform or bismuth.

RADICAL CURE OF FEMORAL HERNIA BY THE INGUINAL ROUTE.

BY

PERCIVAL P. COLE, M.B., Ch.B., F.R.C.S.ENG.,

SURGEON, SEAMEN'S HOSPITAL, GREENWICH; SURGEON, QUEEN MARY'S HOSPITAL FOR THE EAST END; ASSISTANT SURGEON, CANCER HOSPITAL.

DURING the period of the war 509 cases of hernia have come under my care for operative treatment at various hospitals. All types—inguinal, femoral, ventral, umbilical—have been dealt with, and the age of the patients has varied from 9 months to 72 years. Fifteen of these herniae were strangulated, and in the whole series no death has occurred.

Thirty-seven cases of femoral hernia were met with, and of these 10 were strangulated. The method of procedure by the inguinal route is not new. It is attributed to Lotheissen, and was made the subject of a communication by Mr. C. H. Fagge in 1911. I became acquainted with the operation by witnessing it in Paris in 1909, and since that time have consistently adopted it. Mr. Fagge in his paper said:

So many of the younger surgeons now use this method that it is unnecessary for me to give exact details of the process.

Since that date men from nearly every school in the kingdom and from the colonies have acted as my residents, and I can only call to mind one who had ever seen femoral hernia dealt with in this manner. It would appear that the operation is not well known, and reference to standard textbooks supports this view. In Choyce's *System of Surgery*, Cheyne and Burghard's *Manual of Surgical Treatment*, and Burghard's *System of Operative Surgery*, no mention of it is made. Warren mentions it somewhat casually, but gives no details as to its performance. The only two books that give it due prominence are Rowland's and Turner's *Operative Surgery* and Taylor's *Operative Surgery*. Burghard (*System of Operative Surgery*) emphatically maintains that, apart from efficient ligation of the sac, no other procedure is really necessary, but qualifies this by suggesting that some endeavour should be made to close the ring as an extra precaution. R. W. Murray also holds that ligation of the sac is all that is required to obviate recurrence.

In Keen's *Surgery*, however, it is noted that mere ligation is attended by 30 per cent. of recurrences. Particularly emphatic is the statement in Cheyne and Burghard's *Manual of Surgical Treatment*:

In femoral hernia, besides obliterating the neck of the sac, it is necessary to do something to reduce the size of the femoral canal, as otherwise recurrence takes place apparently invariably.

These views are supported by the fact that at one time at one hospital there were under my care three cases of recurrent femoral hernia operated on by London surgeons of repute.

Multitudinous methods are described for dealing with the femoral ring and canal—a sure indication that no particular method can be safely relied on to prevent recurrence. In my opinion, the method of approach from below is unsound. Exposure of the femoral ring from above gives better access to the essential structures concerned. Of the procedures based on these lines the Lotheissen operation is considered the best, both from an anatomical and practical standpoint.

I propose to give a brief description of the operation, as applied to ordinary strangulated cases, with comments on the difficulties encountered and the advantages accruing.

The incision is that employed in the case of inguinal hernia. It should be free and prolonged at its lower end. The external ring is defined together with Poupart's ligament, and the external oblique aponeurosis divided throughout the length of the inguinal canal (Fig. 1). The arching fibres of the conjoined tendon are clearly demonstrated, the cord or round ligament defined and dislocated from its bed. An incision is then made through the transversalis fascia where it forms the posterior wall of the inner part of the inguinal canal (Fig. 2).

The extraperitoneal fat is thus exposed and the deep epigastric vein lying well to the inner side of the artery is

sought for at the outer end of the incision. This vein marks the outer edge of the peritoneal diverticulum that leads to the femoral ring. Separation of it renders easy the definition of the outer aspect of the neck of the sac. Some curved blunt instrument or the finger can then be passed behind the neck of the sac and made to emerge in the wound on its inner side. The condition of the sac as to contents can then be easily ascertained. If it is clear that nothing is passing through the neck from the peritoneal cavity, an

Fig. 1.

Fig. 2.

Fig. 3.

Fig. 4.

Fig. 5.

Fig. 6.

endeavour is made by blunt dissection and gentle traction to deliver the sac from the thigh. In Fig. 3 this has been done, and the unopened sac is presenting in the wound. The sac is opened and ligated at or above the junction with the peritoneal cavity (Fig. 4). The iliac vessels are then pulled outwards and the femoral ring clearly defined. Three sutures are then passed through the conjoined tendon above, and below through the posterior thickened margin of the femoral ring—Cooper's ligament. The outermost of these sutures lies close to the vein and should be inserted first (Fig. 5). The passage of these sutures is quite easy if a small half-circle needle be used. When tied, these sutures bring the conjoined tendon in contact with the posterior boundary of the ring, and a new bed for the cord or round ligament is thus constituted. It will be noted that no attempt is made to close the femoral ring. All direct communication with the peritoneal cavity is abolished—the femoral ring opens into the musculature of the anterior abdominal wall. In Fig. 6 a probe has been passed through the femoral ring into the thigh, the better to demonstrate this altered relationship. The remaining structures are united as in an inguinal hernia.

It happens fairly frequently that, though empty, the sac cannot be delivered easily. In this case the neck is cut across, and the hole in the peritoneum thus resulting is sewn up in the usual way. Taylor (*Operative Surgery*) comments on this difficulty, and advises that the incision be prolonged down into the thigh in order to aid delivery above by exposure and manipulation of the sac from below. This is laborious and quite unnecessary. It was my

custom in such cases to leave the sac in the thigh, but this procedure was abandoned after a somewhat unusual experience. I had left in the thigh a thick sac, which could be readily detected. The patient had an attack of vomiting some time after. This was really attributable to bad food, but she was sent up as an urgent case with a diagnosis of strangulated femoral hernia. The sac can be very quickly removed by undercutting the lower lip of the incision, and since then I have always removed it in this way.

Should the sac not be empty, the peritoneal cavity is opened immediately without any attempt at delivery of the sac, and in any doubtful case this is the easiest and safest course to pursue. The nature and condition of the contents is apparent at once. The most frequent content in an unstrangulated case is omentum and is in my experience the only content that becomes adherent to the sac. No attempt is made to separate these adhesions, but

FIG. 7.

the omentum is ligatured and cut off in the peritoneal cavity. The neck of the sac is then cut across, the sac with its adherent omentum being removed at a later stage in the manner above described.

On one occasion the bladder was encountered not as a hernial content but forming the inner wall of the sac. From below it might well have been included in the ligation of the sac, from above the condition was obvious and readily dealt with.

It has been urged that the operation is complicated, and the only reference to it in Keen's *Surgery* dubs it as such and so dismisses it. That it can be made complicated there is no doubt. The operation is rather more difficult in the male owing to the presence of the cord, but in these circumstances I have completed the operation, without undue haste, in eighteen minutes.

Operation from above has even greater claims in strangulated than in simple cases. The general peritoneal cavity is opened as soon as the extraperitoneal fat has been exposed. The condition of the bowel at the site of and above the constriction can be immediately determined. An abnormal obturator artery can be seen and avoided—it has been encountered on three occasions. When the operation is undertaken from below and the gut is not viable, textbooks on operative surgery recommend, almost without exception, that resection should be performed through a second abdominal incision. By the inguinal route a secondary incision is unnecessary, sufficient exposure being easily obtained to permit of the performance of any procedure that may be deemed necessary. Invagination of a gangrenous patch, excision of a gangrenous area without division of the mesentery, and excision of bowel and mesentery with lateral anastomosis have been accomplished in comfort.

Opportunity has been afforded of noting the nature of the constricting agent. It has been observed that reduction by an upward pull is difficult before the constriction is relieved. In other words, the resistance of the constricting ring to distension is considerable. The anterior, posterior and internal boundaries of the femoral ring are dense resistant structures, but the same cannot be said of the slight fibrous partition of the femoral sheath that forms its other boundary, on the side of the femoral vein. The resistance to distension of any ring can only be that of its weakest segment, and it follows therefore that the rôle of the femoral ring as a constricting agent has been exaggerated.

This theoretical consideration was put to the test in the last strangulated case—a male patient aged 72. Gimbernat's ligament was carefully divided outside the sac to such an extent

that a finger could be introduced between these structures. The reduction of the sac contents—a Richter's hernia of small bowel—was in no degree facilitated. Not until the neck of the sac itself was divided could the bowel be drawn up.

In operating from below the constriction is usually relieved from inside the sac, thus dividing its neck and Gimbernat's ligament at the same time. No due assessment can be made in this way of the relative parts played by each. It is contended that the essential agent in strangulation is the thickened neck of the sac, and that Gimbernat's ligament is merely a reinforcing structure. Another striking feature is the frequent occurrence of a Richter's hernia in association with a hydrocele of a hernial sac. In many of the simple cases dealt with the hernia was irreducible, and in every such case a hydrocele was demonstrated. It was noted that the hydrocele did not involve the neck of the sac, but was terminated at a level distinctly below it, at a situation indeed corresponding to

FIG. 8.

that at which the sac is twisted upward and outward, as it emerges from the saphenous opening (Fig. 7). It would appear that the combined effect of axial rotation and friction determines fusion of the sac walls at this site, and brings about the condition shown in Fig. 8, a condition readily complicated by the occurrence, as depicted, of a Richter's hernia. Irreducible femoral hernia of this type may easily escape notice, but the necessity for operation is obvious, and should be undertaken as soon as the diagnosis is made.

Advantages of Inguinal Route.

The advantages of the inguinal route operation may be summed up as follows:

1. It provides a certain means of cure.
2. It permits direct view of the essential structures.
3. Abnormal conditions can be recognized and dealt with.
4. Resection can be undertaken through the original incision.
5. It is neither difficult nor complicated.

OBSERVATIONS ON THE OPERATIVE TREATMENT OF TUBERCULOSIS OF THE LARYNX.

By DR. THOMAS RÜEDI,

LARYNGOLOGIST, DAVOS, SWITZERLAND.

DURING a period of six years I have treated at Davos 575 cases of laryngeal tuberculosis. On these 575 cases 1,548 operations were performed. In 61 cases curetting only was performed, in 168 both curetting and canterization, and in the remaining 1,319 the electro-cautery alone was employed. In every instance the laryngeal tuberculosis accompanied pulmonary tuberculosis, it being usually a case of stationary pulmonary tuberculosis (of the third degree, according to Turban-Gerhardt's classification) and of tuberculous laryngitis brought by the patient with him to Davos. In 39 cases the pulmonary tuberculosis was clinically demonstrable by the lung specialist as unilateral only (in 25 cases on the right and in 14 on the left). Of the 25 right-sided patients it was found that in 5 cases only the right side of the larynx, and in 4 cases only the left was affected. Of the 14 cases whose pulmonary tuberculosis was left-sided it was found that in 2 cases only the right and in one case only the left side of the larynx was affected. This agrees with the observations of Besold and Gidionsen and others, and argues against homolaterality of pulmonary and laryngeal tuberculosis.