In my opinion, the best method is to make two or three incisions into the cervix, as Dührssen advises. This immediately produces full dilatation, and also allows a loss of blood, which is itself of benefit. The incisions should always be stitched up when the intrauterine manipulation is completed. Any necessary intrauterine manipulation may be at once proceeded with. When the cervix is fully dilated, if any operative interference be required version or forceps will usually answer the purpose; the former is especially indicated when speed is necessary—for example, when a fetal heart is heard.

Such methods I have never known to fail to produce satisfactory emptying of the uterus and cure of the toxæmic condition. One operation remains to be discussed—namely, Caesarean section. Möller, Paterson, and Humbert of Strassburg advocate the vaginal route, but the number of cases recorded by each is small—Möller reports only seven. McCann⁷ described a successful case, and reported a mortality of 50 per cent. from the literature. The indications he gives for performing the operation in toxæmic conditions are (1) frequent fits, (2) labour not yet commenced, (3) contracted cervix, (4) a moribund mother and a live fetus (a rare combination), (5) pelvic contraction. In 1897 Kettlitz collected 28 cases showing a mortality of 50 per cent.; in 1899 Hillmann collected 40 cases with a mortality of 51 per cent., and in 1903 Streicherein reported 26 cases with a mortality of 31 per cent. From these figures it will be seen that the case mortality is really huge; in any case the operation aims at saving the child, if necessary at the expense of the mother, and I strongly hold that the reverse should always be our aim. I have never seen Caesarean section performed for eclampsia, and can see no prospect of ever performing it in this connexion except for mechanical reasons.

I present these opinions regarding the treatment of toxæmias complicating pregnancy in no dogmatic spirit, but merely as the considered results of recent clinical observations.

REFERENCES.

- ¹ Ward: *Amer. Journ. of Obstet.*, March, 1910. ² Zinke: *Loc. cit.*, 1911. ³ Short and Bywaters: *BRITISH MEDICAL JOURNAL*, 1913, vol. i. ⁴ Sellheim: *Zent. f. Gynäk.*, 1911, No. 2, p. 54. ⁵ Williams: *Johns Hopkins Hospital Bulletin*, 1906; *Glas. Med. Journ.*, December, 1912. ⁶ Bumm: *Wien. med. Klin.*, October, 1909. ⁷ McCann: *Lancet*, 1910, vol. ii, p. 787.

MALARIA AND DISEASES OF THE EYE.

BY

J. KIRK, M.D. EDIN.,

OCULIST TO EDINBURGH NATIONAL SERVICE BOARDS; FORMERLY
VISITING OPHTHALMIC SURGEON, GOVERNMENT HOSPITALS,
PENANG, STRAITS SETTLEMENTS.

My excuse for writing this brief note is to attempt to clear up in the light of my own experience the exact position of malaria as an etiological factor in the production of eye disease. The question is of great practical importance at all times, but more so in these days when we have large military forces operating in malarious regions. My attention was specially drawn to this subject during the eight months (1916-17) when I acted as ophthalmic specialist to the troops in Malta. There I was brought into contact with a very large number of soldiers from the Salonica front infected with malaria. A great many of these presented eye symptoms. Interference with vision and night blindness were common complaints, and were generally regarded not only by the men, but also by their medical officers, as being directly due to repeated attacks of malaria. On examination, however, it was found that the symptoms in a very large proportion of the cases were explained by latent errors of refraction which previously had not given trouble, but now, under the strain of the unaccustomed conditions of sun glare, dust, and exposure, enhanced by the debilitating influences of the malarial poison, had become manifest. The night blindness was invariably associated with and accounted for by anaemia. I have found, however, that in tropical countries the hypermetrope seems to be specially liable to this complaint, and a good many of my cases in Malta were associated with this condition. Retinal haemorrhages were seen occasionally. In the examination of the fundus in many hundreds of cases in malarious countries I never found this symptom at all common. A true malarial retinal haemorrhage does occur in subtertian cases, either

as the result of capillary thrombosis following parasitic emboli or as a sequel to definite blood changes of a pernicious nature, coincident with great anaemia, but the haemorrhage in the majority of the cases at Malta was certainly not due to such causes. I have a suspicion that some cases were due to overdosage with quinine in susceptible individuals.

Some American ophthalmologists regard malaria as the cause of much eye trouble, especially in the Southern States, and the statement has been made on the authority of Poncet, who published his results as long ago as 1878, that eye complications occur in 10 per cent. of malarial cases. My experience, as the result of twenty years' practice in a country where a vast amount of malaria was seen, does not coincide with these authorities. Eye complications in malaria appeared to be rare, yet the disease is credited with causing conjunctivitis, keratitis, corneal ulceration, iritis, retino-choroiditis, optic neuritis, opacities of vitreous and lens, nerve paralysis, retinal haemorrhages, etc. It is easy to make such wide and inclusive statements and difficult to disprove them. I would like to point out, however, that all these symptoms occur as frequently in districts in the tropics where malarial infection is not common as in parts where malaria is rife. They appear to be caused by the intense sun glare, dust and heat, the presence of such complicating conditions as trachoma, gonorrhoea and syphilis, debility, bad hygiene, and such intercurrent diseases as dysentery—especially the bacillary variety—a common cause of corneal troubles and iritis. In hospital practice in Penang, where the bulk of the patients were either Tamils or Chinese, though both nationalities were equally exposed to and frequently the subject of severe malaria, it was among the former race that the largest amount of such climatic eye troubles was seen. For instance, the Kling is very subject to all the forms of keratitis and corneal ulceration, night blindness, xerosis-conjunctivitis, optic atrophy of a primary chronic nature, non-specific, and probably due to malnutrition and chronic anaemia. All this is understandable, for the Kling is often an individual of poor physique and low resisting power and will not feed himself properly. The Chinaman, on the other hand, is a person of much vitality and good physique, though he takes all the malaria that is going. The cause of eye trouble in the Kling is, in the great majority of cases, malnutrition, degenerations, and ensuing secondary infections.

What we have to regard as almost certainly due to the direct action of the malarial poison are the following:

1. What is commonly called keratitis dendritica is simply a sequel to a herpes of the cornea, and due to a direct action of the malarial poison on the ganglion trophic cells of the fifth nerve. An exactly similar condition occurs quite frequently in influenza, and is due probably to a similar cause. I saw several cases of this in Malta among soldiers infected with malaria. A practical point is that quinine will not help in this condition unless the cause is diagnosed at the very beginning of the attack and energetic quinine treatment started immediately. Once the disease has a firm hold on the cornea and degenerative changes have occurred, general treatment does not appear to help, yet it will most certainly do so in the beginning.

2. A similar poisoning of nerve elements by the direct action of malaria results in external muscle paralysis. The third, fourth, and sixth nerves may be affected, more commonly, in my experience, the sixth.

3. A malarial neuralgia affecting the branches of the ophthalmic division of the fifth is common and may produce reflex conjunctival inflammation, but a pure malarial conjunctivitis would be difficult to diagnose.

As regards the deeper structures, I have never diagnosed malarial iritis or cyclitis or choroiditis. Neuro-retinitis may arise as a complication in malignant and subtertian cases and may be succeeded by nerve atrophy and blindness; and I have also seen a retrobulbar neuritis which appeared to be of malarial origin. Haemorrhage from the retinal vessels occurs under the conditions I have already mentioned.

Probably the truth of the matter as to the action of the malarial fevers in producing eye disease is that the parasites or their products have no direct power of producing inflammatory results in the eye structures (and this is in accordance with what happens elsewhere in other organs), but they have the power of affecting the

nervous supply of the eye, probably by direct degenerative action on the central nerve elements, and as a result the following may ensue: First, atrophic changes in the cornea producing a typical keratitis. Secondly, degenerative changes in the tissues of the optic nerve producing a true malarial neuro-retinitis. Thirdly, paresis of the extrinsic muscles of the eye. Fourthly, affections of the sensory nerves of supply.

This, I think, represents a fairly accurate and reasonable view of what really occurs. All the other many affections of the eye structures which have been described as being malarial are not truly so, but are probably simply inter-current affections or complications.

A practical conclusion is that in military operations in tropical countries in all cases of chronic eye trouble latent errors of refraction should be looked for. Their correction will in many cases cure or alleviate greatly the condition even though the error may be a small one.

AN OPERATION FOR RECONSTRUCTION OF THE URETHRA IN CASES OF SEVERE OR IMPERMEABLE STRICTURE.

BY

JOHN GUTHRIE, M.B., CH.B. GLASG.,
CAPTAIN N.Z.M.C.

THE following cases illustrate a method of procedure for reconstruction of the urethra in severe or impermeable stricture which will prove, in my opinion, more reliable and easier to perform than any other yet commonly practised.

A patient, aged 43, admitted to Christchurch Hospital in March, 1917, under my care had complete retention of urine, and the doctor who sent him had failed to pass any instrument. It was found that no instrument could be passed owing to a tough stricture in the bulbous portion of the urethra. The site of this stricture corresponded with a scar in the perineum, where an abscess had been opened some years previously. He stated that he had frequently to pass a catheter himself, that numerous doctors had dilated his strictures at intervals during the past eight years, and that the process had been more frequent and difficult of late. He had a high temperature and rapid pulse on admission, and looked pale and ill. Rigors also occurred.

Paracentesis suprapubically failed to relieve him except for a trickle. It was judged that the failure was due to the bladder being full of blood clot. Suprapubic cystotomy was performed without loss of time, and a very large quantity of foul-smelling blood clot and urine escaped. The bladder was washed out and drained. He recovered rapidly, and in a few weeks had put on flesh, but no instrument could be passed through the urethra. I decided to attempt some method of restoration of the urethra.

In Murphy's *Clinics* (December, 1916, vol. v, No. 6, p. 1189) Dr. Hochrein of Chicago described an original operation, performed for the first time in October, 1914, for the cure of an impassable stricture of the penile portion of the urethra by the use of a pedicled attached flap of dartos. I decided to adopt this method with necessary modifications.

Operation.

With the patient in the lithotomy position, a Wheelhouse's sound was passed as far as the stricture and an incision of ample length made in the middle line of the perineum down to and through the urethra on to the end of the instrument. With some difficulty a fine probe was passed through the stricture into the bladder. This was left in whilst the dense fibrous tissue was divided, laying open the "floor" or ventral wall of the urethra. I continued this incision until I was able easily to pass my finger into the bladder alongside the probe. I now dissected away all scar tissue, being careful only to preserve a strip of the "roof" of the urethra, representing, perhaps, one-third of the circumference. This appeared quite normal and was essential for the further steps of the operation. A broad flap of dartos was now cut in such a way as to leave the base attached in the vicinity of the point of the urethra which was first opened—that is, the most distal portion of the stricture. This flap would be $2\frac{1}{2}$ in. long and $1\frac{1}{2}$ in. broad, its rounded free end pointing towards the meatus. When dissected back, it reached easily to the limit of the dissection, which had been carried towards the bladder, as previously described. A rectal tube, the size of the middle finger, was now passed

through the meatus and brought out in the wound. This was re-entered and passed easily along the proximal urethra into the bladder. The flap was brought back over this, and, by slightly displacing the rubber tube, the left side of the flap was secured by numerous plain catgut interrupted stitches to the corresponding edge of the strip of urethral roof which had been left as described. This was repeated on the other side, and the end of the flap was finally stitched accurately to the divided circumferential edge of the proximal urethra. The gap in the dartos was closed with a few catgut stitches and the incision closed. The tube was secured to the retracted foreskin by a stitch behind the corona. It was my intention to leave the tube in for three weeks, but the patient sneezed it out at the end of a week. It was cleaned and sterilized, and I was somewhat surprised to find that it went in again with ease. Three weeks after the operation it was removed, and the patient has passed urine with perfect ease since (eleven months ago). I was able to pass a 14 to 17 Lister's bougie with ease six months after the operation. About seven months after the operation the patient had a typical attack of renal colic, and about a week later he informed me he had passed a round smooth brownish stone as large as a pea.

I have since performed similar operations in two other cases.

In the first of these I had the misfortune, in shaping the flap, to divide the superficial perineal artery on the right side. As a natural result the flap sloughed on that side. I managed to close the resulting fistula at a second operation, and when last heard of the wound was quite healed, and he was passing urine normally. In the second of these cases there was a perineal fistula just proximal to a stricture. The patient had for a considerable time been passing all his urine through the fistula, and his life was a misery to him. In this case I took the flap from the left side and turned it across. This seemed easier at the time, and involved much less time in stitching. The operation in this second case was performed under spinal anaesthesia, owing to the patient's age and debilitated condition. All the patients up to the time of writing are quite healed, and passing urine normally.

I have been induced to put these cases on record because I believe the flap operation is easier to perform and a great deal more satisfactory than the grafting operation. There is no doubt in my mind that the urethral epithelium rapidly grows from the strip left in the roof of the urethra, and covers the entire surface of the dartos flap.

THE INFLUENZA EPIDEMIC IN A CAMP.

BY

BREVET COLONEL C. AVERILL, R.A.M.C.(T.),

MAJOR G. YOUNG, and CAPTAIN J. GRIFFITHS,
R.A.M.C.(T.), R.A.M.C.(T.).

THE widely prevalent influenza epidemic visited this camp recently, 1,439 cases being admitted to hospital between June 21st and July 10th. The admissions on the first day of this period numbered 27, and gradually increased to their maximum on July 1st, on which day 154 cases were admitted. Thereafter the number of admissions gradually decreased.

Most of the cases were of characteristic type—admitted with temperature ranging from 101° to 104° , complaining of headache, backache (lumbar), general aching, and sore throat. The majority stated that they had been suddenly taken ill, with a feeling of great weakness. They looked ill; but apart from the fever, some congestion of the pharynx, and in a few cases slight bronchial catarrh, presented no other symptoms.

The illness ran a short course, the average duration of stay in hospital being five days. Within twenty-four to forty-eight hours of admission the temperature dropped by crisis to normal, or hovered between 99° and 100° for a few days. During the period of elevation of temperature the patients slept most of the time, and did not care to be disturbed, even for food. As soon as the temperature dropped they sat up in bed looking and feeling well, and began taking their food well.

Complications were few. Several patients had obstinate constipation; one or two had mild diarrhoea. Cough in a few cases was troublesome. Sixteen of the 1,439 cases