A Clinical Lecture ON ISCHAEMIC MYOSITIS.

BY
SIR JAMES PURVES STEWART, K.C.M.G., C.B.,
SENIOR PHYSICIAN TO THE WESTMINSTER HOSPITAL.

(With Special Plate.)

CERTAIN injuries which in pre-war days were uncommon have become relatively much more frequent since the present war. Amongst them is the condition known as ischaemic myositis. Before referring to individual cases let me remind you of the chief points in its morbid anatomy and causation.

MORBID ANATOMY.

Ischaemic myositis, as its name implies, is an affection of a muscle, or part of a muscle, resulting from local limitation of its blood supply. It is not due to total cutting off of blood, which would result in gangrene, but the muscle fibres affected, suddenly deprived of an adequate circulation, undergo coagulation-necrosis. There is an early or premonitory stage, during which oedematous swelling occurs in and around the affected muscles. Unless the blood supply is promptly restored by removal of obstruction to the arterial flow, coagulation-necrosis supervenes, which is irreparable. Later, the oedematous effusion becomes absorbed, and the coagulated muscle fibres become replaced by young fibrous tissue which rapidly contracts. Thus the affected muscles undergo shortening, become firm and doughy to the feel, losing their normal elastic consistence, until at last they are hard and board-like. Unlike most other forms of muscular atrophy, the bulk of the muscle is only slightly, if at all, diminished. Together with the muscles, the adjacent nerve trunks may have their blood supply reduced, and fibrous tissue may become deposited around them too.

CAUSATION.

A. In peace the commonest cause of ischaemic myositis is the too tight application of bandages or splints over a fractured forearm or elbow. This is the variety described in the textbooks as *von Volkmann's paralysis*. Those parts of the forearm muscles which are most exposed to pressure (and it is almost exclusively the upper limb which is affected) develop ischaemic myositis. Children or adolescents are the victims, partly, perhaps, because of the greater ease with which their arteries are compressed, but mainly, I suspect, because an adult usually insists on the loosening of uncomfortably tight bandages, whereas less attention is paid to complaints of tight constriction in a child.

B. In war the commonest cause is obstruction of the main artery of the limb by ligature. Thus, I have seen cases following ligation of the subclavian, the axillary, and the brachial artery, on account of primary wounds or of their complications, whilst in the lower limb ligation of the femoral artery sometimes produces ischaemic palsy.

C. Least frequent of all are the cases due to spontaneous obstruction of the main artery, from embolism or from thrombosis.

Let us study examples of each of these varieties.

A. Ischaemic Myositis from Tight Bandaging.

Here the fibrous sclerosis of the affected muscle is usually not a diffuse, but a patchy affair, occurring at the areas of maximal constriction by the splint or bandage. Above and below the sclerosed zone there may be an area of normal muscle fibres, so that the affected muscle comes to have a digastric or even a trigastric form.

Immediately following the application of the splints and bandages there is a premonitory stage of swelling of the hand, with coldness and cyanosis of the fingers. There is also acute pain in the limb, which is sometimes overlooked, being wrongly attributed to the original fracture for which the splints were applied. The oedema and cyanosis subside in a few days, but the fingers become drawn up. Then, after days or weeks, when the bandages and splints come to be removed, it is discovered that the fingers and wrist are stiffly flexed, not from adhesion in the joints or tendon sheaths, but from shrinking of the flexor muscles. Active movements of the hand are diminished or lost, and even passive extension of the

fingers and wrist is impossible. The degree of flexor contracture of the fingers and wrist varies in different cases, and may attain an extreme degree. Often the thumb escapes (having a long flexor of its own), whilst the other four digits (corresponding to the flexor sublimis and profundus) are severely affected. If we increase passively the flexion of the wrist, this slackens the flexor muscles, so that the fingers can now be slightly extended. On the other hand, attempted passive extension of the wrist exaggerates the deformity of the fingers, by pulling on the shortened flexors. The affected muscles feel denser than normal on palpation.

The electrical reactions of the affected muscles in this variety of ischaemic myositis may remain normal, both to faradism and to galvanism, since certain fibres, which have escaped coagulation, retain their normal excitability. Sensory changes in this variety are usually absent, but not always.

The following are illustrative examples:

CASE I.

A boy, aged 10, fell and sustained a fracture in the region of the right elbow. He was taken to a hospital, where splints were applied, reaching from the middle of the upper arm down to the hand. The splints were not removed until a month later, when it was found that he could not move his fingers properly. Assiduous massage was tried at a second hospital but failed to produce any improvement.

He came under my observation ten months after the original injury. At that time there was (Fig. 1: see special plate) a wide dimple or indentation on the flexor aspect of the forearm, apparently in the substance of the flexor muscles, and probably corresponding to a special lump of padding within the splint. There was no cutaneous anaesthesia or analgesia. The forearm muscles, which were not wasted, were contracted, so that all the digits except the thumb were semiflexed at the interphalangeal joints, and could not be extended, either actively or passively, unless the wrist were previously hyperflexed to its full extent, when slight extension of the interphalangeal joints could be carried out. Voluntary movements of the thumb were free in all directions. To faradism all the muscles of the forearm and upper arm reacted, but less briskly than in the healthy limb. To galvanism there was diminished reaction in all the muscles of the forearm, but KCC remained greater than ACC, as in health.

CASE II.

A boy, aged 6, was riding a donkey on the sands at the seaside, when he fell off. The donkey trod on his right arm, causing a fracture of the lower end of the humerus. The limb was bandaged in a flexed position, and subsequently fixed in a rectangular poroplastic splint. Next day it was noticed by the

parents that the child could not move the hand, which was swollen but not cyanosed. From the third day onwards the bandages were removed daily for purposes of massage and in order to dress a superficial abrasion on the front of the elbow, sustained at the time of the injury. Ten days after the accident the child began to complain of occasional pins-and-needles in all the digits except the thumb; this lasted for a fortnight, and then disappeared.

When examined, three and a half weeks after the injury, there was still considerable swelling above and below the elbow-joint. An x-ray photograph showed the fractured humerus in fairly good position.

The elbow could be moved, the biceps, triceps, and supinator longus all contracting voluntarily. All the muscles of the wrist, thumb, and fingers, however, were totally paralysed. The forearm felt dense and doughy. There was slight flexor contracture of the wrist, thumb, and fingers. When the wrist was passively extended the flexion of the thumb and fingers at once became increased. To faradism the upper arm muscles reacted normally, also the supinator longus, but there was no reaction in the flexor muscles of the wrist, thumb, or fingers, and only a trace of response in the extensor muscles. To galvanism there was no reaction to bearable currents in any muscle of the forearm or hand.

There was an area of total cutaneous anaesthesia to cotton-wool and pin-pricks in the "glove" distribution of the hand, front and back (Fig. 2). Joint-sense was lost in the thumb and fingers, normal at the wrist and elbow.

The radial pulse was palpable at the wrist, though for the first few days after the accident it could not be felt.

This case, of comparatively recent date, differs from the first mainly in the presence of anaesthesia of the hand.

FIG. 2.—Case II. Anaesthesia in ischaemic myositis of three and a half weeks' duration.

Such anaesthesia usually clears up in a few weeks or months.

CASE III.

This was an unmarried lady who came under observation at the age of 59 for epilepsy. At the age of 6 she had a fracture of the bones of the forearm. She said that the limb was for three days in tight splints, and that until they were taken off, the fingers were blue and swollen. Ever since, the right wrist had remained fixed in a flexed position, also the fingers, especially the ring finger. (See Fig. 3, in special plate.)

On examination, fifty-three years after the original injury, there was well-marked contracture of the right wrist, which was flexed almost to a right angle, and could not be extended, even passively (Fig. 3). The fingers were similarly contracted, chiefly at the terminal joints, but the thumb was freely movable. The flexor muscles of the forearm were wasted in their lower two-thirds. The surviving third of these muscles formed a distinct lump, above the upper limit of the original splint, contrasting with the atrophied part below. To faradism and galvanism all the muscles reacted briskly and normally. A radiogram of the forearm and elbow showed slight thickening in the region of the head of the radius, as if from an old fracture of its neck. There was no diminution of cutaneous sensibility.

B. Ischaemic Myositis following Ligation of an Artery.

In war wounds it is sometimes found necessary to tie large arteries in order to check primary or secondary haemorrhage, or to arrest the growth of a traumatic aneurysm. In some of such cases ischaemic myositis supervenes. This variety has several clinical differences from the group due to tight bandaging. Instead of having patches or zones of coagulation-necrosis, alternating with healthy zones, in the affected muscles, we find whole muscles undergoing diffuse coagulation-necrosis. The affected muscles come to have a curious hard, board-like consistence, much harder than in the other variety. Together with this, the electric excitability is often completely lost, both to faradism and to galvanism. Moreover, well-marked anaesthesia of the periphery of the limb is the rule rather than the exception, the sensory loss extending up along its outer border, but not in the territory of special nerve areas.

CASE IV.

A soldier, aged 33, had multiple wounds from fragments of high-explosive shell, including a penetrating wound of the right axilla, from a fragment which entered through the right upper arm. The fragment was located and removed, but three weeks later a deep axillary abscess developed, and at the end of the fourth week severe secondary haemorrhages occurred. The axillary artery was clamped and the subclavian artery was then ligatured. When he came under observation, in a hospital at Salonica, five months after the original shell wound, there was total paralysis of the right arm, forearm, and hand, all the muscles of the limb being hard and board-like. The only

FIG. 4.—Case IV. Anaesthesia in a case of ischaemic myositis following ligation of the subclavian artery.

voluntary movements were feeble contractions in the right pectoral, deltoid, and latissimus dorsi, but not enough to move the shoulder-joint. The deltoid and pectoralis major shared in this board-like hardness of the musculature of the limb. The trapezius and all the vertebro-scapular muscles were normal, both in consistence and in voluntary movement.

The elbow was semiflexed. It could be freely flexed and extended, passively, within a short range, but full flexion and extension were brought up sharp by the fibrous hardness of the muscles. All the joints of the hand were stiffly fixed in a lightly extended position, and could not be passively moved at all. There was comparative blunting to

cotton-wool, with total analgesia to pin-pricks in the whole hand, front and back, and along the radial side of the forearm and elbow (see Fig. 4). Joint-sense was lost in the fingers and wrist, normal in the elbow-joint. No pulsation could be detected in the brachial nor in the radial artery. There was slight oedema of the subcutaneous tissues of the flexor aspect of the forearm and outer side of the elbow.

To faradism no reaction was obtained in any muscle of the hand, forearm, or upper arm. A few fibres reacted in the deltoid and pectoralis major; the trapezius and all the vertebro-scapular muscles reacted briskly. No galvanic battery was obtainable to test the galvanic reactions.

CASE V.

A soldier, aged 28, was handling a loaded rifle when it went off. The entry of the bullet was through the left upper arm posteriorly, at the lower edge of the deltoid; the exit was just below the insertion of the pectoralis major anteriorly, in the line of the brachial artery. Next day the brachial artery was ligatured, and the wound was still granulating when he came under observation in Malta four weeks later. The patient complained, ever since the injury, of constant aching pain in his left fingers and palm, extending up to the elbow, also of difficulty in moving the fingers and of deficient sensation in the hand.

On examination, both hands were equally warm. In the left upper limb no pulsation could be felt in the brachial or radial artery. The left elbow was stiffly semiflexed and could not be passively extended. The left triceps muscle was hard and board-like in consistence, and had no voluntary power of contraction. The other upper-arm muscles, including the biceps, were normal. The flexor muscles of the forearm were hard, fibrous and paralysed, but not obviously wasted. The extensor muscles of the wrist, thumb, and fingers were normal in consistence, and contracted voluntarily.

The thenar and hypothenar muscles were paralysed, but not the interossei, for the fingers could feebly be spread out in fan fashion.

To faradism, there was loss of reaction in the biceps, triceps, flexors of the wrist, thumb and fingers, and in the intrinsic hand-muscles, whilst the supinator longus and the extensors of the wrist and fingers reacted briskly. The biceps-jerk and triceps-jerk were absent on the affected side.

There was total anaesthesia and analgesia of the hand, front and back, with a strip extending along the front of the forearm on its outer aspect (Fig. 5). Joint-sense was absent at the fingers, normal at the wrist and elbow. Vibration-sense was lost in the bones of the hand.

In this case the possibility of the anaesthesia being due to a concomitant lesion of the nerve trunks, whether by the original bullet injury or later, at the time of the operation, had to be excluded. But the fact that the supinator longus and all the extensors of the wrists and digits retained their motor power was against a direct lesion of the musculospiral as the cause of anaesthesia of the dorsum of the hand. Moreover, the absence of muscular atrophy of the flexor muscles and their characteristic hard consistence were against a lesion of the median and ulnar nerve trunks. Finally, the distribution of the anaesthetic area closely resembled that of other cases in our experience, in which the nerve trunks had certainly escaped direct injury.

CASE VI.

A soldier, aged 22, was shot through the left upper arm. The entry wound was at the middle of the left biceps; the exit posteriorly, at the same level, an inch behind the line of the vessels and nerves. He was admitted to hospital in England, ten days later, with a localized swelling in the middle of the upper arm, with absence of radial and ulnar pulses, and considerable

weakness of the left wrist and fingers. On the fifteenth day profuse spontaneous haemorrhage occurred from the exit wound. Under an anaesthetic the bleeding cavity was laid open, blood-clot was turned out, and the vessels above and below the aneurysm were ligatured. Next day the extensor muscles of the wrist and fingers were distinctly stronger. The wound healed uneventfully. At the end of five weeks it was noticed that there was total anaesthesia of the fingers, and partial anaesthesia of the lower two-thirds of the forearm.

When examined, two months after the original wound, there was anaesthesia to cotton-wool touches of the left hand,

FIG. 5.—Case V. Anaesthesia in a case of ischaemic myositis following ligation of the brachial artery.

FIG. 6.—Case VI. Anaesthesia in a case of ischaemic myositis following ligation of the brachial artery.

extending higher in front than behind (Fig. 6). To pin-pricks there was analgesia from the wrist downwards, front and back.

All the muscles of the upper arm, forearm and hand contracted voluntarily, except the long flexors of the wrist, thumb and fingers, which were hard and doughy in consistence, totally paralysed, and had no reaction either to faradic or galvanic stimulation. The left hand was colder than the right. Slight blisters formed occasionally on the dorsal aspect of the terminal phalanges.

G. Ischaemic Myositis due to Spontaneous Obstruction of a Main Artery.

Rarest of all are the cases of ischaemic myositis following spontaneous thrombosis of a main arterial trunk. In this variety there is no direct compression of the affected muscle, nor is the artery directly injured either by trauma or by ligation. It undergoes spontaneous thrombosis as a result of extension of some inflammatory process, usually in the peri-arterial tissues. Similar arterial thrombosis may also follow an embolus, especially a septic embolus from the left side of the heart, but such patients rarely survive to show ischaemic myositis.

The following is an illustrative example of spontaneous infective thrombosis of the external iliac artery, following an intrapelvic septic process.

CASE VII.

A married woman, aged 30, who had a large intrapelvic growth, was operated on at Westminster Hospital by my colleague, Mr. William Turner. The operation, which was carried out extraperitoneally, was one of extreme difficulty. To gain access to the growth one incision in the left iliac region and another in the left ischio-rectal fossa were required. Ultimately the tumour, a fibroma the size of a cocoa-nut, was successfully detached from its point of fixation to the left iliac spine, and from the vagina, bladder, and urethra, to all of which it was firmly adherent. The wound became infected, and widespread pelvic cellulitis supervened, but under assiduous and skilful treatment it slowly healed up.

Two and a half weeks after the operation pain and oedematous swelling were observed in the right foot and leg—that is, on the opposite side from the operation wounds. Next day there was pain and subjective numbness of both legs. Small

superficial trophic blisters appeared on both heels. The swelling of the right leg subsided in about a fortnight, but for some time there were painful cramps in the legs, more marked on the right side.

Three months after the operation, on getting up, it was noticed that the right foot was specially weak. At this stage she came under my observation, when the following state of affairs was found to be present:

There was marked feebleness of the right toes and ankle, but no individual movement was impossible. The anterior tibial and calf muscles of the right leg were hard and board-like on palpation. There was slight contracture of the calf muscles, precluding the right heel from reaching the ground when standing or walking. Passive dorsiflexion of the right ankle was impossible. There was no muscular wasting of the right leg; on the contrary, the right calf measured $\frac{1}{2}$ in. more than the left. To faradism and galvanism there was loss of reaction in the anterior tibial muscles and the short extensors of the toes, and marked diminution to faradism in the peronei and calf muscles, but without polar changes to galvanism. The knee-jerks were brisk and equal. The right ankle-jerk was absent, the left was brisk. Both feet were warm, and the posterior tibial arteries were palpable at the ankles.

To cotton-wool touches and pin-pricks there was a "stocking" area of anaesthesia up to a level $2\frac{1}{2}$ in. below the tubercle of the right tibia (Fig. 7). Joint-sense was lost in the right toes and ankle; normal at the knee and hip.

This case is specially interesting, inasmuch as, although the site of the original arteritis was presumably in the external iliac artery within the pelvis, nevertheless only a restricted group of muscles below the knee underwent coagulation-necrosis; moreover, the posterior tibial artery remained patent. It is possible that this was due to collateral anastomosis having restored the circulation in that artery, whilst the sural and anterior tibial branches of the internal popliteal artery remained obstructed.

FIG. 7.—Case VII. Anaesthesia in a case of ischaemic myositis following spontaneous infective thrombosis of the external iliac artery.

In the foregoing cases of ischaemic myositis following obstruction of a main arterial trunk, it is interesting to observe that in all of them anaesthesia of the limb was also present. The explanation of this anaesthesia is not yet clearly established. It may be that the blood supply to the nerve trunks is interrupted together with that to the affected muscles. It is not due to mechanical compression of nerve trunks, for in nearly all the cases anything in the way of direct pressure—for example, by ligature—could be confidently excluded; moreover, the distribution of the anaesthesia is not that of individual peripheral nerve lesions. It is more like the distribution of an ordinary peripheral neuritis, where there is no picking-out of individual nerve trunks, but where all the peripheral nerve trunks share in the general malnutrition.

TREATMENT.

Various ingenious operations have been devised by orthopaedic surgeons to remedy the deformity of that variety of ischaemic myositis which results from tight bandaging and splinting, and which is usually limited to the flexors of the wrist and fingers.

In view of the mechanical shortening of the sclerosed muscles, artificial lengthening of the flexor tendons at the wrist has been tried. The disadvantages of this operation are the extensive dissection which is required and the prolonged after-treatment of the wound.

A simpler operation is one which shortens the bones of the forearm, by excising an inch, or more, of both radius and ulna, so as to allow of the fingers being placed in a position of extension and even slightly hyper-extended. The patient then has a shorter forearm but a more useful hand. The drawback to this procedure is the tendency to the formation of still more scar tissue by operating on an already sclerosed limb.

It is therefore better to attack this class of case, as recommended by Sir Robert Jones, by means of a careful and systematic stretching of the shortened tissues. His procedure is as follows:

First, the wrist is passively flexed by an assistant, thereby allowing the fingers to extend. In this position small metal finger-splints are fixed to each digit by adhesive strapping, leaving the pulps of the finger-tips exposed. Over these digital splints, either at the first sitting or a day or two later, is fixed a flat palmar splint, so as to keep the phalanges and metacarpals in line. Next, at intervals of two or three days, the wrist is gradually extended, a few degrees at a time, being fixed in its turn to a series of suitably angled splints, until at last the wrist and hand are in a position of extreme dorsiflexion. The wasted finger pulps begin to fill up; later, the power of voluntary dorsiflexion gradually reappears. Assiduous massage to the forearm muscles, hot douches, and whirlpool baths will all aid in the softening and stretching of the affected muscles. It is also advantageous to supplement these measures by local hypodermic injections of thiosinamine. By such means a considerable degree of recovery can generally be obtained, even in severe and long-standing cases, although the best results are only to be looked for in the milder and more recent case.

The diffuse hard interstitial fibrosis following arterial obstruction produces less deformity than the first variety, and is therefore less likely to become the subject of the enthusiastic operating surgeon. Here we have to make the most of such muscle fibres as survive, by means of balneological measures, massage, and passive movements. But if a muscle or group of muscles have been practically reduced to fibrous tissue, it is obvious that very little can be expected in the way of recovery of voluntary motor power.

At a recent meeting of the Waffenbrüderliche Vereinigungen Deutschlands und Österreich-Ungarns a lively discussion on tuberculosis took place. Professor Sorgo of Baden thought that typhoid inoculation stirred up latent tuberculosis and that the tuberculous were peculiarly sensitive to foreign proteins other than tuberculin; soldiers should therefore, he argued, not be sent on active service until fourteen days after typhoid inoculation, and those already known to be tuberculous should not be inoculated against typhoid fever. All these views were ridiculed by Professor Kirchner of Berlin. To prevent soldiers from being sent on active service within fourteen days of typhoid inoculation was impracticable, as the need for men was far too pressing.

SIR JAMES PURVES STEWART: ISCHAEMIC MYOSITIS.
(See p. 151.)

FIG. 1.—Ischaemic myositis of 10 months' duration, showing contracture of flexor muscles of wrist and fingers, with wasting in their lower two-thirds.

FIG. 3.—Ischaemic myositis of 53 years' duration, showing inability to extend wrist or fingers, also wasting of muscles in lower two-thirds of forearm.

W. A. CHAPPLE: A MODIFIED STOKES-GRITTI AMPUTATION.
(See p. 158.)

FIG. 1.—A Stokes-Gritti in which the quadriceps had been cut and the edges stitched with catgut only.

FIG. 2.—A Stokes-Gritti in which the quadriceps had not been cut.

LIONEL J. PICTON: ATTEMPT TO BREED TUBERCLE-
IMMUNE CATTLE.
(See p. 157.)

FIG. 1.

FIG. 2.