The patient gradually became more and more toxæmic, and finally died eight weeks after admission.

Post-mortem Examination.

The heart was not enlarged, nor was there any degree of myocardial degeneration. A large cauliflower-like vegetation was found on one cusp of the aortic valve, almost completely closing the orifice of the valve. There was a similar vegetation on the adjoining cusp; the third cusp was free. The summit of the large vegetation was soft and friable, but the base was cartilaginous. The mitral valve had two or three similar but smaller vegetations on it. The right side of the heart was normal.

The spleen was dark, about twice its usual size, and showed two recent infarcts. There were also small infarcts in each kidney, but the rest of the examination showed nothing abnormal.

In a smear made from the aortic vegetation were seen enormous numbers of organisms resembling streptococci, rather large in size, and growing in chains of three to five.

Drawing of interior of left side of heart.

The remarkable features about this case were the insidious and slow progress, the absence of any definite cardiac symptoms, and the fact that the heart never became enlarged, in spite of the severity of the aortic lesion and the long duration of the case. This may be attributed to the fact that the large vegetation prevented any real leaking of this valve.

The patient was never really "cardiac"; there was no dyspnoea or oedema, and the pulse maintained a steady, regular, slow beat till the end. He was in a "typhoid" state throughout the disease, and died eventually of a slow toxæmia.

A CASE OF INTESTINAL OBSTRUCTION.

BY

J. M. WOODBURN MORISON, M.B., C.M.,

RADIOGRAPHER,
AND

L. WHITE, M.R.C.S., L.R.C.P.,

CLINICAL PATHOLOGIST,
THE DISTRICT INFIRMARY, ASHTON-UNDER-LYNE.

THE case here reported was evidently an instance of intestinal obstruction due to the healing process of old tuberculous ulcers of the bowel, possibly secondary to a lung infection.

John S., aged 68 years, was admitted to the Ashton-under-Lyne Infirmary on April 16th, 1918, having been sent in by Dr. Macintyre on account of abdominal pain and distension accompanied by vomiting. At 16 years of age he had "slow fever," and at 25 years had "inflammation of the bowels." He stated that with the exception of these illnesses he had always been a healthy man, and up to three months ago had never had any digestive troubles. The bowels had always acted regularly without aperients.

Three months before admission he began to be troubled with "pain in the stomach," the pain being chiefly located in the epigastrium. Usually it came on suddenly within half an hour after taking food, and ceased suddenly. He vomited occasionally and this relieved the pain; the vomit was "very acid." His appetite had fallen off, and he had lost flesh considerably. The family history was unimportant.

The abdomen was greatly distended and the percussion note tympanitic all over. The liver dullness was normal. On palpation strong peristaltic waves could be set up, causing the intestines to stand out in well marked relief in transverse bands running obliquely downwards from left to right in the umbilical and hypogastric regions, the so-called "ladder phenomenon." There was also marked gurgling on palpation of the descending colon. No tumour was palpable. Though tenderness was general, it was most marked over the epigastrium. The heart and lungs were normal; temperature 98°, pulse 98, respirations 24.

Blood Count.

Erythrocytes	...	...	...	4,664,000
Haemoglobin value	...	...	...	65 per cent.
Colour index	...	...	...	0.7
Leucocytes	...	...	...	12,000
Eosinophils	...	...	...	0 per cent.
Polymorphs	...	...	...	74
Lymphocytes	...	...	...	18
Large mononuclears	...	...	...	2
Transitionals	...	...	...	6
Mast cells	...	...	...	0

Total non-protein nitrogen, 95 mg. per 100 c.cm.

The urine contained a trace of albumin and a few granular casts; indican was present in marked excess.

X-ray Report.

The x-ray examination showed the stomach to be of a partial hour-glass form. The bismuth meal passed freely into and through the duodenum and jejunum, but was obstructed in the ileum, from which it passed slowly into the caecum. The radiogram shows the small intestines filled with the bismuth meal, portions of the intestines being ballooned out with gas. The x-ray diagnosis was "obstruction in the lower part of the ileum."

The bowels acted naturally once or twice daily. The vomiting persisted and the pain increased, so that it became necessary to administer morphine. On the evening of April 15th, his general condition being satisfactory enough to warrant it, it was decided to open the abdomen in order to relieve the obstruction. The diagnosis before operation was obstruction in the ileum or at the caecum, probably due to a malignant growth.

Operation.

The abdomen was opened by a right paracentral incision through the rectus. Numerous loops of distended small intestine immediately protruded through the wound. A rapid examination only was permissible on account of the general condition. No definite tumour or band or other cause of obstruction was found. A Paul's tube was inserted high up in the ileum, drainage being free through the tube.

The day following the operation the patient's general condition improved and the vomiting ceased, but from then onwards he became gradually weaker and died on the third day.

Post-mortem Examination.

This was performed twenty-four hours later. Emaciation was not marked. There were very few adhesions at the site of the wound and little shutting off of the abdominal cavity. There was no definite tumour palpable, but at intervals along the small intestines at the attachment of the mesentery could be felt small hard masses. The largest mass, however, was situated in the descending colon just above the sigmoid, and this would probably have been palpable during life but for the general distension. On removal of the small intestine it was found that at a point 77 in. above the caecum there was extreme stenosis, the lumen of the bowel being contracted to a size sufficient only to admit a wooden match. The stenosis was due evidently to scar tissue formation consequent on a previous ulceration of the bowel, the scarring spreading pincer-wise, with much thickening at the attachment of the mesentery. The bowel above the obstruction was ballooned and its wall markedly hypertrophied. There were no stercoral ulcers—"distension ulcers of Kocher"—present. The peritoneal surface of the intestine in the neighbourhood of the stenotic areas showed numerous yellowish plaques. Between this main obstruction and the caecum there were three similar lesions of varying degrees of stenosis, and in one instance evidence of an old ulcer was distinctly visible on the interior of the bowel by an oval depressed scar with its main axis transverse. A similar condition also existed at a point in the descending colon, just above the sigmoid flexure, but here the thickening and scar formation was much more pronounced. The peritoneal surface of the bowel showed patches of pigmentation. It was not possible to remove the stomach in the usual way as its posterior wall was firmly adherent to the subjacent structures, and after removal by dissection it was found that there was an old healed ulcer which had given rise to much thickening and contraction of the posterior wall and a mild grade of hour-glass stomach. The mesenteric glands were enlarged and one or two showed calcification changes. The

lungs showed at both apices old healed scars. The left base had firm, tough adhesions. The other organs showed nothing abnormal. Histologically the appearance of the lung sections were those of old healed tuberculosis. Sections of the bowel made at the site of stenosis showed thickening of the muscular layers and particularly of the submucosa, with proliferation of fibroblasts. The mucous membrane was in an atrophic condition. Sections from different points stained for tubercle bacilli were all negative.

The points of interest in this case are the length of time lapsing between the primary attack, forty years previously, and the shortness of the period—three months—during which the patient showed symptoms and was extremely ill. This is assuming that the history of "inflammation of the bowels" at the age of 25 was the primary attack of intestinal tuberculosis, and the details he gave of the illness give weight to this assumption, which is further supported by his claim to have enjoyed good health without a break up to the time when his final illness began. Excluding strictures of the rectum, obstruction of the bowel due to cicatricial contraction is a rare condition. A series of 70 cases has been collected (Bernay), and over a period of thirteen years the percentage of cases of cicatricial stenosis causing obstruction was 1.8 out of 669 cases at the London Hospital. According to Nothnagel, the only common causes are tuberculous and stercoral ulcers, but typhoid, dysenteric, duodenal and syphilitic ulceration may be followed by cicatricial stenosis. Tuberculous ulcers usually give rise to multiple stenoses, and in one case described by Nothnagel there were seven stenotic cicatrices.

PYLORIC STENOSIS WITH ACCOMPANYING SPASMODIC DYSPHAGIA.

BY

CAPTAIN E. B. BARTON, M.B., Ch.B., R.A.M.C.,

AND

LIEUT.-COLONEL H. H. C. DENT, M.B., F.R.C.S.,
L.R.C.P., R.A.M.C.(T.F.).

THE following case of pyloric stenosis presented considerable interest on account of an accompanying difficulty and pain in deglutition.

J. W. S., aged 36, previously in excellent health, developed gastric symptoms in August, 1916. He was treated in a military hospital from September 11th to November 10th, 1916, for dilatation of the stomach. At that time he had increasing dysphagia. In December, 1916, on two occasions attempts were made to pass oesophageal bougies. Finally he was informed that he had an oesophageal stricture.

Up to July, 1918, he was in a low category. He had lived on bovril, milk, and slops only, as he was unable to swallow anything solid. He was then admitted to the Clifton Military Hospital in a very debilitated condition. He complained of marked dysphagia, gastric distension and vomiting. The dysphagia had been constantly present since 1916, but was apparently not progressive. There was no regurgitation of food from the oesophagus. Regularly every two or three days he vomited a large quantity of fluid and mucus, sometimes streaked with blood.

A bismuth meal was given and the passage of the bismuth observed with the screen and the x rays. There was a distinct check in the forward peristaltic contraction of the phreno-cardiac portion of the oesophagus. The examination was repeated and the check appeared at a different level. There was very marked gastric delay. Bismuth still remained in the stomach after thirty-six hours. The opinion was formed that the condition was one of duodenal ulcer with pyloric stenosis and a reflex spasmodic stricture of the oesophagus.

On September 4th laparotomy was performed by a para-central incision. The stomach was found very much dilated. The cardiac end was normal. The pylorus was obstructed by a thick fibrous ring which was almost impermeable. Over this ring was a puckered scar suggesting an old duodenal ulcer. Posterior gastro-jejunostomy was performed, an opening being made 2½ in. in length in the stomach and in the jejunum.

He stood the operation well and had hardly any post-operative sickness. He was soon on solid food, and, although the first monthful causes slight difficulty in

swallowing, yet he can eat a full meal with ease. He was up in three weeks. His weight was 8 st. 3 lb. before operation; now (October 10th) it is 8 st. 13 lb. He looks and feels an entirely different man. X-ray examination shows normal swallowing.

INEQUALITY OF THE PUPILS.

BY

T. STEWART BARRIE, M.B., F.R.F.P.S.G.,
GLASGOW.

THE question of the meaning of inequality of the pupils is always being raised for diagnostic purposes, especially in affections of the thorax. According to many authorities, "inequality of the pupils is always pathological";¹ but many observers are not so dogmatic, and admit that the condition occurs in from 1 to 10 per cent. of all cases.²

In order to gain some idea of the frequency of this condition I examined, for this peculiarity, in the course of my work as an ophthalmic surgeon in the Glasgow recruiting area, the eyes of 326 men who had been sent for special examination. Inequality of the pupils was found in 35 cases, a proportion of 10.73 per cent. In none of these cases was there any sign of ocular disease nor manifestation of disease of the central nervous system. The pupils reacted readily to light, and the consensual and convergence reflexes were normal; in each case the inequality persisted in the contracted state.

The inequality in every case was indisputable, the difference between the diameters of the two pupils being from one to two millimetres. This difference is not large, but as pupils are compared by areas it was easier to note differences than to measure them. For example, pupils having diameters of 3 and 2 mm. respectively have areas in the proportion of 9 and 4.

An analysis of these 35 cases gave the following results:

1. The left pupil was the larger in 21 cases, the right in 14.

2. The visual acuity without glasses was the same in 10 cases; the right eye had the higher visual acuity in 8 cases, the right pupil being the larger in 4; the left eye had the higher visual acuity in 17 cases, the left pupil being the larger in 13.

3. The refractive condition of the 35 cases was as follows:

Emmetropia	...	...	...	2
Hypermetropia	...	...	...	2
Hypermetropic astigmatism	...	...	...	2
Myopia	...	...	...	11
Compound myopic astigmatism	...	...	...	1
Mixed astigmatism	...	...	...	3
Anisometropia	...	...	...	9
Not recorded	...	...	...	5

The conclusions which may be drawn from this investigation are as follows:

1. Inequality of the pupils is frequent.
2. It is associated with all refractive conditions, with a tendency to be more frequent in myopic conditions.
3. The visual acuity is not affected adversely by the fact that one pupil is slightly larger than the other.
4. The left pupil is more frequently larger than the right.
5. Inequality of the pupils occurs as a physiological condition.

REFERENCES.

¹ Fuchs, *Textbook of Ophthalmology*, fourth edition. ² Hawthorne, *Clinical Medicine*.

THE Italian Minister of Finance has set up a committee consisting of the deputies Sanarelli, Maggiorino Ferraris, Rava, Landucci, and Belotti, Professors Flora, Vinaj, and others, with the Under Secretary of State as chairman, to investigate and report on the watering places and thermal springs of Italy, and the means of developing their usefulness and increasing their value.

WITH the object of preventing the evils due to the excessively high prices of medicines in Italy, powers have been given to prefects, on the representation of the provincial health council, to fix a maximum price for drugs in most common use. Contravention of this ordinance is punishable by fine up to £40; for a repetition of the offence the penalty is imprisonment up to the term of one year. The shop of the offending pharmacist may be closed temporarily, or in case of repetition permanently, by decree of the prefect.