the attention of those concerned is attracted by the appearance of a rash on the body. The wound is very inflamed, even sloughing, and round the wound the skin is bright red and swollen. The swelling of the skin becomes less the further it is from the wound. The rash on the skin is bright red and confluent in some cases, and where it is not confluent it is a punctate erythema. There is a red and congested throat and a red and swollen tongue. In three or four days peeling begins on the neck and rapidly spreads to the body and limbs. It is prolonged, often over six weeks. It commences from above downwards and finishes in the same order. This class of case closely resembles the first class, but is more severe. Obviously, a greater and more virulent amount of infective matter has found an entrance through the cut surface of the operation wound.

The third class of case is where there is a wound, or lesion of the skin, but there is no local erythema. In this class of case there are the early symptoms—the sore and dirty throat with exudation, the shivering, fever, and malaise, but the rash and tongue are not such prominent features. The rash may be confluent on the back and elbows, but is a discrete and punctate erythema on the body and limbs. It is often evanescent, lasting only a day or two, and in times of stress may be missed, and attention be subsequently drawn to the condition by the discovery of desquamation. In this class of case desquamation follows in the same order as in the first two, but may be later in commencing. Cases occur with little rash where desquamation starts as late as the eleventh or fourteenth day, misses the body, and is found only on the hands and feet.

Operation in this class of case did not produce any recurrence of the symptoms.

All these cases seem to belong to the group of streptococcal infections. They vary in their intensity with the facilities for the admission of the coccus or its toxin into the circulation. I cannot distinguish them from ordinary scarlet fever, and am of opinion that the diseases are identical. These diseases have been treated in the same ward with ordinary scarlet fever cases, and there have been no cases of secondary infections. It is impossible to say that it is safe to keep them in the same ward with general surgical cases, whether clean or infected, but where missed cases have been found late in the disease, it does not appear that they have left a track of infection behind them.

The period of observation has been eighteen months, and the number of cases sixty, and the wounds in most parts of the body.

INCUBATION PERIOD OF INFLUENZA.

BY

PETER MACDONALD, AND J. C. LYTH, M.B.,
M.D., B.S.,
YORK.

THE incubation period of influenza is not yet definitely settled, and although the number of the cases we describe below is small, the opportunities for their observation were so close that we give an account of them, as they go some way towards fixing it within small limits of error.

We travelled from London together on Thursday, October 3rd, by train leaving King's Cross at 5.30 p.m., arriving at York 9.30, and as we were leaving the carriage a young flying officer who had come the whole way with us, and was coughing and sneezing at intervals, informed us that he was ill, and had had "influenza" for several days. On Saturday, October 5th, we both became ill, and both developed typical attacks of influenza. With both of us the illness developed suddenly with laryngitis; in both the first signs were a severe attack of coughing; and in both the time was noted fairly accurately as being between 2 and 2.30 p.m. One case was quite mild, the temperature was never over 101°. The other was more severe; the temperature rose to 104.5°, and the catarrh extended to the bronchi. His wife and two children also developed influenza, and in their cases the first symptoms showed suddenly about 2 p.m. on Monday, October 7th.

Now we are convinced that we became infected from our travelling companion during our train journey, more likely towards the end of the journey; and if we take the time of infection as 9.30 this fixes the incubation period

for both of us at a minimum of forty-one hours, with a maximum margin of error of four hours. But as we may presume that the bacterial content of the railway carriage increased with the time of our journey, we incline to regard our infection as dating from the end of our journey rather than the beginning; so that we take the incubation period to have been nearer forty-one than forty-five hours.

The three cases developing in the family of one of us point to a similar incubation period, as their illness started almost exactly forty-eight hours after his, and as it is likely that infection would not take place until a few hours after the first symptoms, the incubation period in these three cases must have been very nearly the same as in our own two.

It can readily be understood that we were in no condition to conduct extensive bacteriological examinations, but a culture taken from the posterior nares of one of us on October 10th, with a guarded swab to prevent soiling from the flora of the mouth, showed colonies of Pfeiffer's bacillus and of *Micrococcus catarrhalis*.

Reports of Societies.

DISCUSSION ON ENCEPHALITIS LETHARGICA.

At a meeting of the Sections of Medicine, Epidemiology, and Pathology of the Royal Society of Medicine held on October 22nd, Surgeon Rear Admiral Sir HUMPHRY ROLLESTON, K.C.B., R.N., in the chair, a discussion took place on encephalitis lethargica.

Lieut.-Colonel F. W. MOTT gave a summary of the results of the examination of two brains, by M. Marinesco, Professor of Neurology at Bucharest, who had been working for some time past in the pathological laboratory of the Maudsley Neurological Clearing Hospital. Disseminated miliary or punctiform haemorrhages visible to the naked eye existed in the grey matter in the neighbourhood of the floor of the fourth ventricle, the aqueduct of Sylvius, and even the third ventricle, and were also found in the posterior part of the pons and the peduncles. The cerebral cortex, except for congestion of some of the vessels of the leptomeninges, had shown in these two cases neither macroscopic nor microscopic lesions. On the contrary, the first segment of the spinal cord, which was the portion of the cord available, had presented the same histological lesions as the pons, bulb, and peduncles. Microscopical study of the above mentioned regions had demonstrated the existence of four kinds of lesions:

1. Infiltration of the walls of the small vessels and especially the veins, consisting of lymphocytes and plasma cells in the adventitia, disposed in several layers. The endothelium and fibroblasts might also take part in the inflammatory process.

2. Foci of interstitial inflammation consisting of neuroglia cells of several kinds, including large cells with voluminous eccentric nucleus, and many fibrillar prolongations, lymphocytes, and polynuclears. The foci of interstitial inflammation appeared sometimes to be altogether independent of vascular infiltrations, and might occur in the roots of the nerves—for example, hypoglossal, pneumogastric, etc.

3. Lesions of the nerve cells which did not correspond with those usually seen in infantile paralysis. There was dissolution of the "soi-disant" Nissl bodies, relative achromatosis, reduction in volume of the cellular body and of the number of prolongations and multiplications of the satellite cells, but only exceptionally was there evidence of neuronophagia, as was described by Economo.

4. The foci of haemorrhage, the most obvious macroscopic change, were seen to be much more numerous when microscopic examination was made. These haemorrhagic foci remained circumscribed around the walls of the small vessels, and red corpuscles were mingled with the cells of inflammatory infiltration, or they constituted a kind of covering, and floated about around the vessels. In spite of the very great number of haemorrhages the vessel wall did not appear to be necrosed, but sometimes a solution of continuity of the vessel wall could be seen.

From microscopic investigation of these two cases M. Marinesco drew some conclusions upon the nature of the disease. He regards lethargic encephalitis as a disease entirely different from botulism, from the haemorrhagic poliomyelitis of Wernicke, from the poliomyelitis of Heine-Medin, and from sleeping sickness. Like these two last diseases, it was an infectious inflammatory disease, but the nature of the infectious germ had not yet been determined. It was distinguished from botulism by its