fibrous tissue occupying the gap between them are now carefully cut away. Finally, each fragment is bevelled by cutting away a flake of bone from its outer surface with bone forceps. In this way raw bone is exposed at the ends of the gap and on the outer aspect of the fragments for about one inch from its extremity. All bleeding is then carefully arrested and the bone graft prepared.

After experimenting with bone from the ribs, the tibia, and from the jaw itself, the iliac crest was finally selected as the site from which to take the graft. The bone is tough and can be cut with bone forceps without splitting. Further, a graft can easily be obtained of any length or breadth and the natural slightly curved contour of the crest is approximately that of the jaw. The graft should be taken preferably from the same side as the operation wound, thus allowing the patient to lie comfortably on the opposite side.

An incision is made over the crest commencing at the anterior superior spine and extending as far back as required. The muscles are then separated on either side of the crest and pressed back by retractors. The bone is cut by an ordinary Horsley's hand saw. The graft should be two inches longer than the gap to be filled. If a more curved piece of bone is required, as to fill a gap near the chin, the graft is made to include the bone between the superior and inferior spines.

In this way a graft four inches long has been obtained which filled a gap extending from the angle of the

FIG. 2.

mandible on one side to well beyond the chin on the other, and the curve was so accurate that no subsequent modelling was needed.

After removal of the bone the muscles detached from it are sewn together with catgut and the wounds closed. No inconvenience whatever seems to result. The ends of the graft are now bevelled with bone forceps, the bevelled areas lying on the prepared outer surfaces of the jaw fragments. In this way the graft overlaps the gap at each end for an inch. Two advantages result from this: (1) A broad line of bony contact between the graft and the fragments is provided with increased prospect of speedy firm osseous union, and (2) there is practically no risk of separation in the event of the gap being increased by subsequent manipulations during the application of dental splints, as a certain amount of sliding can take place without contact being lost.

No attempt is made to fix graft or fragments by plates and screws, by wiring, or even by dovetailing the graft into the fragments. All these measures have been tried and discarded. The presence of foreign bodies greatly militates against successful healing, a sinus down to the plate or wire almost invariably forming. Attempts also to make the graft act as a splint by dovetailing it between the fragments have not led to satisfactory results.

Our practice now is to keep the graft in place by sewing the soft tissues closely over the graft and the ends of the fragments by hardened catgut. This has the additional advantage of closely surrounding the graft with living vascular tissue and abolishing dead spaces in which blood clot and serum can collect. This improves the nutrition of the graft and diminishes the risk of sepsis.

Finally, the skin is approximated with a few interrupted stitches. No drainage is employed beyond that of leaving spaces between the skin sutures to allow of the escape of serum. A simple dressing and bandage is applied and the patient sent back to bed.

FIG. 3.

No attempt is made to reintroduce dental fixation splints until the wound is firmly healed and the compound fracture has been converted into a simple one. This usually occurs in two weeks, after which the case is treated as one of simple fracture of the jaw. Firm osseous union occurs in from two to four months, but it is inadvisable to fit the final dentures until at least four months have elapsed, and it is perhaps wiser to allow an interval of six months.

We intend to supply notes of cases in a subsequent paper. We desire to record our appreciation of the very valuable assistance at the operations that we have received from Captain Learmonth and Sister Dorothy Jones of the 1st Southern General Hospital.

INFLUENZA: ITS CAUSE AND PREVENTION.

BY

W. FORD ROBERTSON, M.D.,

PATHOLOGIST TO THE SCOTTISH ASYLUMS.

MANY who have read the recent writings on this subject in the *BRITISH MEDICAL JOURNAL* must feel there is great need that those who have had special opportunities for the study of the *Bacillus influenzae* and its pathogenic action should make public any knowledge they have that may be useful at the present time. I therefore hope that the following notes may be acceptable.

The best method of growing the bacillus of influenza seems to be still unknown, though it was described and demonstrated before a scientific society in 1916 and mentioned in a paper published shortly afterwards.¹ It has been constantly used since 1914 in the laboratory of the Scottish asylums, where it was originated by the senior assistant, Mr. John J. Ritchie. It takes advantage of the fact that the bacillus of influenza is stimulated to growth by the mere proximity of certain other micro-organisms, especially those of the *catarrhalis* group and pneumococci. In this method, which may be referred to as the alternate drill method, generally three straight drills of, say, *Micrococcus catarrhalis* are made from bottom to top of a suitable agar surface, one at each side and the third exactly in the middle. Between these drills two drills are made from colonies of *Bacillus influenzae*. Abundant growth, suitable for the preparation of vaccines, can generally be scraped off after twenty-four hours' incubation, but it is of advantage to continue incubation for two or three days.

The best medium is haemoglobin agar, which, though it has been used in the laboratory of the Scottish asylums for many years and repeatedly described, seems still to be rarely used, at least in a correct way. I obtain haemoglobin serum from sheep's blood (secured with aseptic precautions), which has been allowed to clot and from which most of the serum has been decanted. The tubes are placed for an hour or so in a freezing mixture. When thawing takes place the remaining serum contains a strong solution of haemoglobin. About 1 c.cm. of this haemoglobin serum is added to each tube of nutrient agar in the fluid state, while at a temperature of about 60° C. The medium is allowed to set in a sloping position. It is important to use a higher percentage of agar than is generally prescribed for nutrient agar. The tubes should be incubated for twenty-four to forty-eight hours before they are used for the making of cultures, in order to test their sterility. In the preparation of vaccines the culture material obtained by the alternate drill method is scraped off by means of a platinum loop and placed in 1 per cent. carbolic acid in normal salt solution. Twelve to twenty-four hours at 37° C. suffice for sterilization. The preparation of the vaccine is completed by the gravimetric method. It is greatly to be deplored that bacteriologists should still adhere, with false conservatism, to the inaccurate, time-absorbing and inconvenient plan of standardizing vaccines in millions. In the gravimetric method the whole batch of dried vaccine is accurately weighed in the chemical balance and made up to a strength of 1 c.cm. = 1 mg.

Influenza bacillus vaccines for use should consist of a one-tenth dilution of the emulsion in the stock tube—namely, 1 c.cm. = 0.1 mg. The dose of influenza bacillus for therapeutic immunization ranges from 0.005 mg. to 0.1 mg.

In the course of the past five years, by the application of these methods, I have investigated considerably over a hundred cases of chronic infection by the bacillus of influenza. Chronic infections of this kind are very much more common than is generally supposed. The only point I wish to dwell upon in regard to these cases is that of the evidence they furnish regarding the pathogenic character of this bacillus, which apparently is still in doubt on the part of many who are able to give publicity to their views. In nearly all of these cases therapeutic immunization has been carried out. The focal and general reactions produced, and the successful therapeutic results that have been obtained, are consistent only with the view that the bacillus from which the vaccines were prepared was the chief cause of the catarrhal conditions from which the patients were suffering. Correct therapeutic doses commonly caused distinct focal reactions; overdoses were followed by the development of symptoms simulating those of influenza. A continuation of correct therapeutic doses over a period of from six to ten weeks almost uniformly resulted in complete recovery from the malady. If the bacillus of influenza had not been the chief cause of the ailment these results could not have been obtained.

In acute influenza, therapeutic immunization acts, in my experience, with equal success. A dose of 0.02 mg., repeated after twenty-four hours, generally causes the temperature to fall to normal within forty-eight hours. With ordinary vaccines there is apt to be a good deal of reactive disturbance from twelve to twenty-four hours after the first dose. For these and other reasons it is of advantage to use sensitized vaccines in acute cases. For the purposes of the preparation of these sensitized vaccines, two sheep are kept immunized against *Bacillus influenzae* at the laboratory of the Scottish asylums. The administration of a sensitized vaccine is followed by comparatively little reactive disturbance.

The most convincing evidence of the causal relationship of Pfeiffer's bacillus to influenza can be obtained from the phenomena that may be observed in the carrying out of protective inoculation. I made a number of these inoculations at the time of the summer epidemic; none of the cases contracted influenza. I have recently immunized a much larger number of persons, but the evidence of the protection afforded has, of course, still to be obtained. I give, at intervals of from two to three days, successive doses of about 0.03 mg., 0.06 mg., and 0.12 mg., or two doses of 0.04 mg. and 0.12 mg. The smaller doses are given first in order to create a certain degree of tolerance of the toxin before the full dose required for complete protection is injected, and to provide against the possibility of hypersensitiveness, which occurs in all persons who are suffering from a chronic infection by the bacillus. After each of these doses most persons within twelve to twenty-four hours experience in mild degree the symptoms of influenza. If the dose of 0.12 mg. is given at once, severe disturbances are generally experienced. Any one who likes can have a controlled attack of influenza. If any sceptic will submit to the test, I believe I can convince him, by means of an injection of a full dose of the killed culture, that the bacillus of Pfeiffer is the cause of influenza.

It is very important that it should be understood that under existing conditions more is required than protective inoculation against the bacillus of influenza. In Scotland, at least, there has been for two months or longer also an epidemic of coryza, dependent upon infection by a special type of *Micrococcus catarrhalis*. Acute infection by this micrococcus generally causes, in addition to "cold in the head," severe headache, aching throughout the body, and some rise of temperature, especially at night. Correct therapeutic immunization is capable of cutting short the attack in forty-eight hours. Beyond question, any one suffering from this form of coryza is very specially prone to be attacked by the influenza bacillus. Most cases of acute influenza that I have recently investigated have been infected also by this special type of *Micrococcus catarrhalis*. When I am asked to protect against the bacillus of influenza I explain the importance of this other form of infection that is prevalent, with the result that all of the cases are given the double inoculation. The dose of this micrococcus is the same as that of the bacillus of influenza. In ordinary course it would have been about four times larger, but experience has shown that this type

of *Micrococcus catarrhalis* produces much greater irritation at the place of injection than the ordinary strains, already notorious for their unpleasant local effect.

I regret that I feel obliged to criticize adversely the recommendations of the War Office Conference of October 14th, which were published in the BRITISH MEDICAL JOURNAL of October 26th. If the errors contained in these conclusions are not recognized now, I challenge their future comparison with the orthodox teaching as it will stand, say, two years hence. It is not properly recognized that the dosage for therapeutic immunization is essentially different from that required for protective inoculation. The doses recommended are, so far as my own experience enables me to judge, strangely wide of the mark. The 30 million dose of *Bacillus influenzae*, given as the initial one, works out at about 0.005 mg. This is a small therapeutic dose, and neither it nor the second dose of double the amount can be expected to afford any valuable degree of protection against infection. The introduction of pneumococcus and streptococcus protective inoculation is a mistake. These bacteria are not commonly spread from one person to another like the bacillus of influenza. It is quite different, however, in regard to *Micrococcus catarrhalis*, protection against which is omitted in these recommendations. The coryza caused by this organism is extremely infectious, and those who have acquired it are, beyond a doubt, specially prone to be attacked by *Bacillus influenzae*. Therefore it is hardly less important to protect people against this micrococcus than against the bacillus. The doses recommended are also open to criticism. The initial dose of 100 million pneumococci and the second of 200 million represent about 0.03 mg. and 0.06 mg. In my experience, if the patient happens to be suffering from a chronic infection by the corresponding type of pneumococcus, these doses will cause rather violent focal reactions. Regarding their value for protective inoculation I cannot speak. The doses of 40 million and 80 million streptococci (presumably *pyogenes* and *anginosus*) represent about 0.01 mg. and 0.02 mg. These, in my experience at least, are merely small therapeutic doses. Protective inoculation should be directed exclusively against attack from *Bacillus influenzae* and *Micrococcus catarrhalis*. It is a different matter when we have to carry out therapeutic immunization in a case of influenza. It is then of proved advantage to add to the suitable dose of *Bacillus influenzae* (0.01 to 0.04 mg.), small doses of a polyvalent pneumococcus vaccine (0.01 to 0.02 mg.) and of *Streptococcus pyogenes* (0.01 to 0.02 mg.), as well as of *Micrococcus catarrhalis* (0.03 to 0.06 mg.). If possible, the vaccines should be sensitized ones.

REFERENCE.

¹ Journ. of Mental Science, January, 1917.

BLOOD AGGLUTININS IN MENINGOCOCCAL ATTACKS.

By I. WALKER HALL, M.D.,

PROFESSOR OF PATHOLOGY, BRISTOL UNIVERSITY; CEREBRO-SPINAL FEVER BACTERIOLOGIST FOR SOMERSET AND GLOUCESTER.

(Report to the Medical Research Committee.)

A STUDY of the agglutinative action of human serum during an acute attack by meningococci involves the consideration of many factors.

If it be granted, for purposes of argument, that the available technique is beyond criticism, there is still a possibility that the reactive combinations evoked by the acting antigens of living, or lysed, cocci may differ from those which follow the injection of organisms killed by heat or other coagulative measures. The emulsions of cocci prepared for the quantitative estimation of agglutinins may also be placed in the category of variants. Owing to inconstancy of amino-acid nutrients or vitaminic stimuli, or a failure to produce artificially the conditions under which the meningococci proliferate during infection, the cultures may exhibit variations in vigour, of growth or extent of autolysis. It has already been remarked by Gordon, Ainley Walker, Hewitt, and Gibson, that the specificity of the agglutinins induced by the injection of killed meningococci is more or less dependent upon the species of animal used for the purpose.¹ That is to say,