as it was then called. In the first place, the patient's general health must, if necessary, be improved, for hysteria is liable to occur in those in whom it is below normal. Secondly, the treatment must be begun as early as possible, for, quite apart from its being always wise to get a patient well quickly, the longer hysteria lasts the more difficult it is to cure. Thirdly, as just mentioned, the doctor must most carefully avoid, by his mode of examination and by his wording of questions, suggesting symptoms. Visitors and nurses must also avoid suggestion. Nothing is more heartrending for the doctor than the well meaning visitor, who says: "Poor man! what a terrible time you must have had! The wonder is you have any power left in your legs." A sentence such as this will undo hours of hard work on the doctor's part, and it is because of this constant liability to the suggestion of new symptoms that it has been found advisable to treat hysterical soldiers in special hospitals where it is understood how to manage them, and where they are encouraged by seeing other patients getting well. This is an extremely influential part of the cure. Fourthly, and most essential, is the reawakening of the supraliminal consciousness and the making of suggestions powerful enough to overcome the hysterical symptoms. Sometimes this is not performed by the doctor. A soldier was invalided from the front for mutism; he had not spoken for weeks. He was playing draughts with another soldier, who cheated. The mute broke out in a torrent of abuse, and spoke ever after. A woman had lain in bed for years owing to hysterical paraplegia. The house caught fire; she ran downstairs fast enough to save her life. Hypnotism was employed for this part of the treatment, but is not at present much used as it is unnecessary. Persuasion and re-education are now largely practised, with excellent results. Men who have not been able to use one or other of their limbs or straighten their backs lose their disability in quite a short time. The medical journals, books on war neuroses, and the studies from the Seale-Hayne hospital give details of many cases. It is necessary that the doctor should convince the patient that a cure can and will be effected. Many employ electricity as a means of suggestion. We use at Guy's an effervescent mixture containing asafoetida and valerian; it is successful in many cases of mild hysteria; it acts by suggestion that it will cure, especially as the patient eructates into her mouth asafoetida and valerian during a great part of the day, and is therefore constantly reminded that she is going to be cured. The Weir Mitchell treatment, which often works wonders in the wasted hysterics, illustrates these principles. The general health is improved by the overfeeding and massage, the isolation removes the patient from the suggestions of her friends and her old surroundings, and a skilful doctor and nurse avoid suggesting new symptoms and devote their energy to suggesting to the patient that she will recover.

I trust the war has shown that we ought to give our best attention to hysteria, for the study of it is a pleasure in itself. It is very common; if you doubt this go to Lourdes. It is capable of cure; it is not, as some people seem to think, malingering. Remember Sir James Paget's famous saying: "The patient says she cannot, the friends say she will not, the doctor says she cannot will," and it is a curse to the patient, her friends, and her relations. Do you recollect what Fielding says in *Amelia*:

"These fatigues added to the uneasiness of her mind, overpowered her weak spirits, and threw her into one of the worst disorders that can possibly attend a woman—a disorder very common among the ladies, and our physicians have not agreed upon its name. Some call it a fever on the spirits, some a nervous fever, some the vapours, and some hysterics."

"Oh! say no more," cries Miss Matthews, "I pity you, I pity you from my soul. A man had better be plagued with all the curses of Egypt than with a vapourish wife."

GUERRERO, DOMINGO and ARGÜELLES have confirmed, in the *Philippine Journal of Science*, vol. xiii, the value of tartar emetic in the treatment of yaws in the manner described by Castellani in our columns on October 21st, 1916. The dose used was 1 grain. The number of doses varied from 15 to 18, and in those cases which recovered the medication was continued for five to twenty-seven days. Complete recovery took place in about two-thirds of the cases. In some symptoms diminished, but in others no improvement was observed. In five cases relapses occurred in from two to five months after the lesions had entirely healed.

The Montgomery Lectures

IN

OPHTHALMOLOGY.

DELIVERED IN TRINITY COLLEGE, DUBLIN, JUNE, 1919.

BY

GORDON HOLMES, C.M.G., C.B.E., M.D., F.R.C.P.,

PHYSICIAN, ROYAL LONDON OPHTHALMIC HOSPITAL; PHYSICIAN TO OUT-PATIENTS, NATIONAL HOSPITAL FOR THE PARALYSED AND EPILEPTIC, QUEEN SQUARE, LONDON.

LECTURE II.—DISTURBANCES OF VISUAL SPACE PERCEPTION.

WHEN my eyes fall on any visible object, as on this piece of chalk on the table in front of me, I perceive it and I can in addition form certain judgements on its spatial relations to myself and to other visible objects; and if it happens to be similar to something I have seen before I can recognize its nature. That is:

1. I can determine its position in space in relation to myself and consequently point to it if I can control and direct the movements of my arm.

2. I can also form an approximate estimate of its distance from me and therefore touch it if it be within my reach, or move immediately to it if it be further from me.

3. And if another object be within my field of vision, as this second piece of chalk, I can form an approximately correct judgement of the relative positions of the two, that is, whether the one is higher or lower, to the right or the left of, or nearer or further than the other.

4. When one of the objects casts its image on the periphery of my retinae I can move my eyes so that it comes directly into central vision, and, if it is within a short distance, converge and accommodate too so that its image is focussed accurately on my maculae.

5. I can estimate its size, both absolute and relative to that of other objects.

6. I can recognize that it has thickness and depth, or in other words that it is tridimensional.

7. Finally, I can identify it, that is, associate its visual perception with the memory images of similar or allied past impressions, and with other qualities that I have learnt by previous sense experience, as weight, brittleness, etc.

All these judgements that I can form about this piece of chalk, which has merely thrown an image on my retinae, may be disturbed, either individually or collectively, by cerebral lesions. To-day I intend to neglect disturbances of identification which are generally called visual agnosia, and to restrict discussion to affection of the spatial attributes that we associate with the visual perception of the object.

We must, however, consider briefly how we localize in space objects which we merely perceive by vision, and how we attain the spatial attributes with which we endow our visual perceptions. This has been the subject of much discussion. The two chief explanations are those that have been offered by the nativistic and by the empiristic hypotheses. The nativistic view attributes localization in space to innate properties of the sense organs or sensory centres, properties which are inherited and ready for use by the newly born animal. According to the empiristic school, on the other hand, all space perception is of intellectual origin and is acquired only by habit and experience, by the testing and controlling of visual sensations by tactile, muscular, and other sensations from all parts of the body. This empiristic view is of considerable interest here, as it was first put forward more than two hundred years ago by George Berkeley, then a Fellow of this College, in his celebrated "Essay towards a new theory of vision." A study of the disturbances and dissociations produced by disease may in this, as in many other subjects, contribute towards the solution of the problem, though it cannot by itself settle it.

During the late war I was able to observe eight men with gunshot injuries of the brain who had disturbances of spatial orientation. All presented more or less similar symptoms, and as most of them remained under observation for considerable periods it was possible to investigate them repeatedly. The detailed observations on seven of these cases have been published elsewhere.¹ These disturbances cannot be attributed to blindness or amblyopia,

for the acuity of central vision was good in all the patients, and though there were defects in the fields of vision in several these defects bore no relation to the symptoms, which were identical in other men in whom the fields were normal. The symptoms were also unconnected with paralysis and loss of sensation, and there was no serious degree of mental deterioration in any of them. One man only had aphasia of severe degree.

ABSOLUTE LOCALIZATION.

The most obvious symptom was the inability of the patients to determine the position in space, in relation to themselves, of objects which they saw distinctly. When I held up a knife in front of one man he said at once "that's a pocket-knife"; but though his eyes were directed on it he stretched out his arm in a totally wrong direction when he was told to take hold of it. Another man struck my face with his hand as he attempted to point to a pencil which I held two or three feet to my right side. Similarly when asked to pick up from a table objects which they could see and recognize, these men could never bring their hands directly to them, and often made very gross errors in their direction. One intelligent man described his experience when a meal was placed on a bed-table in front of him; when he tried to take a piece of bread he brought his hand under the table rather than above it, and on attempting to seize the cup he found his fingers first in the tea, and in his second attempt on a plate to one side of it.

This inability to recognize the positions of objects seen was obvious whether they were in central or peripheral vision, but in the latter case the errors were much greater; most of the patients, in fact, were eventually able to localize fairly well, though not with normal accuracy, objects at which they looked.

It was only when they relied on sight alone that they could not localize the positions of an object in space. If it touched any part of their own bodies they could always bring their hands to it immediately and correctly; touch gave them the necessary local knowledge that they failed to obtain from vision alone. This was well shown when one man was given a spoon and told to take soup from a bowl on the table in front of him; he succeeded in placing the spoon in the bowl only after repeated attempts, though he could always bring it accurately to his mouth, but when he was allowed to hold or touch the bowl with his left hand he fed himself without any difficulty. His left hand gave him the spatial information about the bowl that he could not obtain from sight alone. These patients also localized the direction of sounds as accurately as normal persons; when blindfolded they pointed in the direction of a noise, and walked towards anyone who called them, as readily and as accurately as a normal person can.

It was the estimation of distance that was most severely affected. When the patients attempted to touch any object near to them they frequently projected their hands far beyond it; or, if it were beyond their reach, they often tried to grasp it before their hands had reached it. One man, for instance, was asked to seize a piece of paper about six yards away from him; he at once extended his right arm towards it, and continued groping for it as he approached it, and even after his hand had passed it. They were also unable to make even approximately correct estimates of the distance of objects which they saw; a tall chimney about 500 yards distant seemed "perhaps twenty yards" from one man—in fact, he remarked: "Everything I see seems to be the same distance from me."

RELATIVE LOCALIZATION.

Not only were they unable to recognize the spatial relations to themselves of objects which they saw, but they could not determine accurately the relative positions of two objects that were at the same time in vision. When a silver and a copper coin, or two pieces of paper of the same size but of different colour, were held up in front of them they were often unable to describe their relative positions, and especially their relative distances from themselves. They either guessed indiscriminately that the one was nearer than the other, or confessed their inability to decide. One patient, for example, could not say which of two men, one dressed in khaki and the other in hospital blue, was the nearer when they were respectively seven and fourteen yards from him; and when he was shown a panoramic view, in which he recognized several promi-

nent objects, he confessed that he could form no conception how they were placed in respect to one another. But when he was allowed to touch or walk to two separate objects, he no longer entertained any doubt of their relative positions.

RECOGNITION OF LENGTH AND SIZE.

Another prominent symptom in all my patients was their inability to distinguish the absolute and relative lengths and sizes of objects they were shown. They could not estimate at all accurately the length in inches or feet of a line or rod, unless it were a familiar object that they recognized. They were always uncertain which of two pencils was the longer unless the difference was considerable, and they could not divide a line accurately; in many attempts one segment was made four or five times as long as the other, yet they were generally satisfied with their attempts. They failed, too, to find the centre of a circle accurately. I asked one patient to divide a large slice of bread equally between us, but the piece he kept for himself was at least six times as large as that which he gave me, and when I pointed out that the division was unfair he failed to see it until he touched the two pieces with his fingers.

IN WALKING.

The inability to localize objects in space by vision alone was very obvious when these patients were allowed to walk about. One man, when only a few yards from his bed, started to walk in a totally wrong direction when told to go to it, and another when asked to sit in a chair to one side of me walked up against me violently, and then explained he had no idea I was so near to him. Each was at first unable to find his way directly to any point to which he was asked to go though he saw it, or succeeded only after taking a devious route, but later most of them could go straight to a point when they kept their eyes fixed on it. In walking about they frequently collided with large obstacles, as they failed to realize their direction or their distance from themselves. Some of the men walked repeatedly up against walls, occasionally so violently as to hurt themselves, as they had no idea how near they were to them, though they saw them distinctly. One interesting feature was the difficulty they experienced in getting around obstacles which they encountered; on running into a chair, for instance, one man stood still for a moment, apparently perplexed as to how to avoid it, then shuffled to one side and when almost free back again, so that it was only after repeated attempts he succeeded in passing it. When, however, he was allowed to place his hands on it he felt his way round it at once, as easily as a normal person can in a dark room.

All these patients had great difficulty in learning their way about. One man who was in the same small ward for three months could not, even at the end of this time, go directly to his bed, though he could see and identify it, if he had to make two or three turns on his way. Another was tested day after day in an open space divided by a partition in which there was a gate, but even after weeks of training he could rarely go straight through the gate to a point in the other division which he knew. The visual impressions out of which they would normally construct their ideas of space were devoid of spatial qualities, and this made it impossible for them to obtain clear pictures of the topography of even small areas. The topographical memory of ways and places with which they were acquainted before they were injured was also severely involved, though the patients possessed good memory pictures of things that had been familiar to them. Those who were questioned on this point were unable to say how they would find their ways about in their old haunts. One man, who had lived in a small provincial town all his life, remembered that the railway station was near his home, but he could give no idea of how he would go to it from there, or to his workshop, which he knew was close by. Another man had a vivid memory of the trenches in a sector in which he had served for a considerable time, but he had no idea of the positions of the different trenches in relation to one another, or of how to go to any point in them. Though they could describe what they passed on the way, and even the general appearance of the streets, these patients retained no idea of the relations in space of the separate objects or streets to one another.

COUNTING AND READING.

The inability to recognize the position in regard to self of objects seen and the relative localization of several objects, as well as to retain a correct picture of the spatial relations which is normally acquired by sight, produced many other symptoms. My patients could rarely count correctly four or five similar coins or matches placed irregularly on a table in front of them. They generally turned their eyes from one to another and back again, counting and recounting till they became hopelessly confused, since they failed to recognize the mutual spatial relations of the coins and consequently had no means of distinguishing those they had already counted, as all were similar in appearance. When coins of different sizes and colour were employed they could distinguish those they had already included in their tally and consequently succeeded. And as touch gave the local information that was lacking in visual impressions they counted the coins quite readily when allowed to feel them with their hands.

The same disturbances accounted for their difficulty in reading. Though the patients could see the smallest type and recognize the written words they were often unable to read the words of a line in their proper order, and their eyes jumped irregularly from line to line, or even on to another paragraph or column. Their most constant difficulty was to find the beginning of the next line when they had finished reading the words of one. Writing was affected by the same factors. None of my patients, except one who was aphasic, had any trace of agraphia; they could write words and sentences correctly, but in the letters they attempted to write the individual words were placed so irregularly, and the lines ran so frequently across one another on the page, that the whole was scarcely intelligible.

RECOGNITION OF MOVEMENT.

The appreciation of movement was accurately investigated in only one of my patients. This man was never certain whether a moving object approached or receded from him when it moved in the line of his visual axes, but he detected the slightest lateral or vertical movement and recognized its direction at once. The smallest movement could be also detected by peripheral vision, but he was always uncertain of its direction unless its range was much larger than the normal person requires for its instant recognition.

VISUAL ATTENTION.

Another symptom which all these patients presented was a disturbance of visual attention. Every object in central vision seemed to absorb attention, so that images which fell on the periphery of the retinae were often unobserved; attention was incapable of noticing two or more visible objects at the same time. When the observer's hands were held up, one on each side of the patient's visual axes, he could instantly see the movement of either the right or the left, but not the simultaneous movement of both hands. When asked to look at a needle placed on the table he often failed to detect a pencil placed on one side of it, or if there were two pencils he could see only the one or the other at one time. This affection of visual attention was obvious in many other ways too. While sitting in the ward it was noticeable that the patients usually saw only what their eyes were directed on and they took little interest in what was happening around them, unless their attention was called to it by sounds or other sense impressions. In walking, too, they frequently passed without observing them things that would have excited the notice of normal persons.

This visual inattention was not due to blindness or other defects in the fields of vision, as no defects that could explain it were discovered after careful examination, and images were frequently perceived on portions of the retinae where similar images passed unnoticed immediately before or after. There was no regularity in perception. This is a condition that results not infrequently from lesions of the parietal lobes, and it may be a unilateral or a bilateral phenomenon. It is commonly found apart from disturbances of visual localization and is entirely independent of this. It plays a part, however, in some of the symptoms we have already dealt with, as if the visual impressions of objects do not attract attention the patient fails to notice landmarks by which he could later find his way, and by which he could construct a geographical picture

of his surroundings. In attempting to count objects, too, he is liable to miss out some owing to their images not exciting attention.

OCULAR MOVEMENTS.

There were also abnormalities of the eye movements in all my cases, though none of the ocular muscles were paralysed and the pupillary reflexes were preserved. The most prominent of these symptoms was a disturbance of fixation. When spoken to suddenly the patient first stared blankly in front of him, and then his eyes roamed round the room till they fell, as if by chance, on my face. Similarly it was only after some time and searching that he could bring his eyes directly to an object on one side of his visual axes at which he was asked to look. This difficulty in fixation was a natural consequence of defective spatial localization, for if the position of the object that threw its image on the periphery of the retinae could not be recognized the eyes could not be moved immediately to it. The fact that most of the patients could turn their eyes promptly to points that they could locate in space, as to parts of their own bodies that were touched or to objects held in their own hands, shows that in these cases this was the explanation of the symptom. But as in a few of my cases and in others that have been recorded the subjects were unable to deviate their eyes directly to points on their own bodies this symptom may be also due to disturbance of the central mechanism of the ocular movements, a disturbance on the effective side which is closely parallel to the perceptual affection on the sensory.

Accommodation was similarly affected; when an object at which they gazed was approached slowly towards their faces their eyes failed to converge, their pupils did not contract, and as accommodation did not occur the object became blurred or indistinct when nearer than eight inches or so. This was also mainly due to the spatial perceptual disturbances, but as some of the patients could not accommodate on their own fingers accurately an affection of the central mechanism that controls it must be also assumed.

The blinking reflex was also abolished. These patients never reacted by blinking or withdrawing their heads when anything was suddenly swung towards their faces, or to threatening gestures. In most of the cases this was probably secondary to the loss of spatial perception; some indeed explained that they did not recognize the nearness of the threatening object, but in others the cerebral centres which are concerned in this reflex were probably involved.

STEREOSCOPIC VISION.

Stereoscopic vision, or the power of recognizing thickness and depth in solid objects, was severely affected in only one of my patients. To this man everything appeared flat and bidimensional; he identified familiar things only by associating their areal shape with previously acquired experiences, just as the normal person interprets drawings and pictures as the objects they represent. He described a man who stood in front of him as flat, "I can only see the front of him, I do not notice that he is thick; I cannot tell the depth in anything." A box seemed to him a piece of flat cardboard, no matter at what angle he saw it, and when it was placed in his hands he was surprised to discover it was a box. And a glass tumbler looked like a bit of flat glass which changed in shape as it was moved about. When he was placed in front of a flight of steps he saw only a number of straight lines on the floor. Everything seen was flat, nothing had thickness or depth in it; stereoscopic vision was completely abolished and remained so till he passed from under our observation three months after the infliction of the wound.

POSITION OF THE ANATOMICAL LESIONS.

In all my eight cases the condition described above was associated with gunshot wounds of the head which involved the posterior and upper parts of both parietal lobes. *Post-mortem* examinations were obtained in two cases; in both the lateral surfaces were injured in the neighbourhood of the angular and supramarginal gyri, while the missiles passed through the mesial surfaces immediately dorsal or posterior to the splenium of the corpus callosum. Craniometric measurements in the remaining cases indicated that in them the same parts of the brain was damaged. As similar symptoms have been observed in cases in which there were vascular

lesions limited to the lateral surfaces of the hemispheres in the region of the angular gyri, we can probably attribute the disturbances of visual space perception to destruction of this portion of the parietal association area. It is interesting that certain observations made thirty years ago by Sir E. Sharpey-Schafer on monkeys from which he removed the angular gyri, can now be interpreted as the result of affection of space perception, similar to that with which my patients suffered.

NATURE OF THE SYMPTOMS.

In attempting to analyse these symptoms and interpret their nature and significance we must consider the physiological factors on which spatial localization by sight and tridimensional vision depend.

The simplest case is localization in a bidimensional plane represented by a hollow sphere, the centre of which would correspond to the pupil. When the eye is focussed on any point on this sphere the concave surface of the retina receives images of all objects that lie on the arc within its field of vision. Then we can recognize the relation in space of any one of these objects to that on which our eyes are directed, and the relation of any other two objects on this arc to one another. This we do by virtue of the local signs of the stimulated retinal points, just as by means of the cutaneous local signs we can localize any contact on the surface of our body.

But such local signs can serve us only in relative localization to some central point or to our visual axes; they cannot inform us of the absolute position in space of any object with reference to ourselves, as they take no account of the position of our eyes, of our heads, or even of our whole bodies. This additional information we must obtain from proprioceptive impressions from the ocular muscles, from our labyrinth, and from the muscles of our bodies; and when by virtue of these we can become aware of our position in space and the direction in which our eyes are looking, we can, by using the retinal local signs, determine the absolute position in space of the object we see in relation to ourselves. Even in this simple case the association of retinal and various proprioceptive afferents is necessary, and the loss of any of these factors may disturb this function. In the cases described here, the afferents from the labyrinths, and from the ocular and other muscles, were unimpaired, and the symptoms must be attributed either to a disturbance of the local sign functions of the retina, or to the failure of the brain to associate and integrate these with other afferent impressions. Many direct observations showed that the retinal local signs were affected.

We have to deal with a more complex condition in considering the perception of distance, as there are several other factors concerned in this. Binocular vision undoubtedly plays a part, and the proprioceptive impressions from the intrinsic and extrinsic ocular muscles that are excited by accommodation and convergence contribute to it when the object is near. The estimation of distance is also aided by the apparent size of known objects, by their distinctness and light intensity, and by the parallax that is obtained on movement of either our heads or of the objects. There is no evidence that any of these separate factors were disturbed in my cases. The inability to estimate distance must be ascribed to failure to correlate and associate the various afferent impressions on which perception of the third dimensions may be based, and to assimilate them with past experiences and thereby supply to consciousness the data on which a correct intellectual judgement can be made.

Loss of stereoscopic vision might be regarded as a natural consequence of inability to appreciate relative distances since, it might be argued, this would make it impossible to discriminate the depths of the different portions of a tridimensional figure. But stereoscopic vision was affected in only one of the eight cases in which the perception of distance was seriously disturbed. The physiological factors on which it depends have been the subject of much discussion; the appreciation of relative distance, difference in light and shade, the parallax and the sense of perspective, all undoubtedly play a part, but it is to the fusion of the non-corresponding images formed in the two eyes and their integration with these other sensory data into a single concept that we mainly owe the appreciation of depth in objects seen. There was no gross disturbance of any of these individual factors in my

cases except the appreciation of relative distance, which the intact faculty of stereoscopic vision in the other cases shows was not sufficient. An affection of the sense of perspective may have contributed, but as this faculty is of intellectual origin and acquired by training and experience, we are ultimately forced to regard loss of stereoscopic vision, too, as the result of a failure to combine and associate the different sensations on which the appropriate intellectual judgements can be made, rather than as due to the loss of any of the physiological factors on which it depends.

REFERENCE.

¹ Holmes: *Brit. Journ. of Ophthalmol.*, 1918, vol. ii, p. 449. Holmes and Horrax: *Archiv. of Neurol. and Psychiatr.*, 1919, vol. i, p. 385.

A CRITICISM OF CERTAIN RECENT CLAIMS TO HAVE DISCOVERED AND CULTIVATED THE FILTER-PASSING VIRUS OF TRENCH FEVER AND OF INFLUENZA.

BY

J. A. ARKWRIGHT, M.D.

(From the Lister Institute of Preventive Medicine.)

In recent papers by Bradford, Bashford and Wilson^{1 2 3 4 5 6}, it is claimed (1) that six diseases including trench fever and influenza, which have up to the last twelve months baffled investigators, have been shown conclusively to be caused by organisms which pass filters of such a quality that they hold back ordinary bacteria; (2) that in the case of seven other diseases (thirteen in all) the virus has been cultivated and can be recognized microscopically when stained by suitable methods.

Most of the work has been published only in the form of condensed epitomes, but rather fuller details on which these claims are based have been recorded for two diseases—namely, "epidemic polyneuritis" and influenza.

These papers have attracted considerable attention on account of the extent and importance of the results claimed.

OTHER RECENT ATTEMPTS TO CULTIVATE A FILTER-PASSING VIRUS OF TRENCH FEVER OR OF INFLUENZA.

A good deal of work in recent years has encouraged the view that the viruses of trench fever and influenza are of such a kind that their filtrability may be ultimately established, and has prepared the way for further researches on the cultivation of filtrable viruses.

1. The American Red Cross Committee on 'Trench Fever published evidence' that the virus of trench fever would in some instances pass through a Chamberland filter "L" with a pressure of 760 mm. of mercury.

2. The War Office Trench Fever Committee in London (1918-19) has made five inoculations of filtrates of emulsions of infected louse excreta into volunteers. The filters used were in one case a Chamberland "F," in the others British "Berkefelds." Only one of these experiments was definitely successful, and in this case the filter used was not satisfactory. The results were as follows:

(a) *Experiment 153*.—20 c.cm. of a 1 per cent. emulsion filtered at a negative pressure of 350 to 400 mm. of mercury through a Chamberland "F" filter. Result, negative.

(b) *Experiment 187*.—10 c.cm. of a 1 per cent. emulsion filtered at a negative pressure of 200 to 300 mm. of mercury through a Berkefeld were injected intravenously. The man had a rise of temperature to 103° C. and abdominal pain on the eighth day from the inoculation, but no further symptoms suggestive of trench fever.

(c) *Experiment 190*.—10 c.cm. of a 1 per cent. emulsion filtered at a negative pressure of 300 to 350 mm. of mercury through a Berkefeld were injected intravenously. The result was negative.

The above filters kept back *B. prodigiosus* under conditions like those of the experiments.

(d) *Experiment 199*.—35 c.cm. of a Berkefeld filtrate of a 0.5 per cent. emulsion were inoculated subcutaneously. The filtration took place at a negative pressure of 600 to 740 mm. of mercury. The man developed trench fever on the sixth day and had several relapses.

(e) *Experiment 200*.—49 c.cm. of a Berkefeld filtrate of the same emulsion as that used in the last experiment was inoculated subcutaneously. The filtrate was obtained at a