the bone and the temporal muscle, and the wound closed in the usual manner.

The next day the pulse was 86, the respirations 22, and previously subnormal temperature had risen to 99.2°. By the second day the plantar reflex had become flexor in type.

On the third day he said that "there was a little man kept running along the bed-rail"; "everything has a red appearance"; "the room is queer-shaped—there are so many corners in it." The auditory cortical centres were also disturbed, as he heard "some one talking to him behind the bed-head." The hallucinations ceased in a few days, and might have been prevented by operating on the right side, had this been feasible. In the case of another right-handed patient, published in 1912,¹ the wall of the posterior horn of the right lateral ventricle ruptured spontaneously during the operation, but no such symptoms occurred during convalescence.

Three months later vision was as follows:

R. $\frac{3}{6}$; 1 Jaeger at 13 in. L. $\frac{3}{6}$; 10 Jaeger at 13 in. No signs of papillitis. Right homonymous hemianopsia still present.

The patient sought relief from the disfigurement of a large hernia over the temporal area which displaced the upper part of the pinna downwards and outwards; he was quite prepared to undergo any operation which would open up fresh channels of escape for the cerebro-spinal fluid. Cushing's operation of retroperitoneal drainage through the lumbar theca was contraindicated, since lumbar puncture had demonstrated that the foramen of Majendie was blocked. We decided to explore the right cerebellum for tumour, and, if no tumour were found, to open up the posterior subarachnoid reservoir (Parkin's operation), and to utilize the deep tissues of the neck as a drainage area, if needful.

Second Operation.

The right lobe of the cerebellum was explored with a negative result. There was a gush of fluid on opening up the posterior extremity of the cisterna magna.

Recovery was uneventful, and, when re-examined two years later, the hernia had disappeared. The stretched skin over the left temporal area could be rolled between the fingers and thumb. The right suboccipital area was not prominent. The sight of both eyes had improved but the hemianopsia persisted. He has resumed his pre-war duties, the chief of which is that of driving a motor car.

The etiology, as is not unusual in these cases, remains unknown.

My thanks are due to Dr. Squire for much assistance in the conduct of the case, and to Mr. Haigh, of the Royal Infirmary, Halifax, for taking post-operative skiagrams of the decompressed area of the skull.

REFERENCE.

¹ BRITISH MEDICAL JOURNAL, March 16th, 1912.

FLAT-FOOT.

BY

LIEUT.-COLONEL HENRY SMITH, C.I.E., I.M.S.,

AMRITSUR.

A GOOD deal written on the subject of flat-foot has been far too much to show that the leg muscles attached to the tarsus were intended by Nature to support and maintain the arch of the foot. It seems to me that Nature no more intended the leg muscles attached to the tarsus to maintain its form than that the corresponding muscles attached to the carpus were intended to maintain its form. The corresponding muscles attached to the carpus are plainly intended to move the joint and nothing else, and in my opinion the same holds as regards the leg muscles attached to the tarsus. They are merely intended by Nature to move the ankle-joint and the joints of the tarsus. If these muscles had to support the weight of the body or a portion of that weight they would very quickly lengthen and be useless for that purpose. The strong ligaments of the sole have something to do with the maintenance of the arch, though if the weight of the body is excessively thrown on them, as in the development of flat-foot, they too give way and lengthen. The cause of the evil in flat-foot and in over-high instep is to be found in the skeleton formation. Normally the heel, the outer side of the tarsus, and the metatarsals carry the weight at rest and in motion. If

the formation is such that the weight of the body is thrown too much on the inner side of the foot we have flat-foot; if too much on the outer side of the foot we have over-high instep, which cripples a soldier in marching as much as a flat-foot. When for knock-knees we cut the thigh bones in a child of, say, 10 or 12 years of age and set the legs straight, we would have reason to be anxious about the loose ankle-joints, due to the undue looseness of the external ligaments. When we see the patient five or six years later we find that Nature has "taken up the slack" and that the ankle-joints are perfect, as well as the general symmetry of the limbs. Here we have merely dealt with the skeleton. We have adjusted it so that the weight is conducted as in a normal case, and Nature takes care of the rest.

I know a farmer in Ulster whose amusement from boyhood was to shoe a crooked-footed horse. He had no teacher in the art, studied the issues himself, and never read books on the subject. He had experimented to such a degree that I doubt if there is a professional or other horse-shoer in the empire—including veterinary surgeons—who could shoe a crooked-footed horse better than he, and exceedingly few who could do it as well. He was a born mechanical genius. I often talked the matter over with him, and though he had never heard of the centre of gravity or of muscle attachments, he had got to the bottom of the subject, and was scientific to the last degree. A few years ago I was spending an evening with him, and a flat-footed workman came up to us. I requested him to strip the feet, and asked the farmer how he would "shoe" him. He had not previously devoted any attention to orthopaedics—a term he had never heard of. He asked the man to stand in an attitude which a soldier calls "attention," and walked round him, looking at the legs and feet. He drew my attention to the fact that the inner side of his feet was not straight, but bulged inwards at and below the ankle. He then said the cause was plain; that flat-foot was due to the formation of the skeleton, which threw the weight of the body too much on the inner side of the foot, and that the remedy was plain. I then asked him about an over-high instep. He replied that it was the opposite of an over-low instep, and the cause must be the reverse, and the remedy the reverse. He said he would get the bootmaker to make the insole (which is usually made of thin leather all the same thickness) of the heaviest sole leather for the flat-foot, full thickness from heel to toe on the inner side and pared away like a wedge to the outer side, and the reverse for an over-high instep. By this means he said he would throw the weight further out on the foot in the case of a flat-foot and further in in the case of an over-high instep. In the case of the over-high instep he said the instep would sink under this treatment. He also said it would require some experimentation to find the most suitable thickness of the thick side of the insole to suit the individual case. I then explained to him the steel plates which orthopaedists use for flat-foot, and that they claimed that they supported the arch. He laughed at the idea, and said that if the plate supported the arch, it would press on it and would certainly lame the man, and that any benefit it might have was due to raising the whole inner side of the foot, and thus throwing the weight of the body further out.

I am personally convinced that the farmer was right both as regards the causation of these two conditions and as to the method of treating them, and I would respectfully invite the attention of those who have to deal with these conditions to this aspect of the question.

KEIFFER has published in *Le Scalpel* the results of a study of the physical and mental state of Belgian women during the war. The effects of privation began to show themselves in 1915 in disturbances of nutrition, wasting, enlargement of the glands of the neck, and rickety lesions. Tuberculosis caused great ravages, especially in young girls. In married women, long before the menopause, amenorrhoea was very common; with the signature of the armistice it diminished. Keiffer attributes it to the insufficiency and bad quality of the food, as well as to the mental suffering caused by the German occupation.

THE Massachusetts Commission of Mental Diseases states that 200,000 persons in the United States suffer from epilepsy, and that only 3.5 per cent. of these are cared for in institutions. Thirteen States in the Union have colonies providing for the care of 7,000 epileptics.