Cortex Adrenalis and the Reproductive Function.

There is evidence to show that one of the functions of the adrenal cortex, the cells of which are filled with lipid substance, is to provide lipid for the building up of the myelin of the developing nervous system. The evidence in favour of this is as follows:

1. A correspondence in size of the adrenal cortex and the brain in the animal series.

2. The great diminution in size of the adrenals in anencephalous monsters.

But it has other functions—namely, the storage of lipid—which, as occasion arises, can pass into the blood and keep constant the supply of lipid to the reproductive organs, where it constitutes the raw material for formative nuclear activity. This functional correspondence of the adrenal cortex and sexual glands is indicated by certain pathological facts—namely:

1. Adenoma of the suprarenal cortex has been found associated with marked sexual precocity.

2. Among my cases an active spermatogenesis in a man aged 69, who died after many years' residence in the asylum suffering with dementia of arrested general paralysis, was associated with a large adrenal cortex.

3. In four male cases of well-marked dementia praecox, with regressive atrophy of the testis, the adrenal cortex was narrow and the cells contained much less lipid than in other cases of mental disease dying in the asylums. I have only paid attention to this matter lately, but it is being followed up to see if there is a correlation.

4. The cortex adrenalis in the teleostei and elasmobranch fishes is a separate gland. The medullary substance, with its chromaffin, is a part of the sympathetic nervous system.

5. Rats fed upon suprarenal gland showed hypertrophy of the testes; other organs did not show hypertrophy (Swale Vincent).

6. The cortex adrenalis is developed close to, or forms part of, the genital ridge, and in one case of dementia praecox, where there was only one testicle, the adrenal gland of one side weighed less than one-half that of the side in which the testis was present, and this difference in weight was mainly due to a diminished cortex.

I have observed in a large number of cases where the patient has died from a disease in which there has been microbial infection and consequent toxæmia a marked diminution of the amount of stainable lipid in the adrenal cortex, and I agree with Elliot that this adrenal lipid probably passes into the circulation to form antitoxins. Now, it is a remarkable fact that however much the lipid may have diminished, on this account, in the adrenal cortex, it does not seem to have diminished in the testis. Possibly this may account for the fact that, in all manner of chronic microbial infections with toxæmia, active spermatogenesis may nevertheless be seen in the seminiferous tubules. This supports the premise that these lipoids form a protective barrier by virtue of the free atomic oxygen liberated upon their surface; thus affording a special provision of nature to protect the germ cells from injury, as well as providing at the same time the oxygen and raw materials necessary for the most fundamentally vital cell activity in the body.

The Oxidase Reaction Applied to the Cortex Adrenalis.—According to Marinesco, the cortex adrenalis does not give the oxidase reaction because the contents of the cells stain uniformly and not as granules, but some cells from which some of the lipid has escaped do show granules of lipid and these stain in a similar manner to the granules in the testicles.

(To be continued.)

SURGICAL TREATMENT OF FACIAL PARALYSIS.

BY

GEORGE FENWICK, F.R.C.S.,

MAJOR, N.Z.M.C.,

AURAL AND OPHTHALMIC SURGEON, No. 2 NEW ZEALAND GENERAL HOSPITAL.

EVERY large military hospital has had experience of traumatic facial paralysis, where restoration of function by nerve repair or nerve grafting has not been possible. In cases of comminution of the petrous bone end-to-end repair cannot be effected, although grafting or anastomosis may still remain practicable; in lacerated parotid wounds the main trunk has already broken up into the parotid plexus, and repair of the diverging branches does not come into the realm of practical surgery.

The deformity is a very terrible one, and any treatment that offers hope of cosmetic improvement should be welcome to surgeons.

Early in 1917 I attempted to do something for these patients by grafts from neighbouring muscles, and in successive cases to employ the method, modifying the procedure by the result of experience.

I describe the operation in the form I finally adopted (Fig. 1):

Preliminary to anaesthesia, the non-paralysed side of the face is inspected, and the position of the lower half of the nasolabial furrow marked in; a second vertical marking is made almost in the centre of the cheek in the position of the dimple that becomes evident when laughing. A third marking is made, beginning below the external canthus and taking the direction of one of the lines of the crow's foot. The side of the head is shaved.

An incision is made through the skin in the hair line from

the zygoma to the upper limit of the temporal fossa, directed slightly backwards so as to be parallel to the underlying fibres of the temporal muscle. The skin is then undermined forwards and backwards to expose the temporal fascia; two parallel incisions are then made, again from the zygoma to the limit of the temporal fossa through the fascia and muscle down to the bone; these incisions should run in the direction of the muscular fibres and should include a strip of muscle as thick as a man's thumb; it is important for the preservation of the nerve supply and for the cosmetic result that fully this bulk of muscle should be used. A smaller slip anterior to this is similarly isolated, and both are detached from the underlying bone.

An incision is now made under the eyelid in the marking representing one of the lines of the crow's foot, the skin is undermined, the anterior smaller slip of muscle drawn through,

made taut, and sutured to the fibres of the orbicularis palpebrarum with catgut (Fig. 1, T, and Fig. 3, T₁).

Incisions are made in the line of the cheek furrow and the lower half of the naso-labial groove, and the skin tunnelled to make a continuous passage from the temporal wound to the corner of the mouth. The large slip of muscle is then drawn down over the zygoma through the channel, sutured with

catgut to the superficial fascia exposed in the cheek wound, and to the muscular fibres of the orbicularis oris below and slightly mesial to the corner of the mouth (Fig. 3, T₂).

The facial incisions are sutured with horsehair: the cut margins of the temporal muscle are brought together as far as possible with catgut (it is extremely difficult to get apposition), and the temporal wound closed with silkworm gut.

Paradism and massage should be employed early, and the patient should daily exercise his facial expression before a glass. The grafted straps of muscle will continue to function.

In the first operation of this kind which I attempted I grafted a slip of the temporal muscle into the orbicularis palpebrarum and a slip of the masseter into the orbicularis oris (Fig. 1, M); Stenson's duct was guarded by a probe inserted and tied to a tooth. Considerable difficulty was

FIG. 1.—2. Denuded area in temporal fossa. T, Slip of temporal muscle inserted into orbicularis palpebrarum. M, Slip of masseter inserted into orbicularis oris.

FIG. 2.—2. Denuded area in temporal fossa. T, Temporal muscle. T₁, Slip of temporal muscle inserted into orbicularis oris.

FIG. 3.—2. Denuded area in temporal fossa. T₁, Slip of temporal muscle inserted into orbicularis palpebrarum. T₂, Slip of temporal muscle inserted into orbicularis oris.

experienced in getting the slip from the masseter, and there are no compensating advantages. I have performed the temporal muscle graft without making use of a cheek incision, but I prefer the method described, and the additional scar, lying as it does in a natural skin fold, is almost imperceptible.

The cavity left in the temporal fossa after swinging down the slip is of considerable dimensions, and to obliterate it by approximation of the cut edges of the muscle is almost impossible. I intend in the next case to implant gluteal fat.

I have had good results before in paralytic ectropion by means of the ordinary Kuhnt-Dimmer operation; I think, however, that the sling of living muscle is more likely to ensure permanent improvement than the mere shortening of the lower lid.

I have not attempted to do anything for the expressionless brow; the patients I had were instructed to refrain from raising their eyebrows and frowning, and were very successful in abandoning this expression of emotion; in a deeply wrinkled forehead where the facial expression could not be adequately controlled I should feel inclined to excise strips of skin and fascia in the line of the frontal furrows. The immediate cosmetic result of the operation is good in repose. The drooping corner of the mouth is slung up into a natural appearance by the strap of inextensible fascia and living muscle, and the hollowness of the cheek is obliterated by the bulk of the graft. It is not to be hoped that the implantation of a graft from a mandibular muscle will restore the infinite variety of facial expression commanded by the complex sheet of muscle it replaces, but an appearance of tone and life is conferred that gratifies both patient and surgeon; and by practice before a glass the patients learn in some degree to associate the contraction of the temporal slips of muscle with that of the zygomatics and the risorius of the unaffected side.

Two years ago a writer in the *Annals of Surgery* contended that the neuro-muscular spindles of a graft multiplied and extended into the fibres of the paralysed muscle, and illustrated his article by microscopic sections that seemed to prove his point. I cannot feel optimistic about the likelihood of this occurrence, but if any such process took place the cosmetic gain would be further enhanced.

Much has been learned in military surgery that is destined to be of value in civil life; cases of facial palsy will be always with us, and I believe that where nerve repair is impossible, or has failed, the operation I have described will be the recognized method of treatment. The diagrams explain themselves.

OBSTETRICS: PAST, PRESENT AND FUTURE.*

BY

ALEXANDER STOOKES, F.R.C.S. EDIN.,

HONORARY SURGEON TO THE LIVERPOOL MATERNITY HOSPITAL AND TO THE LIVERPOOL SAMARITAN HOSPITAL FOR WOMEN.

In this paper on Obstetrics, the Cinderella of the Medical Faculty, I wish to present to the society the changes which have taken place, and more especially those which are now going on—changes not so much in technical points which are more suitable for presentation at a meeting of specialists, but changes affecting the community and our profession as a whole. Those of us who were concerned with the discussions and committee meetings before the National Insurance Act was passed will remember how difficult it was to get at what the medical profession wished or at any plan upon which we could agree. Since the Act came into force I have never met any one who was satisfied with its working. Even men with a panel of 5,000 are not pleased.

At present even greater changes are looming ahead, and in obstetrics have already taken place without the bulk of the profession realizing how we are drifting. In our archives we have a most interesting record of obstetric work done in Liverpool. Henry Park was a pupil of Percivall Pott at St. Bartholomew's, and studied in Rouen; he was surgeon to the infirmary, a partner with Alanson, and was the inventor of the operation for excision of

joints. His midwifery record, which we possess, extends from 1769 to 1830, and enumerates 3,900 cases extending over fifty-one years. The first 1,000 cases took fourteen years, the second thirteen, and the third twelve years. It was not until he had done 3,300 that he slackened in this work, and the last 100 cases extend over eight years. The record says nothing of mortality, but singularly few abnormal cases are noted. There were only two forceps cases, one a brow presentation; the difficulty in the other case is not recorded. There were five crotchet or craniotomy cases, four cases of placenta praevia, and only two cases of convulsions. Evidently child-bearing in those days in Liverpool was more simple than in these times.

A later record, of even more interest from the obstetric point of view, is the very detailed account left by Francis Ramsbotham of the Royal Maternity Charity from 1820 to 1850. From 1828 to 1850 in 48,996 deliveries there were 220 deaths (4.5 per 1,000), 162 were puerperal deaths, 58 non-puerperal; 66 of the former were from haemorrhage. There were only 43 cases of eclampsia (about 1 per 1,000); of these 27 were before delivery with one death, 16 after delivery with 3 deaths. Stillbirths came to 2,723, equal to 5.4 per cent.

These two records cannot be considered as typical of obstetric practice of the time, but show what could be done. Dr. Matthews Duncan, writing in 1866, took 31,277 cases occurring in the private practices of the best known obstetricians and found there were 272 deaths; he concludes that 8.2 per 1,000 might be considered the average mortality of childbirth. It was well known that hospital practice gave a much higher mortality in most instances. Duncan came to the conclusion from investigation of the returns of well managed hospitals, such as the Dublin Rotunda, that the mortality was about the same as in private cases. Florence Nightingale in 1871 said that childbirth in a hospital in London was four times more dangerous than in the towns of England, where the mortality was 5 per 1,000 births.

Death Rates in Hospitals in London, 1865-1875.

British Lying-in	...	...	19.4 per 1,000
City of London	...	...	14.3 " "
Queen Charlotte's	...	...	22.3 " "

Obstetrics has always been, as I said, the Cinderella of the Medical Faculty. It was considered a branch of the profession hardly respectable. The College of Physicians precluded practitioners of midwifery from their Fellowship; the College of Surgeons would not allow such a one to sit on its council until 1828. It was only in 1886 that an amendment of the Medical Acts required all registered practitioners to qualify in obstetrics as well as in medicine and surgery.

TABLE I.—Mortality of Mothers per 1,000 Births Registered (England and Wales).

Decennium.	All Causes.	Sepsis.	Accidents of Childbirth.
1851-1860	4.7	1.5	3.2
1861-1870	4.7	1.70	3.00
1871-1880	4.72	2.18	2.54
1881-1890	4.73	2.68	2.05
1891-1900	5.09	2.36	2.73
1901-1910	4.01	1.76	2.25
1911-1917	4.02	1.4	2.59

When we turn to the Registrar-General's statistics for the past fifty years we find there is little if any less risk for a woman in childbirth to-day than there was in 1851.

The fall in mortality from causes other than sepsis has extended through five decades, whereas that from sepsis dates only from 1881-1890. Over the whole period the fall from non-septic causes is greater than from sepsis.

In the decade 1851-1860 the mortality from all causes was 4.7 per 1,000 births—1.5 from sepsis, 3.2 from accidents; if we take 1911-1917 the total mortality is 1.4 from sepsis, 2.62 from accidents, a total of 4.02 per 1,000.

No one event has produced so much effect in midwifery as the passing of the Midwives Act, 1902, whereby it became compulsory for any woman who practises as a midwife habitually and for gain to be on the midwives'

* A paper read at the Liverpool Medical Institution, November 21st, 1919.