2. Arrest at Puberty.

The following are examples of arrest at or soon after puberty:

Boy, aged 16, died of chronic morbus cordis of long standing. The tubules were atrophic and showed miniature development. As in the testis at birth, the tubules were separated one from another by abundant interstitial tissue. There was no evidence of formation of spermatocytes or spermatozoa; in fact, the epithelium retained to a great extent its embryonic character, and there was little evidence of mitosis of nuclei. The membrana propria was thickened, and chronic passive congestion had given rise to an interstitial fibrosis; no interstitial cells could be discovered.

As a contrast the following acute case of morbus cordis may be cited:

A young man aged 21 died suddenly of cerebral haemorrhage caused by aneurysm due to infective embolism. There was ulcerative endocarditis. The seminiferous tubules are of normal size and are packed together with very little interstitial tissue; normal spermatogenesis with abundant lipid in the interstitial cells, the spermatogonia, and cells of Sertoli.

Chronic disease of long standing before puberty is associated with complete arrest of spermatogenesis at puberty. This fact has been shown by an important research of Kyrle¹ on the disturbances of the development of the male generative organs at all ages from birth to adolescence. He has demonstrated the fact that chronic constitutional diseases, such as tuberculosis, are associated with arrest of development. One hundred cases were examined; among these, seventy-six were highly undeveloped; of the remaining twenty-four by no means all were normal; more than half showed distinct underdevelopment. Only ten were certainly normal.

Kyrle states that at the first glance it would seem that the under-development of the testes in chronic tuberculosis was due to the effect of the toxin. He points out, however, that out of thirty-nine dying in the first year of life, twenty-nine had undeveloped testes, but only one of these died of tuberculosis. He thinks this proves that the undeveloped organ is already undeveloped when the individual is affected by the disease.

He concludes that a considerable number of children are born with undeveloped reproductive organs. The development of the whole organism is directly under the influence of the reproductive organs. Individuals with undeveloped reproductive organs have less vital resistance to disease than those with normal reproductive organs. Among the children dying with undeveloped seminiferous tubes far the greater number were physically undeveloped. The greater number of those dying in early life had undeveloped testes. Kyrle asks the question whether this is a kind of natural selection by which individuals of little genetic worth are cut off in early life.

REFERENCE.

¹ Kyrle, Dr. Josef: Ueber Entwicklungstörungen der menschlichen Keimdrüsen in Jugendalter. *Wien. klin. Woch.*, November 10th, 1910, No. 45. Aus dem pathologisch. Institute in Wien.

(To be continued.)

An Address

ON

CANCER.*

WITH DEMONSTRATION ON THE EPIDIASCOPE.

BY

PROFESSOR J. RUTHERFORD MORISON, F.R.C.S.

A WEEK ago, when I was at the War Hospital, Gosforth, Major Reader, on our round, showed me five cases of malignant disease in soldiers. The first had a cancerous stricture of the oesophagus; the second had cancer of the stomach; the third had intestinal obstruction due to general cancerous peritonitis; the fourth was a man with hopeless inoperable carcinoma of the rectum; and the fifth a man with cancer of the prostate and secondary deposits.

These soldiers, of course, were not old men. The man with cancer of the rectum was only 20 years of age. We used to think of cancer as being associated with old age. So it is; but there is an increasing amount of cancer in the country, and it is increasing more in the younger people than in the older, I think. When I saw the wounded

soldiers and saw those hopeless cancerous soldiers together, I thought what a blessing it was to be wounded instead of to have been caught with cancer.

The fact is this, that tubercle and syphilis and cancer—all of them very closely related—kill more people every year than were killed in any year of the war. We ought to face these facts and make war upon them. We have finished with the Huns, now we ought to make war upon cancer; and how is it to be done?

My brother, the medical officer of health for Cumberland, wrote a paper about a year ago in which he advised that periodical examination of every one in the country should take place at least once a year—he said once in six months, but at least once every year. That is the only way to deal with and to prevent these scourges, or discover them early enough for successful treatment. Sanatoriums for tuberculosis have been proved, almost proved, to be useless except for the individual. The treatment of syphilis in the new venereal centres may be satisfactory enough for the individual, but is of small national benefit.

What we want to do is to have everybody examined once a year and all the syphilitics to be sent to an island and to be kept there until they are cured; nobody to be allowed to enter or leave the island without a doctor's passport, and nobody to enter England without being examined by the Port Authorities. This would eradicate syphilis in two years as hydrophobia was banished. We know that this is true, and, whether popular or not, we ought to have the courage to preach it. That is the only way to prevent, or at all events to recognize, early cases of cancer.

We know about cancer a few things; we do not know the great thing—its exciting cause—but we do know a large number of predisposing causes, so that in many cases it is possible to prevent cancer. Then we know that cancer can be removed by operation, and, if a good operation is done early enough, with a large and steadily increasing percentage of success. I do not say operation is the last word in cancer; it is not. There will eventually be something better, but meantime we are to deal with what we know, and this is that from 75 to 90 per cent. of early cancers can be cured by a satisfactory operation.

PREDISPOSING CAUSES.

The following specimens and photographs, shown by the epidiascope, will impress upon you the predisposing causes of cancer and the importance of early diagnosis.

1. Here is a clay pipe smoker's epithelioma of the lip. It at first appeared as a little scaly, corny patch, which may have existed for three, four, five, six, and even ten years, and then it got a little crack in it. Even the crack could heal with some ointment, and things returned to their normal condition; but there came a day when this crack spread and it developed into an epithelioma of the lip.

2. Here is a chimney-sweep's cancer. This is a very unusual condition, because there are multiple primary foci. Now if you see multiple sores you think it cannot be cancer, but it is so in this case. From irritation of soot in the breeches we may get multiple foci of cancer.

3. This photograph is from an infirm patient of mine. He was a paraffin worker and he got a paraffin burn, which developed, as you see, into an epithelioma.

4. Here is another example of chronic irritation—chronic superficial glossitis. This tongue spells syphilis, tobacco, alcohol, leucoplakia, and cancer.

5. A case of leucoplakic tongue, one developing carcinoma in the middle. The tongue had been like this for ten years. Some leucoplakic tongues are curable by x-ray or radium treatment—about the only thing as yet which seems to do them any good.

6. A case of leucoplakia, also of long duration—very long duration—but no cancer. Of course you would not advise this man to have his tongue out. You would knock him off his tobacco and alcohol, and give him some antisymphilitic remedies and x-ray treatment, and get him to come up for observation every three months; at the least sign of cancer you would have the tongue out.

7. Here is an instance of leucoplakia of the vulva. Leucoplakia does not develop only in the tongue; it may come on the vulva and on the penis, and wherever it is it is apt to develop an epithelioma.

8. Here is a burn scar. In my own experience of old burns, if the scars do not heal very few of the cases escape cancer. This man had his leg burnt when he was young, and at the age of 30 an epithelioma developed on that burn. The scar never was quite healed. It broke down from time to time and discharged a little and then skinned over, but it never was a sound scar. That scar ought to have been removed and the scar area replaced by skin grafts or flaps from the other or the same leg.

X rays, of course, are another well-known cause of burn giving rise to epithelioma. I have seen quite a number of

* Given before a Scientific Meeting of the North of England Branch.

lupus cases treated by x rays that have or have not been cured of their lupus, but have developed epithelioma in the same area.

Cancer in China is very common in the pharynx and in the oesophagus, and for a long time it was difficult to find any explanation of it. Eventually it was discovered that it occurred only in men, that women almost never developed either cancer of the pharynx or of the oesophagus, and the explanation of that was probably found. It was this, that at meals men are served first, and they get, and swallow, their rice boiling hot; ladies have to wait until the men are satisfied, and they get their rice cold.

9. The case of an old field lady. She worked in the fields, and her left eye was not very good at holding tears; the tears ran down the face, and here you see that the chronic irritation from cold, exposure, and tears has caused an epithelioma.

10. Here is a curious oesophagus with a stricture at the lower end, a congenital stricture, and here at the upper end is a cancer which has burrowed its way from the oesophagus into the trachea. Owing to the chronic irritation of food in the oesophagus, cancer has developed at the upper end of the dilated oesophagus.

11. Here is a carcinoma of the penis with the usual predisposing phimosis.

Simple Tumours.

Another predisposing cause of cancer is the presence of simple tumours. We separate simple tumours and malignant tumours, and we think we have done a clever thing; but there are many simple tumours that tend very strongly to malignancy. Most simple tumours predispose to it to some extent, but there are simple tumours so closely associated with cancer that I want to draw your attention to them.

Some of you who are old enough will remember the case of the ex-Emperor of Germany's father, and how the surgeons, English and German, quarrelled about it. He had either a simple papilloma or a malignant papilloma of the larynx. Now a papilloma is almost always, in a person above 40, potentially malignant; and it proved so in the German Emperor's case. He died of cancer of the larynx. On removal of a portion for microscopic examination it was reported to be simple.

These specimens will illustrate my point:

12. This is a woman with scirrhus of the breast. She had got a tumour of the thyroid. Now I have been struck with the frequency with which women who have thyroid tumours develop cancer of the breast.

13. Here is another papilloma that is very often indeed followed by cancer. I have operated upon a great number of these cases, and 75 per cent. of them have come back, after some years even, with cancer on the site of the papilloma of the bladder. Now these cases are very easily discovered. Cystoscopy will show you "red seaweed trees" growing into the boracic lotion in the bladder.

14. Here is another form of papilloma that is very apt to develop into carcinoma, a papillomatous condition of the ovary. This ovary is cystic, and in connexion with the cysts there are papillomata.

15. Here is a condition in the rectum described by proctologists as mucous polypi. These mucous polypi and adenomata and papillomata in the rectum and colon are very frequently indeed associated with cancer, and it follows their development.

16. This is an ordinary thyroid tumour. In a man you think that is a simple growth, and so it is. It is now not doing the slightest harm, but a certain percentage of these in later years develop into carcinomas. If a patient with a thyroid tumour finds that it is beginning to grow larger, and especially if it grows larger quite quickly after he is 40 years of age he is very likely developing a carcinoma or a sarcoma of the thyroid.

Another familiar type of tumour that is apt to end in a carcinoma is what we call parotid tumours. You find, the first time you see such a case, a nice movable tumour the size of a half-orange, and you cut it out without any difficulty, and you send it to the pathologist. He writes back to tell you that it is a myxo-chondro-myo-adenofibroma. The patient comes back to you after having it removed, with a recurrence, three, four, or even five years perhaps, and next time the tumour is a little more fixed—more difficult to get out; then it recurs more quickly still, and in the end that is the condition—a malignant growth of the parotid. So these parotid tumours, however simple they may seem, ought to be all removed in their earliest encapsulated stage.

We come now to another thing—sarcoma. Of course sarcoma is cancer.

17. This condition has many names—von Recklinghausen's disease, neuro-fibromatosis, molluscum fibrosum; all these things are the same. This woman came out in little spots, and

they look like little soft fibromata, but when you take them between your finger and thumb they melt away—there is no resistance. These patients are apt to develop large sarcomatous nerve tumours.

18. This is another instance of the same condition. People are not familiar with them. I am surprised to find that very often even the best of our students have never seen such a case. Look at the patchy soft tissue on her hand; her breast has thickened patches, and the skin is discoloured in spots and patches.

19. This is a different type of disease—what the pathologists and other people call keloid. It is a fibroid tumour in cicatrix, and some say a fibroid sarcoma in the cicatrix. As a matter of fact it is malignant, because if you excise it you get a bigger and bigger thing. Every time you excise you get a bigger growth.

20. Here is a photograph sent by Dr. Burman, an old house-surgeon of mine now in South Africa. This native came to Dr. Burman because nobody would employ him. He begged to have these enormous keloid tumours removed. Dr. Burman removed one and sent me this photograph. The patient had one ear tumour taken away, the other left. The tumours grow from earring holes. You see the keloids on the rest of his body. These, Dr. Burman tells me, are the result of native medical treatment. The natives and lay doctors put lances and punctures into everything; they are where we were fifty or one hundred years ago.

How Cancer Commences.

Cancer begins always in one of two ways: as a tumour or as an ulcer, or I should say as an ulcer and as a tumour. Characteristics of the cancerous ulcer are these—that there is growth in excess of destruction.

The cancerous tumour has four characteristics: its consistency is hard, its surface is nodular, its edge is merging off into the surrounding parts (not like a walnut or potato in the tissues), and it is generally fixed and infiltrating the surrounding structures. These are the characteristics you look for in cancer.

21. Here is a cancerous ulcer. This is the anus and a cancerous ulcer of the rectum. The rectum is split open. You see the thickened edge of the ulcer all round. There is growth, there is destruction in the middle, there is growth in excess all round.

22. Here is what is called a Cock's tumour, a rare form of epithelioma. I think this is the only case I have seen. I got this photograph at the infirmary. A very exceptional case. You will see it is like a cock's comb, but Cock was the name of the surgeon. This is between an ulcer and a tumour. The tumour is ulcerated on its surface and nodular.

23. Here is a breast showing how cancer can commence. If it begins in the breast you always think of cancer there as a tumour, because in 99 cases out of 100 it is a tumour; but it occurs as an ulcer, and when it does (Paget's disease or Paget's eczema) there are ulcerated red patches round the nipple, which in this case is retracted. That woman had a tumour as well as an ulcer.

24. This is a typical scirrhus—a large, stony, hard nodular tumour in this breast, with shelving edges and tucking the breast up. In the majority of women the left breast is rather lower than the right. The right breast is much lower than the left in this case and the nipple is retracted—quite a characteristic appearance, sufficient for you to make a diagnosis.

Method of Spread.

We now come to the method of the spread of cancer. Of course, to know anything about the prognosis and treatment of cancer, you want to know how it spreads. It spreads just the same as tubercle and sepsis do, in the same four ways:

1. By continuity.
2. By contiguity.
3. By the lymphatics.
4. By the blood.

25. Here you see carcinoma of the stomach. This is a cancerous ulcer of the pylorus which spread all the way along, involving the whole of the pyloric end of the stomach by continuity. It lies next the colon, close to the transverse colon, which is involved by contiguity. The lymphatic glands along the lesser and greater curvatures are evident.

26. Here is an example of probable blood infection. Some people say that cancerous bones are the result of the lymphatic spread. In this case the humerus is broken in the middle, through a cancer secondary to scirrhus of the breast. It is very common to get bone cancer secondary to scirrhus of the breast. I have seen quite a number of cases operated upon, patients who complained of "rheumatism" in their arms, "rheumatism" in their back, and they had cancerous growths. You must not operate upon a patient who complains of "rheumatism," however favourable the case, until you have had an x ray to see if there is any growth or not in the bones.

27. This girl had a pigmented wart on her back. It grew hard and was cut off. Within six weeks she was in this condition, dotted all over her body with deposits of melanotic sarcoma or carcinoma.

28. You see here the metacarpal bone of a child. Here is the lower epiphysis of the radius. That child had sarcoma of its testicle removed and developed a growth in the bone.

Diagnosis of Cancer.

It would be correct to say that no case of cancer begins with pain. There is a popular delusion that cancer is always a hideously painful thing, and people come with advanced disease who say they did not come before because they never had any pain.

The early diagnosis of cancer is one of the more difficult problems. A patient over 40 years of age comes with haematuria. What is to be done? Send him for cystoscopy straight off. In eight times out of ten the man of over 40 with haematuria has cancer somewhere in his genito-urinary tract.

If a woman over 45, or after menstruation has ceased, has haemorrhage, and no simple cause can be found, the sooner she has her uterus out the better. Nothing may be found in one case out of twenty, but in the other nineteen there is carcinoma somewhere.

In nineteen times out of twenty, if a man comes to you over 40 years of age and he cannot swallow, and he shows you he cannot swallow, he has got carcinoma of his oesophagus. X ray with bismuth meal will clear this up.

The first symptom of carcinoma of the stomach is "indigestion." A man of 50 who has never had indigestion all his life and comes to you complaining of indigestion, which Gregory powder has not cured, nearly always has carcinoma. You must have him examined with a test meal to show whether he has got a normal acidity and have him x-rayed by a specialist. Now examination of the stomach by the radiologist is sometimes derided and said to be of no value, but that is because people have not got the right man to examine the stomach. The x-ray stomach specialist can diagnose cancer of the stomach in almost every case which comes early.

Stricture of the sigmoid colon is a very common result of carcinoma. These patients say they have never had any trouble with their bowels before. Now they have to take medicine and the longer the time the more the medicine, and the longer time the more the rumblings inside. If a patient of 50 comes to you saying he has got constipation and cannot get rid of it, and he has rumblings inside enough to keep his family disturbed or even to disturb himself—think of carcinoma, because if it can be diagnosed early it can be treated with great success.

ENLARGEMENT OF THE THYROID GLAND IN MALARIA.

By J. B. HUME, M.R.C.S., L.R.C.P.,

DEMONSTRATOR OF ANATOMY AT ST. BARTHOLOMEW'S HOSPITAL;
LATE CAPTAIN R.A.M.C.

ENLARGEMENT of the thyroid appears to have hitherto been an unrecognized complication of malaria. Some thirty cases of it came to my notice during the East African campaign; and I recorded some preliminary observations in a paper before the Dar-es-Salam Medical Society on May 18th, 1918. My interest was the greater because in June, 1918, during an attack of malaria, I developed a marked enlargement of the gland myself.

The condition is characterized by an acute enlargement of one or both lobes of the thyroid gland during an attack of malaria, and occurring some twelve to thirty hours after the fall of the temperature.

All the cases were among white troops, soldiers with varying periods of service in German East Africa, the majority of whom had had several attacks of malaria. None of them gave any history of goitre, or even of fullness of the neck previously; in several instances I had made a definite note of the absence of thyroïdal swelling on admission. In no case could I obtain a family history of thyroid disease.

Malarial parasites were found in the blood in every case, usually in rather scanty numbers. Subtertian forms predominated, but benign tertian, and mixed infections also occurred. No opportunity presented itself of studying the condition *post mortem*.

In a typical case the enlargement of the thyroid appeared suddenly, in the course of an attack of malaria, from twelve to thirty hours after the fall of temperature. The previous

pyrexia had lasted from one to three days, and no subsequent rise of temperature occurred. One or both lobes became markedly swollen, and somewhat painful and tender to touch. There was no redness of the skin, the consistency of the swelling was uniform, and it showed all the characteristics of a generalized enlargement of the gland.

Dysphagia, in varying degrees, was a prominent symptom. Pain and throbbing in the head were frequent. There was usually a slight muscular tremor. In only two cases was there any exophthalmos. Tachycardia was very constantly present, the pulse rate being usually in the neighbourhood of 100-110, but sometimes as high as 130. Some alteration of the heart sounds at the pulmonary base was common—either a systolic bruit, or an accentuation of the second sound. The spleen was enlarged at some stage of the attack in about 70 per cent. of cases.

The treatment was usually carried out on routine lines. Quinine bihydrochloride was given by the mouth, and in a few cases also by the rectum. The usual procedure was to give 5 grains, three-hourly, omitting one dose in the night, over the first four days. Then a smaller dose, 15 to 20 grains daily, combined with arsenic, increasing to the full dose over the relapsing period on the ninth, tenth, and eleventh days.

In most cases there was, in from four to ten days, a marked reduction of the swelling, but in my own case it has not yet entirely disappeared. In subsequent malarial attacks it has invariably enlarged for a time.

The following cases occurred in No. 84 General Hospital, Dar-es-Salam:

CASE I.

Lieut. J. M. C., aged 30. Temperature 101° on admission. Very scanty subtertian rings present. Slight enlargement of thyroid in an attack six months previously; scarcely any enlargement on admission. Nine or ten attacks of malaria altogether. Enlargement of both lobes occurred on the third day, twenty-four hours after fall of temperature. Dysphagia, dyspnoea, pain, and a fine tremor were all present. Pulse 120. No exophthalmos. Spleen enlarged. The usual quinine treatment was given. On the fifth day 3 c.cm. of adrenalin given subcutaneously. On the sixth there was a definite increase in the thyroid. There was a decrease from that time to the fourteenth day, when, following another slight rise of temperature, the gland again enlarged. When the officer was invalided it had not disappeared.

CASE II.

Pte. P. S., aged 20. Temperature normal, but pyrexia reported by medical officer on day before admission. Many previous attacks of malaria. Both lobes of thyroid enlarged, mainly right. Both benign tertian and subtertian parasites present. Dysphagia, slight dyspnoea, tremor, and some tachycardia. Pulse 103.

CASE III.

Spr. S., aged 20. Temperature 103° on admission. Subtertian parasites present. First attack of malaria. Bilateral enlargement appeared on second day. Slight exophthalmos and tremor present. Pulse 100-110. Spleen enlarged. Marked reduction in ten days, under quinine treatment.

CASE IV.

Capt. J. B. H., aged 25. Temperature 101.6° on admission. Second attack of malaria. Scanty subtertian rings. Pyrexia lasted twenty-four hours; enlargement of both lobes, especially left, occurred twelve hours after fall of temperature and reached its maximum twenty-four hours later. Dysphagia and dyspnoea present; neck painful and tender to touch. No tremor nor exophthalmos. Pulse 110. Spleen slightly enlarged. There was a definite reduction in size in seven days, but the swelling has never entirely disappeared. It increased three months later, following another attack of malaria, and was invariably larger after slight attacks. In July, 1919 (one year from first appearance), enlargement of left lobe was still present, but without symptoms.

The following three cases present rather different features, in that the tachycardia was much more marked and the thyroid enlargement, though of old standing, had in every case first arisen as a sequel to an attack of malaria:

CASE V.

Pte. T. P., aged 33. Eleven previous attacks of malaria. Subtertian parasites. Noticed choking, tight sensation in throat, and dyspnoea in four previous attacks. Bilateral thyroid enlargement present. Marked and persistent tachycardia; pulse about 140. Spleen slightly enlarged. Thyroid decreased under quinine and arsenic treatment. Tachycardia decreased very slowly, in spite of absolute rest in bed, Pulse about 110 when invalided.

CASE VI.

Pte. F. G. A. B., aged 22. Six previous attacks. Admitted with tachycardia. Was in another ward in No. 84 General Hospital with subtertian malaria three weeks previously. Noticed enlargement of neck before readmission, when in Camp