It is only fair to bear in mind that many cases were fitted with glasses without having recourse to retinoscopy, but in this note the results are built up from cases in which it was thought advisable to dilate the pupils.

THE ADMINISTRATION OF OXYGEN.

BY

LEONARD HILL, M.B., F.R.S.,

LECTURER ON PHYSIOLOGY, LONDON HOSPITAL MEDICAL SCHOOL.

OXYGEN is usually given in a thin stream through a nozzle or small funnel held more or less near the mouth or nose of the patient. I have seen a nurse hold it 6 in. away. This method is more than inefficient; it is absurd. The oxygen diffuses into the atmosphere, and the patient inspires air enriched very slightly with oxygen. Analyses of the pulmonary alveolar air obtained from subjects to whom oxygen was administered in this way showed an increase of only 1 to 2 per cent. If a most wasteful current was used and the funnel held close to the mouth, the increase was at most 10 per cent. Patients who are ill and fretful, say with pneumonia, cannot bear a tight-fitting mask. Thus oxygen cannot be administered in the most economical way by means of an anæsthetic mask fitted with breathing bag and caustic soda to absorb the exhaled CO_2 . If such a method could be borne, a 20 ft. cylinder would give a supply for continuous breathing of oxygen, which would last all day, for a patient in bed uses only about 250 c.cm. per minute, and there are about 56 litres in the cylinder.

Ludicrously inefficient as the nozzle and funnel methods are, I have known of a practitioner going a step further in absurdity, in allowing the oxygen to escape in the patient's room! A man might as well hope to affect the composition of the sea by emptying his bladder into it.

It has been ascertained that in places situated at high altitudes, such as Mexico City, it gives the patient who has pneumonia much the better chance if he is rushed down to the plains. This shows the importance of the partial pressure of oxygen in maintaining the strength of the heart, etc., while the invading microbe is defeated by the rise in the immunizing powers of the blood. It therefore seems well worth while to try the effect of a still higher partial pressure of oxygen on patients situated at sea level.

It has been ascertained by Lorrain Smith, and by J. J. R. Macleod and myself that high pressures of oxygen produce inflammation of the lung. This is only brought out in mice by a very prolonged exposure (two or three days) to a partial pressure of 70 to 80 per cent. of an atmosphere—that is, four times as much oxygen as normal. The miners who practise wearing the Fleuss rescue apparatus for use in mines at the Howe Bridge and other rescue stations now set up in the colliery districts breathe 80 to 90 per cent. of oxygen for two hours or more, and never suffer in the least.

There is abundant evidence that the breathing of this partial pressure of oxygen has no ill effect. Börnstein has even breathed oxygen at two atmospheres pressure for half an hour without harm. Undue alarm as to the poisonous effect of oxygen has been felt by the medical profession. Oxygen does not influence the metabolism of the resting man at all. The common idea that the processes of combustion in the body are accelerated by breathing oxygen is erroneous. A man doing very

hard work is benefited by breathing oxygen because he does not then suffer from a shortage of oxygen, and so works more easily. Shortage of oxygen leads to the production of lactic acid, and this adding its effect to that of the carbonic acid in the blood increases the dyspnoea. The same must hold good with men ill of pneumonia. The problem is to contrive a mask which will allow a patient with pneumonia to receive an efficient supply of oxygen and yet will not render him uncomfortable or fretful. Messrs. Down have made for me a celluloid face-piece, to which a curtain of washable material is attached. The handle of the mask is also the inlet tube for the oxygen. The current of oxygen flows over the mouth and nose of the patient. The current is turned on until an agreeable cooling effect is obtained, and no sense of closeness is felt when the mask is applied to the operator's face. The operator, by applying the mask to his own face, will soon get to know the right degree of hissing noise in the tube. The patient inspires the atmosphere confined under the mask, and as a current is used sufficient to blow away the CO_2 , a very high percentage of oxygen is obtained in the alveolar air—for example, over 70 per cent.

Messrs. Down have also contrived for me a heating box which can be used with the mask for evaporating alcohol and adding vapour of alcohol to the oxygen. An incandescent electric lamp is placed at the bottom of the box; one of the lamps

in the ward or room can be inserted into the socket at the bottom of the box, and the wires from this attached by a plug to the socket from which the lamp was taken. A pot containing whisky or brandy, 2 parts, and water, 1 part, is placed in the box above the lamp. The current of oxygen is led through this box. There is no need to use a heating box or to wet the oxygen, unless alcohol or some other vapour is required to be given. Using the mask alone, I find patients like the cool stream to play upon their face. The

stream has to be made forcible enough to have a cooling effect, otherwise the patients push away the mask.

In the figure a water manometer is shown attached to the heating box. I devised this as an index for measuring the supply of oxygen, but experience has shown me that it is troublesome in use; the uninitiated cannot control the tap properly and blow out the water from the manometer. A gauge is not needed. It suffices to turn on the current until a distinct pleasant cooling effect is produced on the face, and learn to use the hissing noise of the current as an index. This method of giving oxygen is thoroughly efficient, and has produced most striking results in two cases of pneumonia on which I have tried it. The irregular pulse with its extra-systoles becomes regular, stronger, and less rapid, the ashen colour of the lips becomes red, and the patient recovers consciousness. In the case of an old retainer in my household dying of pneumonia, the administration of oxygen time after time restored her to consciousness. Aroused from that sleep which ends in death, she said, "Why, you have woken me up," and answered questions, only to relapse again soon after the oxygen was withdrawn. The return to consciousness makes the patient sensible of his illness, it improves his condition but he feels his distress, and therefore may ask for the oxygen to be discontinued.

In the struggle with the invading micro-organism oxygen helps to maintain the strength of the heart and so preserve the patient until the process of immunization is established and the disease arrested. To give oxygen most efficiently we require to place the patient altogether in a superoxygenated atmosphere, wherein he can live and be

Oxygen mask and heating box (Down Brothers). A, The mask, which can be used with or without the heating box. C, The cup in which alcohol and water are placed above the electric lamp. The ingoing current of oxygen blows over the cup. D, The manometer, the use of which I have discarded.

nursed and fed. This is a very difficult condition to realize. I am now experimenting with an oxygen bed—an airtight chamber into which the bed and patient can be placed. An account of this I hope to give in a subsequent note.

These researches are assisted by a grant from the London Hospital Research Fund.

Memoranda:

MEDICAL, SURGICAL, OBSTETRICAL.

SPHYGMOMANOMETRY AND PACHON'S OSCILLOMETER.

I was interested in Dr. Watson Wemyss's observations on Pachon's instrument (BRITISH MEDICAL JOURNAL, December 2nd, 1911, p. 1472). It may interest some of your readers to know my own experiences.

I have for some time been using this instrument, and have made several hundreds of observations with it during the season at Vittel and since at Beaulieu. In order to make it comparable I coupled it up with a Riva-Rocci mercurial manometer; we worked out a table of corrections, and these show very much higher readings for the Pachon than for the mercurial, and, like all instruments constructed on the anæroid principle, the errors are not proportionate. With the oscillogram it is quite as troublesome and uncertain to find the two critical points as with the Riva-Rocci. Furthermore, the time taken up in reducing the pressure to find these points is much greater, and if patients have high tension they complain a good deal of pain of the prolonged compression. I am about to try Oliver's auscultatory method for the Riva-Rocci, and a double armlet suggested by Amblard but not yet carried out, in which the lower cushion is used to determine by an indicator the first passage of blood through the arteries as the arm is decompressed by the upper cushion.

I think that the actual amplitude of the oscillations of Pachon's instrument may in the future give some valuable information, and I am collecting observations with that object.

The armlet of the Pachon is incorrect—first, on account of its narrowness, and, secondly, because the application to the forearm does not cut off the circulation between the radius and the ulna. I soon abandoned this, and replaced it by the broad armlet on the upper arm. The old strap method of fixing this I found took too much time, and was too fumbling a process. I tried glove pressure buttons, but the pressure on the arm, when closing them, was too painful for the patient. I then designed three blunt hooks and eyelet holes on a broad canvas strap, and this was carried out by Hawksley. It is ideal, being put off and on each in one movement. I find I economize one minute in each of these processes, which mounts up to near upon an hour when one has twenty or thirty observations to make of a forenoon.

Another question still unsettled is, What is an *average normal* blood pressure? I have adopted an empirical method which, though not scientifically correct, has working advantages. It is this: For every year of age after 15 allow 1 mm. of mercury, and add 100 to the number. Thus, a person 15 years old, 115 mm.; one of 35 years, 135 mm.; one of 60 years, 160 mm.

Beaulieu.

H. J. JOHNSTON-LAVIS.

CATARRHAL JAUNDICE OCCURRING IN EPIDEMIC FORM.

LAST summer I saw an outbreak of acute catarrhal jaundice among children, occurring in a small village in Suffolk, similar to that reported by Dr. Holdernes in the JOURNAL of December 9th, 1911, p. 1533.

The symptoms were the same in the majority of the 10 cases, namely, jaundice coming on suddenly, with slight rise of temperature and pulse-rate, pale stools, and bile in the urine. Slight pharyngitis was noticeable in each case, while vomiting occurred in only 3. As regards age, the eldest was 12 years old, and the youngest 6. In one family, all sleeping in the same room, only two contracted the disease.

The important point in these cases was the early occurrence of acute pharyngitis, which in every patient was getting better when the jaundice came on, showing,

I consider, the strong possibility of the pharynx being the primary point of infection. I have come across similar cases in the Fens of Cambridgeshire, and always, as in the present instance, in hot, dry summers.

Walsham-le-Willows. RALPH N. POIGNAND, M.B. Cantab.

TREATMENT OF GONOCOCCAL ARTHRITIS.

WHILE agreeing with Dr. J. Bain¹ in his contention that in gonococcal arthritis treatment of the urethra is of the first importance, I would like to supplement his remarks with regard to the best method of carrying it out.

That lavage of the whole urethra is by far the most effective local treatment is certain, but the solution of silver nitrate which he recommends—that is, 10 grains to the pint or about 1 in 1,000—is far too strong to begin with in the majority of cases. The fluid by the irrigation method fills the urethra under considerable pressure and is forced into all its nooks and crannies—lacunae, gland ducts, prostatic ducts, etc.—and while being very effective is also very painful. It is not uncommon to excite an immediate seminal emission, due probably to the fluid finding its way into the ejaculatory ducts. I consider that a strength of 1 in 5,000 (2 grains in the pint) is sufficiently strong to begin with, or even 1 grain in the pint in sensitive individuals. This can be rapidly worked up to 1 in 2,000, or 1 in 1,000 if necessary, in very resistant cases. London water destroys about 1 grain in the pint of silver nitrate, so that unless distilled water be used this should be allowed for in making up the solution. I am also of opinion that irrigation by the hydrostatic pressure method, the reservoir being raised about 5 ft. above the couch, is superior to that by any form of syringe, for the reason that the stronger solutions of silver nitrate are very liable to excite a spasm of the bladder in spite of the use of cocaine. This is evidenced by the fluid rising in the glass reservoir. On removing the nozzle from the meatus the patient will be found to be in the act of emptying his bladder. Again, in the more acute or irritable cases, a violent spasm of the bulbo-cavernosus muscle may occur, the spurting-out fluid forcing the nozzle away from the meatus, and unless the irrigator be held with a light hand an enormous tension in the urethra will ensue. In chronic conditions, no doubt, no evil result follows, but the same cannot be said of the acuter cases. Gonococcal rheumatism very frequently begins when the local disease is in a subacute and not a chronic condition. A large number of these patients have a varying amount of prostatitis, and a strong silver irrigation would undoubtedly be followed by an increase of the prostatic inflammation. A lavage with mild silver nitrate or potassium permanganate solution after a gentle massage of the prostate per rectum would be indicated. To ensure the gentleness of the proceeding $\frac{1}{2}$ drachm of a 2 per cent. eucaine lactate solution may be injected and "milked" back into the posterior urethra as indicated by Dr. Bain.

Great caution must be observed in commencing instrumental treatment in these cases of generalized gonococcal invasion, and they should not be used until the joint or other inflammation has partially subsided. The pressure caused by the use of a sound or a Kollmann's dilator, while securing the absorption of inflammatory deposits, also favours the entrance of living micro-organisms into the circulation. To minimize this evil the whole urethra should be thoroughly flushed with a warm 4 per cent. solution of boric acid or a 1 in 10,000 mercury oxycyanide before the use of the instrument, and the usual silver nitrate or permanganate after.

London, W.

W. WYNDHAM POWELL, F.R.C.S.

NERVOUS RETENTION OF URINE.

A YEAR or two ago it occurred to me that in certain cases of retention of urine—hysterical and after labour—a rectal injection of glycerine might be efficacious, and since then, when such cases have occurred, I have tried the plan with success. I am reminded of it by a district nurse telling me she found it "a great success" after trying it, on my instructions, in a case of retention after labour.

For all I know, the idea may not be new, but I have not heard of it.

HARFORD EDWARDS, L.R.C.P., M.R.C.S.

Markyate, near Dunstable.

¹ BRITISH MEDICAL JOURNAL, December 30th, 1911, p. 1695.