After-History.

The boy was returned to bed with little hope; his colour was ghastly, his breathing feeble and gasping; his pulse was 110, very soft, but much improved by the saline infusion. I did not give him more saline, as I feared pulmonary oedema or exudation. As a last resort I gave him strychnine mij , ether mij , hypodermically. He was devotedly nursed. Strychnine mij , ether mij , was given again in four hours, and half a pint of saline was given per rectum later as often as he would retain it.

Next morning he was a little better, but extremely weak; pulse 108. In view of the fact that I had not been able to sponge out the blood, which must have been contaminated with renal and hepatic exudation, I feared peritonitis, so gave him calomel gr. v, followed by mag. sulph. 3j hourly; an enema at 4 p.m. was without result. He had further saline rectal injections, and in the night strychnine and ether were again given. Next morning the calomel and mag. sulph. were repeated with good result.

On August 29th he seemed better; the bowels were opened again; the general condition of the abdomen was good. The exudation, which had been very copious, was less.

On August 30th he was better; the pulse varied from 86 to 92.

Second Operation.

C.E. was given by Dr. Crofton. I reopened the wound, sponged out sero-sanguineous fluid from the loin, then, with my finger on the vessels, gently removed the clamp; no bleeding occurred. I replugged the liver rent, put a cigarette drain down to the region, and sutured the wound in layers with silk and, for the skin, salmon gut. The peritoneum had retracted far, and was very difficult to bring together.

The general peritoneal cavity had not been shut off by adhesions; it was clean and dry; the extravasated blood and fluid had entirely disappeared; there was no peritonitis.

The patient stood the second operation most satisfactorily. The sinus healed in a few weeks, and the boy left the hospital on October 25th.

I saw him on December 7th. He was strong and well; the scar was perfectly sound.

The following points seem to me very striking:

1. The severity of the injury: (a) The kidney was ruptured completely away from its pelvis and vessels and partially split; (b) the liver was extensively torn, and a portion completely detached.

2. The fact that the renal vessels were only bleeding comparatively slowly, suggesting wonderfully efficient re reaction.

3. The slowness of the pulse at the time of admission and the slowness of its rise. I think that this may be accounted for by rupture of the suprarenal with absorption of its secretion.

4. The capability and activity of the peritoneum in dealing with a considerable quantity of blood and exudation so completely in four days.

ON ACETYL-SALICYLIC ACID, WITH SPECIAL REFERENCE TO ITS VALUE IN TYPHOID FEVER.

By GRAHAM CHAMBERS, B.A., M.B.,

ASSOCIATE PROFESSOR OF CLINICAL MEDICINE, UNIVERSITY OF
TORONTO; PHYSICIAN, TORONTO GENERAL HOSPITAL.

ACETYL-SALICYLIC acid is formed from salicylic acid,

by replacing the hydrogen atom of the hydroxyl group by the radicle acetyl, CH_3CO . The chemical formula of acetyl-salicylic acid is therefore

The aspirin is protected.

Many of the physical and chemical properties of acetyl-salicylic acid are very similar to those of salicylic acid. Like the latter, it is a solid, crystalline, very slightly soluble in water, and freely soluble in ether and alcohol. It also resembles salicylic acid in being a monobasic acid, but differs from the latter in being less stable. For instance, it is decomposed by boiling with an alkali or a mineral acid into a salicylate or salicylic acid. Martindale and Westcott state that heating the drug in presence of moisture decomposes it into salicylic acid, and that the solution in water effected by means of sodium bicarbonate contains sodium salicylate and sodium acetate. They also state that when the drug is administered by the

mouth it passes unchanged through the stomach, decomposing only on reaching the alkaline intestinal juices. Cushny holds a similar view, for he states that the systemic action of the drug is dependent upon its decomposition in the intestine into sodium salicylate. According to these authors, therefore, the actions of salicylic acid and acetyl-salicylic acid, administered by the mouth, are the same, except with regard to their local actions in the stomach. This, I believe, is incorrect; and I shall report observations in this paper which show that the pharmacological action of the two drugs, after absorption, are not the same, and that acetyl-salicylic acid is absorbed, at least in part, before decomposition occurs. Solutions of the drug, which should be similar to those which occur after the drug is administered by the mouth, were tested for salicylic acid, making use of a solution of ferric chloride, which gives a violet colour with salicylic acid, but not with acetyl-salicylic acid. This is the usual qualitative test for a salicylic acid, and it was used, in addition, to determine quantitatively, by a colorimetric method, the amount of salicylic acid or sodium salicylate formed from acetyl-salicylic acid.

Summary of Chemical Experiments.

1. A solution of the drug in cold water gives no violet colour with ferric chloride.
2. A solution of the drug in boiling water gives a violet colour with ferric chloride.
3. If one dissolves the drug in cold water, and then boils, the intensity of the ferric chloride reaction increases for several minutes.
4. The drug, dissolved in water, at the temperature of the body, gradually decomposes, the amount decomposed varying with the length of time for at least twelve hours. A colorimetric method showed that the amount of acetyl-salicylic acid decomposed in twelve hours was four times greater than that in three, and that very little was decomposed in half an hour.
5. The drug dissolves in a solution of baking soda or in a 1 per cent. solution of sodium carbonate without decomposing into a salicylate. If either of these solutions is kept at the temperature of the body decomposition gradually occurs, but very little is decomposed at the end of half an hour.
6. The drug dissolves in a 0.2 per cent. solution of hydrochloric acid without decomposition. If the solution is kept at the temperature of the body decomposition gradually takes place.

These experiments indicated that the decomposition of acetyl-salicylic acid in a medium such as the gastric or intestinal juice is a gradual process, and that thirty to forty minutes after the drug is exhibited, at which time its pharmacological action is frequently evident, very little of the drug is decomposed. I believe that this is true, because I can show that the analgesic and antipyretic actions of acetyl-salicylic acid are more potent than those of salicylic acid, which could not be the case if acetyl-salicylic acid were decomposed by the intestinal juice, as Martindale and Westcott, Cushny, and others state. I should judge, therefore, that as a rule the greater part of acetyl-salicylic acid is absorbed undecomposed.

Pharmacology.

The principal difference in the pharmacological action of the two drugs—salicylic acid and acetyl-salicylic acid—is due probably to the presence in the latter of the radicle acetyl, because the introduction of this group into a compound as a rule augments both the analgesic and antipyretic actions. For instance, acetanilide, which is the acetyl derivative of aniline, and acet-phenetidine (phenacetine), which is the acetyl derivative of phenetidine, are more analgesic and antipyretic in character than aniline and phenetidine respectively.

The Analgesic Action.

Nowadays the analgesic property of acetyl-salicylic acid is fully recognized, and it is extensively used in the relief of pain. It has the advantage over acetanilide, antipyrin, and phenacetine of being less injurious to the blood and other tissues of the body.

The principal indications for the exhibition of acetyl-salicylic acid as an analgesic are the relief of the milder forms of pain, such as that of sore throat, tonsillitis, myalgia, pleurisy, etc. My surgical colleagues tell me that it is very valuable in the relief of pain of diseases of bone and joints, and I have found it, in doses of 5 grains every three hours, of considerable value in the relief of pain of malignant disease.

The Antipyretic Action.

Salicylic acid may be said to act as an antipyretic in two ways, namely, by its action as an internal antiseptic, and by its action on the heat centres, sweat apparatus and other parts which enter into the regulation of the temperature of the body. In acute rheumatism the lowering of the temperature after the exhibition of large doses of salicylic acid is probably partly due to its antiseptic action, because in many cases there is an abrupt fall of temperature to the normal or subnormal, very similar to that of a crisis of an acute infection; usually this is followed by very little rise of temperature after the administration of the drug is discontinued. On the other hand, there are cases of fever in which the administration of salicylic acid produces sweating and a transient fall in the temperature, which, as soon as the drug is discontinued, rises to its former height. The antipyretic action of salicylic acid in this kind of case is no doubt dependent on the action of the drug on the heat-regulating apparatus.

Acetyl-salicylic acid also may be said to act as an antipyretic in two ways, similar to those of salicylic acid; but it is usually not administered in doses sufficient to exercise its internal antiseptic action, and the antipyretic action of the drug must therefore be due to the action of the drug on the heat-regulating apparatus (sweat glands, heat centres, etc.), which property is present to a much greater extent than in the case of salicylic acid.

The following observations afford evidence that acetyl-salicylic acid is a more potent antipyretic than salicylic acid:

To 10 patients with typhoid fever of moderate severity we gave 6 grains of sodium salicylate every four hours, and carefully observed the course of the temperature, etc. The skin did not become moist, and we were unable to detect any result from the administration of the drug. After five days the salicylate was replaced by acetyl-salicylic acid exhibited in 3-grain doses every four hours. The result of the change was almost immediately apparent. The skin became moist, and the temperature was lowered; the degree of sweating was variable. In some cases it was very slight, while in others it was profuse. In cases in which there was little sign of action the dose was increased to 4 or 5 grains.

Acetyl-salicylic Acid in Typhoid Fever.

In the prognosis of typhoid fever the course of the fever is an important consideration. A very high temperature, say above 104° F. every evening, maintained for one or two weeks, and unaccompanied by considerable morning remissions, should be considered a danger signal. On the other hand, a low or comparatively low course suggests a favourable outcome. Indeed, in my experience, the mortality in the latter cases has been practically nil. Again, most physicians agree that in the treatment of typhoid fever it is essential to keep the fever under control, because high fever in itself, irrespective of its cause, is harmful. The question whether cold baths or cold spongings are superior to antipyretic drugs, etc., cannot be discussed here, but in private practice, and sometimes in a hospital, either from want of proper facilities or insufficient aid in nursing, it not infrequently happens that hydrotherapeutic measures prove insufficient to control the fever. The question then arises whether it is better to allow the fever to run its high course or call to use antipyretic drugs. Two years ago I decided in favour of the latter view, because it was found that the exhibition of acetyl-salicylic acid in small doses, 3 to 5 grains every four hours, combined with tepid or hot sponging, was generally an effective means of lowering the temperature. The exhibition of the drug in this dosage to typhoid patients does not alter the blood pressure or produce any appreciable ill effects. The skin remains moist, and in some cases there is profuse sweating. The greatest effect on the temperature is obtained by sponging the patient about half an hour after the administration of a dose of the drug, in this way combining the antipyretic actions of the medicinal and hydrotherapeutic agents. It is probable that other beneficial effects are derived from the exhibition of acetyl-salicylic acid. The increase in the amount of perspiration should diminish the toxæmia, although it is very difficult to estimate how much endotoxin is excreted with the sweat. As a clinical observation I may mention that the patients were all of the opinion that they felt better during

a moderate degree of perspiration, and to me they seemed clearer mentally and more willing to carry out other parts of the treatment, such as the drinking of water, etc.

I have been especially impressed with the marked diaphoretic and antipyretic actions of the drug. Three grains every four hours had frequently considerable effect, and in most cases 5 grains at the same periods produced profuse diaphoresis. In only one case were we compelled to increase the dose above 5 grains. In two cases—one treated during September, 1909, and the other during October, 1910—5 grains every four hours appeared almost too active, as the temperature charts show a fall in temperature of 9° and 10° respectively during the exhibition of the drug; there was no untoward effect during the fall of the temperature, but in the case of antipyretics which act as acetyl-salicylic acid does, one does not like to see such precipitous action. In giving acetyl-salicylic acid to typhoid patients, it is desirable, therefore, to commence with a dosage of 3 grains every four hours; if this should not be sufficient, the quantity should be gradually increased until the desired effect has been obtained. The reason why acetyl-salicylic acid is more effective as an antipyretic in typhoid fever than in most other fevers is, probably, that the fever of typhoid fever is frequently labile in character. This theory also affords an explanation of the undoubted value of hydrotherapy in the treatment of the disease, because it is found that the more readily the temperature can be reduced by a bath or sponge the better the prognosis.

The Mode of Administration.

When a drug is given dissolved in water, its action is usually quickly manifest, because it almost immediately passes into the intestine and is absorbed. This principle is of special importance to the exhibition of acetyl-salicylic acid, because the shorter the stay in the gastrointestinal tract the less the decomposition of the drug. The drug should therefore never be administered in tablet form, but in solution. It is only slightly soluble in cold water; it is freely soluble in water containing sodium bicarbonate; the application of heat to a solution in water tends to decompose the drug.

PASSAGE OF A CAST OF THE BLADDER PER URETHRAM.

BY

JEHAN M. BARLET, M.D.BRUX., M.R.C.S., L.R.C.P.

THE patient in the following case, a married woman aged 26, first sent for me on November 2nd, 1911, and stated that for two or three days she had had severe pains in the lower part of the abdomen, which were like labour pains; she had also been losing large quantities of blood, with clots, and had passed a large mass, which she said was like a piece of flesh.

State on Examination.

Bimanually, the uterus was found to be enlarged to about the size of a three months pregnancy; the os was patulous, and admitted the tip of the index finger, and she was losing freely.

Treatment.

She stated that her periods had been perfectly regular, but thinking she had had a partial miscarriage, and that the placenta had been retained, I dilated the cervix with Hegar's dilators to No. 16, and I then inserted my finger into the uterus, and, to my surprise, I came upon a fetus enclosed in the membranes. This being the case, I did not rupture the membranes, but after washing out the vagina with cyllin, I plugged the vagina with iodoform gauze. The latter I removed the next day.

Result.

The patient went on exceedingly well; all the pains and haemorrhage ceased. No miscarriage took place, and on November 12th she got up perfectly well.