CONCLUSIONS.

1. The condition found after the non-union of a cut nerve trunk is very variable.

2. These varying conditions render any definite statement as to time limit to secondary suture difficult, if not impossible. I have had successful cases of secondary nerve suture in the human subject after many (six) years of non-union.

3. Where regeneration after secondary nerve suture has not taken place, the main cause (apart from local conditions at the suture area, such as sepsis) is lack of regenerating power in the nerve cell.

4. When all factors have been taken into consideration and it is concluded that restoration of function after secondary nerve suture is unlikely, there is still another surgical procedure open. This is the suture of recently divided central nerve fibres to the distal trunk of the degenerated nerve—that is, by nerve anastomosis to an adjacent healthy nerve.

My best thanks are due to Professor Osborne for many suggestions, and to Mr. Jona, who assisted me at most of the experiments.

The expense of this research was partly defrayed by a grant from the Royal Society.

REFERENCES.

¹ Bowlby, *Lancet*, 1902, p. 197. ² Sherren, *Injuries of Nerves*. For best results in anastomosis see paper by writer in *BRITISH MEDICAL JOURNAL*, September, 1905

PNEUMOCOCCAL MENINGITIS.

BY

HORSLEY DRUMMOND, M.D.,

SENIOR ASSISTANT PHYSICIAN TO THE ROYAL VICTORIA INFIRMARY,
NEWCASTLE-UPON-TYNE.

THE causes of acute inflammation, with purulent lymph exudation, of the pia arachnoid are various forms of pus-producing micro-organisms, and not infrequently more than one kind of organism is found in the same case. The *Meningococcus intracellularis* of Weichselbaum usually suggests itself in the first instance, especially when cases of cerebro-spinal meningitis are known to be in the neighbourhood, though the claims of the pneumococcus for recognition as a common cause are well known to pathologists. The impression that this organism is a more common cause of acute meningitis than is generally supposed, and that involvement of the cerebral membranes is a more frequent complication of fatal cases of pneumonia than is commonly believed, must be my excuse for publishing the following cases. For the notes of the third case I am indebted to Dr. J. W. Smith of Ryton, who has kindly given me permission to use them:

CASE I.

James S., aged 15, a factory storekeeper, was in good health until February 1st, 1911, on which day he was struck on the head by a heavy iron bolt which fell 12 ft. He was not rendered unconscious, but it produced giddiness and headache. He was able to attend to his work next day, and little was thought of the injury until three days later, when he fell in a fit whilst at work. He was unconscious for upwards of two hours, and on recovering complained of pain in the right side of the chest and severe headache.

He was admitted into the Royal Victoria Infirmary on February 5th, when it was noticed that he was drowsy, had a temperature of 101°, a pulse of 95, and that his face twitched. Next day he was restless and delirious and complained constantly of severe headache, chiefly frontal. There was photophobia and slight nystagmoid movements and the pupils were dilated, but there was no ocular paresis. There was slight rigidity of the muscles of the neck, but no retraction of the head. The skin of the trunk and legs was hyperaesthetic. The knee-jerk was normal and the plantar reflex flexor. There was no optic neuritis.

On the third day of the illness the breathing became accelerated and a hyper-resonant note was detected over the left lung. This was, however, soon replaced by dullness, and harsh breathing by tubular, so that it was evident by the fourth day at least that the illness was pneumonia complicated by nervous symptoms. Lumbar puncture on the third day led to the withdrawal of clear fluid free of organisms at increased pressure.

There was little change in the symptoms until the sixth day, when the respirations and temperature rose, the former to 62, and the latter to nearly 104°. Indeed, for a day or two the signs of meningitis became less evident, and the case assumed in a more pronounced form the aspect of a very bad case of croupous pneumonia. On the seventh day the nervous symptoms

resembled the delirium of enteric at the end of the third week: the patient was wakeful, muttered constantly to himself, and occasionally uttered loud shouts. He rapidly sank and died on the eighth day.

The urine was free from albumen throughout the illness, and there was no vomiting.

Post-mortem Examination.

Body poorly nourished. Rigor mortis general. No signs of injury to scalp or skull. The heart showed nothing abnormal worthy of note.

The pleural cavities contained small quantities of turbid fluid. Evidence of recent pleurisy over surface of left lung, both lobes of which were consolidated. The right lung was intensely congested, and the lower lobe contained patches of bronchopneumonia.

The spleen was soft and acutely congested. The liver and kidneys were in an advanced state of cloudy swelling.

Examination of the brain showed general leptomeningitis, best marked over the vertex, where there was a deposit of greenish fibrinous lymph. The exudate was also well marked over the base, especially on the under surface of the frontal and temporal lobes and over the pons, medulla, and cerebellum. Films from the cerebro-spinal fluid showed typical pneumococci.

CASE II.

James B., aged 38, cartman, was admitted into the Royal Victoria Infirmary in a semi-conscious condition on January 27th, 1911.

The history of the case was very indefinite. The friend who brought him to hospital stated that he had been found in a lodging-house that morning in a state of delirium, and had only been ill a few hours, having fallen in a fit, which was apparently the commencement of the illness.

The cheeks were flushed and the pulse and respirations quick. There were temporary intervals of consciousness, when his speech was slurred and slow and he appeared to have some difficulty in articulating. The temperature was 103.4°, the pulse 114, and the respirations 28. There was no paralysis. The tongue was furred and dry. The patient was restless and delirious and was with difficulty kept in bed. There was nothing definite to be detected in the chest. The reflexes were normal.

On the morning following his admission he became very violent and excited, and this was followed by a period of collapse, in which the pulse became weak and rapid (130), and a few hours later paresis of the right side of the face and right arm and left eye were noticed. The temperature was 103°, and the respirations 44.

As the day advanced he picked at the bedclothes and became more and more drowsy. The urine had to be drawn off, and was found to contain albumen and sugar. Then ptosis of the left eyelid appeared and the paralysis extended to the right leg with anaesthesia, on which the plantar reflex was extensor. The next development was paralysis of the left side of the face, motor and sensory.

A lumbar puncture was made, and turbid fluid flowed under marked pressure. It contained flakes of lymph, was highly albuminous, and did not reduce Fehling's solution. Pneumococci, streptococci, and lymphocytes were in abundance. A blood count showed no leucocytosis. The optic discs were normal.

He passed rapidly into a state of profound coma, and died on the morning of the fourth day.

Post-mortem Examination.

The body was well nourished. Rigor mortis present. Post-mortem lividity marked.

Brain.—Extensive suppurative leptomeningitis. The exudate extended to the vertex and over the base (temporal lobes, pons, and cerebellum). It was best marked over the right hemisphere. It passed down the spinal canal and affected chiefly the cervical and dorsal regions (back and front). A considerable quantity of turbid fluid escaped from the base of the brain and spinal canal. Stained films showed numbers of pneumococci.

Lungs.—Acute inflammatory changes in bronchi and extensive patches of bronchopneumonia on both sides.

Heart.—Muscle soft and flabby. Recent vegetations on both segments of the mitral valve indicating infective endocarditis.

There was cloudy swelling of the liver and kidneys and the spleen was acutely congested.

CASE III.

J. T., a man aged 42, was taken suddenly ill on January 23rd, 1911, with intense headache, vomiting, aching in the back and limbs, and general febrile symptoms (temperature 103°, pulse 132). On the third day delirium, and on the fourth twitchings of the legs and arms, with rigidity of the muscles of the back of the neck and coma were observed. Next day there was general rigidity with opisthotonos; the knee reflex was active and the plantar response extensor, and the pupils were widely dilated. He died on the sixth day of the illness.

Post-mortem Examination.

The lungs were congested and oedematous at the bases, but there was no pronounced consolidation. The heart was normal.

The soft membranes of the brain were covered by a greenish purulent exudate, which passed down the spinal canal. A bacteriological examination of the exudate demonstrated pneumococci.

There is abundant evidence in medical literature of the connexion between pneumonia and meningitis, dating from the publication by Immermann and Heller in 1868 of 9 cases of purulent meningitis found in a series of thirty pneumonic sections, and yet it cannot be said that the subject has received the attention it merits.

It would appear that our present knowledge does not justify any positive statement with regard to the frequency with which meningitis complicates pneumonia, as the statistics of writers vary within wide limits. Netter, for example, quoted by a writer in the *Lyon Médical* (1909), reports numerous statistics varying from 43 per cent. to 1 per cent. Beddard¹ suggests 1 per cent. above the age of 10 years, and from 1 per cent. to 2 per cent. below that age. Holt² met with meningitis four times in 500 cases. Osler³ gives 8 per cent. of fatal cases.

But it is manifest that these findings cannot be relied upon, and I suspect that when the question has been gone into more carefully in the light of more extended statistics and improved methods of diagnosis, and latent cases are recognized, it will be found that meningitis, especially in children, is more common than is generally supposed.

An important paper by Liebermeister⁴ on the frequency of meningitis with pneumonia makes the fact clear that the meningitis may easily be overlooked. In three out of eleven consecutive fatal cases of pneumonia occurring in adults ranging from 45 to 71 years, he found reliable microscopic evidence in the brains and spinal cords of acute meningitis, without any gross naked-eye appearances suggestive of membrane involvement; in none of these did the clinical symptoms and signs justify the diagnosis of meningitis. This author alludes further to the connexion between pneumococcal meningitis and outbreaks of cerebro-spinal meningitis, an association to which Immermann and Heller had already called attention, and which has not escaped the notice of other writers. Additional evidence of the frequency of meningitis in cases of acute pneumonia is to be found in the record by Hugo Meyer (to which Liebermeister refers) of 5 cases of latent purulent meningitis in 11 pneumonic necropsies.

The importance of Liebermeister's observation that the meningeal lesion was often microscopic is corroborated by the records contained in a paper on meningitis and meningo-encephalitis by Ludwig Kirchheim,⁵ who refers to 500 cases of pneumonia, all in young people; in 13 there was, in his opinion, clinical proof of meningitis, but ending in recovery; and in 4 only were there signs of purulent meningitis. In the case of the thirteen he was apparently content with the diagnosis of meningeal irritation (*meningiale Reizung*).

The opinion obtains very widely that cases of pneumonic meningitis are nearly always fatal, and probably very few would venture upon a favourable prognosis. Against this opinion may be set Kirchheim's contention that the milder cases recover; and without any desire to minimize the risk or to call in question the correctness of the common belief that the prognosis is grave, I may quote the following case recorded by Dr. Hemenway⁶ as affording some ground for the more favourable view.

A child aged 2 was admitted to the "Babies' Hospital" New York, with complete consolidation of the left lower lobe, but without cerebral symptoms. On the twelfth day of the illness twitchings were noticed with some retraction of the head. Lumbar puncture was performed and 1½ oz. of clear fluid at high pressure were withdrawn. There was no deposit, but cultures of the fluid showed an abundant growth of pneumococci. The signs of meningitis continued, but on the fifteenth day of the illness the temperature suddenly fell, and the infant rapidly improved and left the hospital cured after an illness lasting twenty-four days.

I venture to think it would not be difficult to procure further evidence in this direction, for the contention that meningitis is not really a rare phase of pneumonia would seem to warrant the deduction that the milder cases recover, but this naturally raises the question how far we are justified in including cases without naked-eye pathological changes in the category.

REFERENCES.

- ¹ Beddard, *Allbutt and Rolleston's System of Medicine*, vol. v, p. 229.
² Holt, *Arch. of Ped.*, 1893, vol. x, p. 1112. ³ Osler, *Modern Medicine*, vol. ii. ⁴ Liebermeister, *Muench. med. Woch.*, 1909, p. 750. ⁵ Ludwig Kirchheim, *Med. Klin.*, 1908, p. 1461. ⁶ Hemenway, *Arch. of Ped.*, 1906, vol. xxiii, p. 276.

TRANSIENT HEMIPLEGIA FOLLOWING PARTURITION.

By HUGH GILLIES, M.R.C.P. EDIN.,
NEW ABBEY, DUMFRIES.

THE following cases of hemiplegia following confinement occurred with such a short interval between them that they could not fail to arouse attention, and a desire for an explanation of the very interesting phenomena they presented.

CASE I.

Mrs. M. H., aged 22, was confined of her second child on June 6th. She had "fitted" on the preceding day from a farm distant thirty miles from her new home; the journey had been done perched upon a quantity of household furniture on an open cart. About 8 p.m. on June 5th she felt her first pain, and I was sent for at 4 a.m. I found the head on the perineum, and the child was born in a few minutes; the placenta was expelled ten minutes afterwards. The whole confinement was perfectly natural; the child was a poor thing, at about the seventh month. The course of after-events was normal. I paid my last visit to the mother on June 14th, and allowed her up the next day.

On July 2nd, nearly a month after the birth, while sitting at the fireside nursing the child, the mother "was seized with a fit, her face was working, and she was all twisted to one side, her eyes were rolling in her head." I found her in bed; the face was pale and covered with a cold perspiration, the breathing was stertorous; the head, face, and eyes were drawn to the right side (conjugate deviation); the left arm and leg were flaccid, and when lifted dropped on to the sheet heavily. The pupils were dilated, and the conjunctival reflex was not quite abolished, though lazy. The sounds in the heart areas were normal, and there was no sign of any valvular trouble. The pulse was 78, the vessel wall soft, and the wave full and regular.

Next day she was completely conscious, and four days after the power had to a great extent returned in the affected parts, the hand being the last part to recover. A month after, the recovery was perfect in every part, with the exception of the thumb, which is to-day weak and shrunken.

On inquiring into the history after her recovery, the only thing of any importance was a story of chorea when she was 9, otherwise there was no neurotic history. There was no albumen in the urine either then or at the confinement.

CASE II.

J. W., aged 25; her second child was born on August 1st; the confinement perfectly normal, and the puerperium uneventful.

On September 27th, while sitting by the fire with her baby, she fell off the stool, dropping the baby beside her on the floor; in this position she was found by a neighbour, who procured assistance and had her lifted into bed. She never lost consciousness, but remembered all that happened quite well. Paralysis affected the left arm and leg, the face and eyes being quite normal. She complained of most intense headache, which seemed to be of maximum intensity over the right side and above the right ear. The heart impulse was in the fifth interspace and powerful; a murmur was heard, of greatest intensity in the aortic area and systolic in rhythm; the pulse was 90.

The paralysis of the limbs lasted for three days, and then quickly recovered. She is now quite well. In this case also there was a history of chorea during girlhood, but nothing else of any importance.

Here we have two very interesting cases of something producing a sudden and powerful temporary effect upon the brain, and almost as quickly being withdrawn, with complete restoration of function to the parts involved; and it may be permitted to indicate briefly the probable causation of the attacks. In Case I, though the patient made a complete picture of a cerebral haemorrhage, yet there was an indefinable something which made me hesitate to give a too grave prognosis—it may have been the symptoms of haemorrhage into the brain with a tranquil pulse; it is not easy to say. After the first case the second was much easier to define.

That it was neither an apoplexy nor an embolism is proved by the almost complete restoration of function in either case, and, though local palsies and unconsciousness are not uncommon in uraemia, yet the known good health of the patients, the absence of albumen, and of all other signs of kidney trouble, make it necessary to exclude it also.

It seems to me that both cases were caused by a spasmodic contraction of the cerebral arteries—an angiospasm—such as is known to occur in migraine (in which case the retinal arteries on ophthalmoscopic examination can be seen as thin threads). The occurrence of previous attacks of chorea in both cases adds to the soundness of the