Time and carefully kept records alone will enable a judgement to be formed of the permanency of such results.

I look forward to the time when the resistance of the units of the nations has been so raised that it will be only exceptionally that the tubercle bacillus will be able to live in human tissues.

REFERENCE.

¹ *The Fight against Tuberculosis and the Death-rate from Phthisis.* By Karl Pearson. London: Dulau and Co.

NOTES ON CLINICAL TRIALS WITH MARMOREK'S ANTITUBERCULOUS SERUM.

BY

JOHN P. MITCHELL, M.D.,

HOUSE-PHYSICIAN, SEAMEN'S HOSPITAL, GREENWICH.

RECENT discussions on the value of tuberculin have shown that much diversity of opinion exists as to its efficacy. My experience with tuberculin has been unsatisfactory, but I have obtained rather more success with another agent—namely, Marmorek's antituberculous serum. This serum has been before the profession for a number of years, but no definite opinion of its utility has been formed in this country though many foreign observers have spoken very highly of it.

I here record the treatment and results of 10 cases which were treated by me in the Royal Waterloo Hospital, London, S.E. For allowing me facilities for this purpose I have to thank Dr. Guthrie Rankin.

Two methods of injection are used—hypodermic and rectal. The former is supposed to be beneficial in acute and rapidly advancing cases, while the latter is more efficacious in chronic and apyretic cases. The great drawback to the subcutaneous injection is supersensitiveness, which is common to the injection of all serums.

The dose used was 2½ c.cm., given hypodermically, until anaphylaxis was seen, when 5 c.cm. was given rectally daily for fourteen to twenty-one days. This was followed by an interval of seven to ten days without serum, which was again followed by daily injections for fourteen to twenty-one days. This routine was carried on as long as the case demanded it. Marmorek recommends that even after recovery seems complete weekly doses of 5 c.cm. should be given for several months.

CASE I.—W. S., aged 3, admitted January 9th with great dyspnoea, a harsh, dry cough, almost incessant sunken features, wasted and in an advanced stage of disease, seemed in some measure improved after four injections. By February 8th there were signs of cavitation at the right apex, and death took place on February 24th. Serum had been given daily.

Post mortem: The right pleura was greatly thickened and covered with tuberculous deposit. At the right apex a large cavity with extensive infiltration throughout the lobe was found. The left lower lobe was consolidated, and the bronchial and mesenteric glands were tuberculous.

It was notable that, in spite of the steady progress of the disease, the temperature and pulse were steadied, the symptoms were less severe, and the patient was more comfortable.

CASE II.—I. H., aged 3, was admitted December 21st with history of having recently had pneumonia, while phthisis was rampant in his family. The entire right lung was affected. The usual treatment was adopted, but the disease made rapid progress, and when serum treatment was begun on January 19th the child was almost moribund, with a temperature swinging between 101° to 104° F. By February 12th the child was eating and sleeping well, cough and dyspnoea had disappeared, and physical signs were clearing up, and on March 4th she was sent home much improved, with normal temperature and pulse. Seen a month later, improvement was well maintained, and there had been great increase in weight.

CASE III.—L. D., aged 10, admitted February 25th with tuberculous history and well-defined physical signs throughout the right lung. Serum was commenced on March 1st and continued up to April 4th, when her condition was so much bettered that she was sent to a convalescent home. The physical signs had not altered, but the temperature was normal and the weight steadily increasing. Reported on two months later the child was at school and thriving well.

CASE IV.—S. B., aged 7, admitted February 3rd, complained of rapid wasting, night sweats, cough, and dyspnoea, and was found to have definite signs in the right lower lobe, while bacilli were found in great numbers in sputum. Serum was given from February 8th to March 5th, and he was discharged on March 26th with practically no physical signs in the chest, greatly increased in weight, no cough, and no night sweats. A report on this child could not be obtained two months later.

CASE V.—G. G., aged 3, admitted February 23rd, in a very much emaciated condition, with extensive physical signs all over the left chest. Serum commenced on admission and continued up to March 26th without effect. Death took place on April 3rd, and *post-mortem* examination showed typical miliary tuberculosis.

CASE VI.—A girl, aged 6, admitted January 6th with a large indurated mass of submaxillary and cervical glands, which, in spite of antituberculous treatment for over eighteen months, had persisted. The general condition was good and the lungs were healthy. Serum was commenced on February 2nd and continued up to March 26th, when she was discharged. During this time the glands, which were about as large as walnuts, hard and adherent, diminished to the size of small almonds, and became simultaneously soft and discrete.

CASE VII.—S. E., a boy, aged 7, was admitted on January 7th with swelling around the ankle and a sinus on the external aspect of the os calcis, which had been discharging for about a month. The sinus was opened up and cleaned on admission, but when serum was commenced on January 28th had not ceased to discharge. Injections were continued up to April 2nd, when, after much coaxing, the sinus cleaned up and healed, and good movement in the joint was obtained. One cannot think that in this case the serum had much curative effect, as with the usual surgical treatment a good result would have been anticipated as soon.

CASE VIII.—S. W., aged 2½ years. Admitted February 27th with swelling in right knee. Skiagram showed definite rarefied patch in the diaphysis of the lower end of the femur, with what appeared to be a sequestrum in it, while the joint capsule was much thickened. Serum injection was commenced on admission, and continued up to April 3rd, when the patient contracted measles and had to be discharged from hospital. Before discharge a skiagram of the bone showed very slight improvement, but clinically there was no change.

CASES IX AND X.—Two girls suffering from tuberculous hip disease, with considerable deformity, and, as shown by skiagrams, considerable destruction of the heads of the femora and acetabular cavities. Serum in each case was given for about two months and a half, but recovery was not more rapid than would have been expected with strict surgical treatment alone.

At the Seamen's Hospital, Greenwich, I continued the treatment of phthisis with Marmorek's serum. Eight cases were treated, all in an advanced stage of disease, and, with the exception of one case, no improvement was noted. In this case the patient was a lad of strong physique, in whom weight and chest measurements rapidly increased, and in whom the fever and average weekly weight of sputum diminished correspondingly.

At the time of discharge, however, the sputum was still muco-purulent, and contained bacilli, and the physical signs had not much altered.

It is claimed for this serum that under its influence tuberculous sinuses of old standing rapidly cease to discharge and heal. But in those cases selected for trial this was not found to be so. It is, however, but fair to state that, besides having these discharging wounds, the three cases were in an advanced stage of pulmonary phthisis, and, although treatment extended over three months, no change took place in the amount of discharge from, in the first case, an ischio-rectal sinus; in the second, an old tuberculous empyema wound; in the third, a submental sinus.

Anaphylaxis was a troublesome feature in the administration of the serum by the hypodermic method, and the rectal method was chiefly employed.

The dose for adults was 5 c.cm. for hypodermic and 10 c.cm. in saline solution to 3ij for rectal injection. The frequent rectal injection was much resented by the patients, and this was a decided objection to the treatment.

I cannot claim on these results the success which some Continental writers have claimed. But, in my opinion, the following conclusions are warranted:

1. There is no danger in administration apart from anaphylaxis, which can be overcome by adopting the rectal method.

2. In certain cases the general improvement is most striking, although physical signs are not altered to a similar extent.

3. The undoubted success obtained in some of the cases justifies further clinical trials.

THE next examination of candidates for the Royal Naval Medical Service will be held in London on April 1st and following days. Fifteen appointments will be offered for competition. Application forms, which must be returned by March 21st, and the regulations for entry can be obtained from the Medical Director-General, Admiralty, S.W.