autoinoculation test we possess the means of diagnosing tuberculosis in its earlier stages, of the further fact that this test is so extraordinarily reliable, that on its verdict alone treatment may be confidently applied or as confidently omitted, and, lastly, of its utility in directing the course of our treatment and of deciding the moment when cure is complete.

REFERENCES.
¹ Wright and Reid, *Proc. Roy. Soc., B.*, vol. lxxvii, 1905. ² Wright, Herter Lectures, *Lancet*, August 17th and 24th, 1907. ³ Studies in Connexion with Therapeutic Immunization, Wright, Douglas, Freeman, Wells, and Fleming, *Lancet*, November 2nd, 1907. ⁴ Patterson and Inman, *Lancet*, January 25th, 1908. ⁵ Inman, *Lancet*, December 17th, 1910.

THE PATHS OF RHEUMATIC INFECTION AND THEIR PROTECTION IN CHILDREN.

BY

J. ROSS MACKENZIE, M.D. ABERD.,

ABERTILLERY, MON.

THE admirable work of Drs. Poynton and Paine has clearly demonstrated that a micrococcus, possessing distinct morphological characters, can be isolated from the tissues of the rheumatic patient, cultured on nutrient media, inoculated into animals, producing anatomical lesions identical with those found in acute rheumatism; and further, that this micrococcus can be isolated in pure culture from the blood and tissues of these infected animals, completing the essentials of bacteriological science. The researches of Drs. Poynton and Paine,¹ Beattie,² Ainley Walker,³ D. B. Lees,⁴ Apert,⁵ Wassermann and Malkoff,⁶ Dana,⁷ and others have done much to dissipate the scepticism manifested as to the identity of the *Micrococcus rheumaticus*. The paths by which the specific agent finds access to the system as well as the prevention of such invasion still demand attention.

I. THROAT.

(a) Local Invasion.

The most important, as well as the most frequent, path of infection is undoubtedly the throat. It is occasionally difficult, however, to decide that diffuse congestion of the tonsils and pharynx have any connexion with the rheumatic agent, even when the manifestations of rheumatic toxæmia are more or less general. A relationship is at once suggested by the large percentage of children suffering from rheumatic phenomena who, at the same time, present enlarged tonsils and hypertrophied tissue in the naso-pharynx. The contention that such a condition of the throat increases the susceptibility to rheumatic infection is rational, but enlarged tonsils never produce rheumatism until they become infected with the specific agent, and even then the infective agent may remain localized in the tonsils or pharynx, producing toxins which are projected into the system in variable quantities and at variable intervals. This, I think, accounts for my frequent failures to find the *Micrococcus rheumaticus* in the blood, and also for the erratic behaviour and latency of rheumatic manifestations in children.

There is evidence accumulating which goes to show, further, that in the absence of hypertrophied adenoid tissue in the throat or naso-pharynx there is frequently inflammatory congestion in the region of the soft palate, pillars of the fauces, and pharynx (Holt,⁸ Hutchison,⁹ and Cheadle¹⁰) in the earliest stage of rheumatic infection, and from such a rheumatic throat the micrococcus has been isolated by Poynton.¹

(b) General Infection.

It has been suggested that the infective agent finds entrance from the throat to the general system through the lymphatics. In many cases the lymphatic glands at the angle of the jaw are enlarged, indicating infection of the lymph channels, due, however, not to the absorption of rheumatic infection through the lymphatics, but to enlarged and infected tonsils. In the purely rheumatic throat the lymphatic glands are not enlarged. This fact, together with the evidence that the *M. rheumaticus* has an affinity for the endothelial cells lining the capillary blood vessels, and the sudden onset of acute rheumatic phenomena when the infective agent comes in contact with an injured surface, force me to the conclusion that the path-

from local invasion to general infection is direct into the blood stream and not through the lymph channels.

CASE I.

A girl, aged 9 years, previous to the operation of tonsillotomy was in excellent health, presenting only the usual signs of enlarged tonsils and adenoids. Within a few hours of the operation she complained of pains in the knees and ankles. The temperature was then 99.5° F. and the pulse 100. Next day the temperature was 101° F., pulse 120, and there was marked pain and tenderness with effusion into the knee-joints and pain in the elbow-joints. The symptoms subsided after some days under sodium salicylate, but not before cardiac dilatation had supervened. This case may seem a pure coincidence, but I have had similar experience on several occasions, and the interpretation of this sudden onset of rheumatic phenomena is, I consider, the disturbance of a local and latent focus with rapid absorption of the micrococcus into the circulation from the bleeding surface of the tonsillar bed or the pharynx. It is interesting to note (1) that a swab taken from the throat and smeared upon agar, blood serum, and gelatine, gave on each a culture of staphylococci; (2) that a particle taken with aseptic precautions from the centre of the left tonsil after enucleation gave in broth a culture of staphylococci with numerous streptococci in short chains.

To emphasize this point I record here four consecutive cases. They are important inasmuch as none of them, apart from occasional attacks of sore throat and growing pains, exhibited other than the concomitant signs of hypertrophied tonsils and adenoids. The tonsils, on removal, were put in sterilized bottles. With strict precautions a particle from the centre of each was inoculated in broth. After thirty-six hours' incubation films were made, and stained with methyl thionine. A more or less pure culture of streptococci in short chains was found in each case, while on some films were seen scattered nodules of lymphoid tissue with numerous cocci arranged in pairs. The microscopical and morphological characters of each culture were compared with those of the original *M. rheumaticus* described by Drs. Poynton and Paine, and were found to be identical.

Age.	Previous History of Rheumatism.	Remarks.	Bacteriological Result.
8 yrs.	Negative	Frequent sore throat	Pure culture of cocci in short chains with diplococci.
6 yrs.	Mild attack of arthritis six months previously	Occasional sore throat and growing pains	Some staphylococci with numerous streptococci in short chains.
9 yrs.	Growing pains	Sore throat occasionally. Some deafness from recent otitis media	Pure culture of cocci in short chains with numerous diplococci.
10 yrs.	Negative	Frequent mild attacks of tonsillitis	Streptococci in short chains, diplococci and staphylococci.

II. PULMONARY MUCOUS MEMBRANE.

It will not be seriously disputed that many cases of acute and subacute rheumatism in children show no evidence of sore throat, and of late my attention has been directed to the possibility not only of certain mild catarrhal conditions of the bronchial mucous membrane owing their inception to infection with the *Micrococcus rheumaticus*, but also to a general infection taking place through the impaired mucous membrane.

(a) Local Invasion.

Satisfactory experimental evidence of the local invasion of the pulmonary mucous membrane by the *Micrococcus rheumaticus* in the form of bronchial catarrh or bronchopneumonia is wanting. If a mild bronchial catarrh may proceed to actual bronchitis, and bronchitis to capillary bronchitis and ultimately consolidation of the alveoli, it is difficult to understand why rheumatic bronchopneumonia may not arise in a similar way, and clinically I am persuaded that it does.

CASE II.

A girl, aged 10, was brought to me on account of a persistent cough with occasional muscular and articular pains. There was a history of acute rheumatism and a well-marked family history. No indication of sweating or wasting. The heart was normal, and there was no evidence that this was a throat cough. The only physical sign found in the chest was a harsh respiratory murmur more marked on the left side, and especially between the scapulae and below the angle of the left

scapula and below the left clavicle in front. The pulse and temperature were normal. I could not find any indication either of enlarged bronchial glands, or ear trouble, or gastrointestinal disturbance, and the mother assured me there was no expectoration. A provisional diagnosis of bronchial catarrh was made; after some days the child had a temperature of 99.2° F., with a dry, hacking cough and indefinite pains. Under sodium salicylate the whole condition, including the cough, disappeared within forty-eight hours.

(b) General Infection.

During a recent epidemic of measles I observed that some children with an ordinary attack had a rather prolonged convalescence. In several cases noted there were varied combinations of mild polyarthritis, cardiac dilatation, anaemia, and erythema nodosum, with slight elevation of temperature and some bronchial catarrh accompanied by cough. There was no trace of throat invasion in any of these cases.

CASE III.

I attended two children suffering from measles, sisters, aged 4 and 6 years respectively. They had a rheumatic diathesis and a strong maternal family history of acute rheumatism. So far as could be made out by examination the throat remained healthy in both children, but there was some bronchial catarrh. On the fifth day both seemed well, with normal temperature and pulse, though the younger presented some twitching movements of the hands and both had an irritating cough. On the sixth day definite choreic movements were noted in the younger child, upon which supervened cardiac dilatation with a mitral systolic murmur, and, after some weeks, fatal pericarditis. On the ninth day the elder child had pain and swelling of the knees and ankles, temperature 100.5° F., with spots of erythema nodosum, from which she made a complete recovery.

Quite recently I had two cases of well-marked erythema nodosum in children suffering from pertussis. In neither case was there any possibility of the symptoms being due to alimentary disturbance, for both at the time were upon invalid diet, and there was neither constipation nor diarrhoea. Further, the only alteration in treatment was the addition of sodium salicylate. I am of opinion that the path of infection in these cases is through the impaired bronchial mucous membrane, and this conclusion has been confirmed occasionally in my experience of bronchopneumonia, for I can recall more than one case accompanied by polyarthritis, peliosis rheumatica, and cardiac dilatation.

III. INTESTINAL WALL.

(a) Local Invasion.

Manifestations of rheumatism frequently arise in children who have neither rheumatic throat nor bronchial catarrh, but who show disorders of the intestinal canal, usually in the form of enteritis or mucous colitis.

It is an everyday incident to see a child between 5 and 10 years brought because of wasting and loss of appetite. There is a history of previous rheumatism, the tongue is dry and furred, breath foul, bowels confined, and faeces enveloped in mucus, or there is diarrhoea alternating with constipation, night terrors and perhaps pain in the upper and left half of the abdomen, or, and much more frequently, muscular or articular pains in the legs. I have treated many such cases strictly on the lines of mucous colitis, with only partial and temporary improvement until anti-rheumatic remedies were exhibited. From such clinical evidence, together with the therapeutic test, it would be difficult to dismiss the possibility of this condition, occasionally representing, in some degree at least, a manifestation of rheumatism, and the primary condition of the bowel would certainly foster the *Micrococcus rheumaticus*.

It is recognized that rheumatic pleurisy occurs, but the theory of the rheumatic origin of appendicitis and peritonitis has been received with suspicion; yet such cases as the following are not, I believe, very rare:

CASE IV.

A boy, aged 9 years, had severe pain over the region of the appendix, a temperature of 101° F., pulse 105, and definite resistance and tenderness around McBurney's point. On the third day of illness there was acute pain in the right knee and ankle joints. Two days later there was arthritis, with effusion into the right elbow and left wrist joints. On the tenth day a definite mitral murmur was heard at the apex, accompanied by dilatation of the left ventricle, with profound anaemia. The family history was positive; a younger sister had had two attacks of acute rheumatism, and the whole condition, with the exception of the mitral murmur, which still persists (nine months after), cleared up under the influence of salicylates and local applications.

C

A similar case in a young boy has been described by Dr. Eustace Smith;¹¹ that I now record is one of several cases of which I have had experience with abdominal pain and tenderness over the appendix, accompanied by articular pains and pyrexia, all of which quickly cleared up with antirheumatic remedies.

Whether mucous colitis and enteritis are ever actually produced by the rheumatic agent I am not prepared to say, but I am of opinion that they are conditions which predispose to the absorption by the impaired intestinal wall of the *Micrococcus rheumaticus*, and its toxins accounting for the appearance of rheumatic appendicitis and peritonitis.

PROTECTION FROM LOCAL AND GENERAL INFECTION.

The importance of this cannot be exaggerated, and the time has come when the mind of the profession, and more particularly the general practitioner, must turn to the possibility not only of treating rheumatic phenomena in children early, but of preventing their occurrence.

Throat.

(a) Hypertrophied adenoid tissue in the throat and nasopharynx should be removed in the quiescent stage.

(b) Simple congestion of the pharynx, palate, and fauces in a child with a rheumatic family or previous history, or with a rheumatic facies, should always be looked upon seriously, and met with local applications of salicylic acid preparations, together with sodium bicarbonate, sodium salicylate, potassium chlorate, and aperients. A 5 per cent. to 10 per cent. of sodium salicylate applied to the tonsils, palate, and pharynx gives a protective film from further contamination, and does not impair the defensive action of the tissues; or a gargle containing 20 to 40 grains to the ounce is equally efficacious. Care should be taken that decayed teeth are stopped or extracted, and the toothbrush and antiseptic powder should be insisted upon daily.

Pulmonary Infection.

Inhalation for half an hour, three times a day, of 10 m. of a solution of equal parts of creosote and carbolic acid with a Burney Yeo inhaler, is I believe, the best method of protecting the pulmonary mucous membrane.

Intestinal Mucous Membrane.

My experience is that sodium salicylate combined with sodium bicarbonate and rhubarb powder is by far the best protective treatment in cases where there is any indication of excess of mucus in the intestine, the alkali acting as a solvent, the rhubarb clearing the offending material away, and the salicylate acting as a sedative and healing agent to the mucous membrane.

CONCLUSIONS.

1. The *Micrococcus rheumaticus* takes the path of least resistance.

2. This may be an unhealthy throat, absorption from which frequently gives rise to general rheumatic infection, including peritonitis and appendicitis, directly through the vascular system.

3. Or it may be localized in the bronchial tubes and give rise to pneumonia, with polyarthritis and endocarditis.

4. An unhealthy condition of the intestinal wall may excite to activity the rheumatic agent, setting up acute rheumatic phenomena with peritonitis or appendicitis as part of a general infection.

5. A mild catarrh is produced at the seat of inoculation, and one or more of three factors in each case are present and promote the inroads of the micrococcus. Either

- (a) The physical resistance, or
- (b) The protective properties of the local tissue, or
- (c) Defensive agencies of the blood, are below par.

6. The distinction between acute and subacute or latent rheumatism is mainly due to general infection with the actual rheumatic agent in the former and with the toxins only in the latter.

REFERENCES.

- ¹ *Lancet*, September 22nd, 29th, 1900, pp. 861-932; *ibid.*, May 4th, 1901, p. 1260; *BRITISH MEDICAL JOURNAL*, June 4th, 1910, p. 1524. ² *BRITISH MEDICAL JOURNAL*, December 3rd, 1904. ³ *Practitioner*, February, 1903, p. 185. ⁴ *Brit. Journ. Chil. Dis.*, March, 1909, p. 97. ⁵ *Comptes rendus de la Soc. de Biol.*, Paris, 1898, vol. v, p. 128. ⁶ *Berl. klin. Woch.*, 1899, No. 29, p. 638. ⁷ *Amer. Journ. Med. Sci.*, 1894. ⁸ *Diseases of Children*, 5th edition, p. 1141. ⁹ *Diseases of Children*, 3rd edition, p. 125. ¹⁰ Harveian Lectures, *Lancet*, vol. ii, 1889. ¹¹ *Common Remedies*, Eustace Smith. ¹² *Lancet*, 1906, vol. xxii; Annual Report Loc. Gov. Bd., vol. xxxix, p. 393.