help of a fellow-practitioner I performed version and delivered rapidly. The child was dead, and had a lower dorsal meningocele. This patient made a slow but uneventful recovery.

CASE III.

The third patient, also a primipara, aged 34, called me to see her at the eighth month for severe epigastric pain. I found a large quantity of albumen in the urine, although none had been present some weeks previously. Some opium quieted down the symptoms, and a fortnight's strict rest in bed, milk diet, careful purgation, and hot baths, considerably diminished the quantity of albumen; and the oedema which she had in her legs went down. In this case, especially as I suspected the pelvis of being of small size, I induced labour at the end of the fortnight by means of the insertion of a bougie; and when labour had progressed far enough for the os to be sufficiently dilated, gave chloroform, turned, and delivered a small living child. Both mother and child have done well, though the convalescence was rather slow.

Recent examinations of the urine from these three cases proved that the albumen has cleared up in all except the last.

I would especially draw attention to two interesting points in these cases: the variability of the presence of albumen previous to the symptoms, and the symptom of epigastric pain.

As a matter of routine I always examine the urine of women who engage me to attend them in confinement, but it would seem that this examination has to be made at fairly frequent intervals to avoid being taken unawares by the advent of albumen. With regard to the sudden onset of pain in the epigastrium, I was put on my guard by my father many years ago, and feel very grateful to him for the hint. It does not seem to be mentioned with much emphasis in books on midwifery or known very widely among practitioners.

Two other cases of albuminuria in pregnancy have come under my notice in recent years. The first was an elderly primipara who had convulsions and lost her first child, but recovered, herself; when, some years afterwards, she asked me to attend her, she went through her labour satisfactorily after being kept on a careful diet after the fourth month, with fairly frequent administration of aperients. The other was a lady whom I saw convalescing from severe eclampsia, and who still showed some partial aphasia, partial blindness in the left eye, and some anaesthesia of the hand. This patient suffers from chronic pyelitis.

Dr. Gowland of Faversham has published in the *St. Mary's Hospital Gazette*, November, 1911, two terrible cases of haemorrhage, and another case of albuminuria with a fatal ending, all three with albuminuria in pregnancy. I am thankful to say that I have never met haemorrhage complicating these cases, but some years ago I lost an elderly primipara from internal bleeding after confinement, due, I think, to an extensive tear of the os uteri into the peritoneal cavity. I can remember feeling woefully helpless in trying to restore this poor woman, and to check the bleeding.

Dr. Pinniger of this town very kindly gave me the advantage of his help and advice in two of my cases.

CHICKEN-POX DURING THE PUERPERIUM.

BY

BERNARD MYERS, M.D. EDIN.,
HAMPSTEAD.

As the occurrence of chicken-pox during the puerperium is fortunately rare, I have been asked by an obstetrician to publish the following case:

A primipara, aged 30, had an uneventful parturition. The puerperium was normal also up to the tenth day. Then a vesicle, containing apparently clear fluid, appeared on the right buttock. In a few hours a second spot was seen upon the right breast immediately below the nipple. These spots were not itchy. The temperature was raised to 99° F.; the patient remained quite well in herself. As a married sister of hers was just convalescent from chicken-pox, I was on the look-out for any evidence of this complaint, more especially as my patient had kissed her sister three days before her confinement. I may mention that her sister was not aware at the time that she was suffering from chicken-pox, but her four children also subsequently developed the complaint, which they undoubtedly caught from their mother.

Twenty-four hours after the two initial spots manifested themselves the patient had a crop of spots on the chest and forehead which were typically chicken-pox. They were particu-

larly itchy. The temperature rose to 101° F. The lochial discharges, which had practically stopped three days previously, now reappeared. In the course of two or three days from the appearance of the first spots the labia majora and breasts were literally covered with pocks. There were several on each nipple. A fair number were also seen on the face, body, legs, neck, ears, eyelids, and palate. Although there were not many pocks on the legs, a good crop was seen on the perineum and around the anus. She looked and felt distinctly ill on the fourth day of the rash; the temperature remained at 101° F.

Fearing that the first spot on the buttock might turn out to be chicken-pox I immediately stopped the child being fed by its mother; it was weaned and brought to another room.

Although much troubled by the intensity and irritation of the spots on the vulva the mother made an excellent recovery without any ill effects. Each spot was treated with 1 in 40 carbolic oil night and morning.

The baby developed a temperature of 100° F. for three days after leaving its mother and showed a tiny papular erythema upon its chest. It took its food well, had no vomiting or diarrhoea, and seemed to be quite undisturbed in any way by the rash, which vanished after four days. Was this an abortive attack of chicken-pox? The little papules came out in crops, and the condition was not like any of the ordinary skin rashes which babies are subject to.

The peculiarities of the case, as far as the mother was concerned, were the predilection of the pocks for the labia majora and breasts, parts which were, one may presume, receiving a more generous blood supply due to her recent pregnancy than would otherwise obtain. The appearance of the first spot on the buttock was unusual. This spot, although at first unlike chicken-pox, subsequently went through the usual changes in a typical manner.

Memoranda:

MEDICAL, SURGICAL, OBSTETRICAL.

ABSCESS OF THE SPLEEN COMPLICATING MALARIA.

THE report by Lieutenant MacGregor, in the *JOURNAL* of February 3rd, recalls a case I met with many years ago in India. It was that of a British soldier who had served some years in India, and was much broken down by climate and service.

His final admission to hospital was for "ague," but after some weeks of irregular fever, it developed a remittent character, and was, in fact, of a more or less hectic type. This was long before the days of the microscope as an aid to diagnosis, and the true nature of the complaint was a matter of much speculation. He grew gradually weaker, and then complained of a pain in the left side. This was more or less referred to the base of the lung, but there were no signs or symptoms of pneumonia. Some friction sounds were heard, and it was thought he might have some dry pleurisy about the base of the lung. The spleen was not much enlarged (as made out by palpation), in fact, much less so than might have been expected considering the duration of the fever. He died somewhat suddenly one night, and at the post-mortem examination multiple abscesses of the spleen were discovered, which had never been suspected.

The case made a great impression upon me at the time, as I had never heard of a similar one, nor have I since, and, as noted by Lieutenant MacGregor, abscess of the spleen is not generally mentioned in textbooks. I could find no note at that time (it was in 1889 or 1890 at Lucknow). I write entirely from memory, which accounts for my somewhat "scrappy" notes. As a remarkable coincidence I may note that I quoted this very case when discussing rare and unusual cases with a medical friend this morning, only a few hours before reading Lieutenant MacGregor's case in the *JOURNAL*.

West Kensington, W. GEO. T. MOULD, Major I.M.S. (ret.).

NERVOUS RETENTION OF URINE.

WITH regard to the treatment of nervous retention of urine by injection of glycerine into the rectum, I should like to state that this treatment entirely failed in a recent case. A fortnight ago a married lady who has had two children, and who was curetted seven months ago, and who has been quite well since, consulted me for retention. This was complete, and I had to use the catheter daily. On the fifth day she was examined under chloroform by a gynaecologist. The pelvic organs were found healthy.