these paralysed muscles which are tending to recover dis-
able them again. An excellent article by Dr. Wittek
(Graz) deals with this matter in detail.

REFERENCES.

¹ Leiner, *Wien. klin. Woch.*, March 3rd, 1910. ² Levaditi, *Journ. R. Inst. Pub. Health*, xix, 1, 2, 3. ³ Hounsfield, *BRITISH MEDICAL JOURNAL*, December 2nd, 1911. ⁴ Levaditi, *loc. cit.* ⁵ Wickman, *Beitr. zur Kenntniss der Heine med. Krankheit*, 1907. ⁶ Krause, *Deut. med. Woch.*, 1909, xxxv, 1822.

ACUTE ANTERIOR POLIOMYELITIS IN SOUTH- WEST NORFOLK.

By G. F. CROSS, M.B.DUNELM.,

MEDICAL OFFICER OF HEALTH, DOWNHAM URBAN AND RURAL
DISTRICTS.

As several outbreaks of acute anterior poliomyelitis have
occurred during the past year in various parts of the
country, a short account of ten cases that occurred in this
district may possibly be of some interest.

Locality of Outbreak.

The outbreak occurred at the end of September in the
Downham Rural District, which is situated in South-West
Norfolk, and embraces both upland and cultivated fen
land, and the part of the district affected borders on the
county of Cambridge. All the cases occurred within a
radius of six miles of Downham Market.

During the months of July and August several cases of
this disease were noted in the adjacent parish of Little-
port, in the Rural District of Ely.

Age Incidence.

The ages of those attacked, and the districts in which
the cases occurred were as follows:

Group I (Southery)—

Case I	...	...	...	6 years
Case II	...	...	...	11 months
Case III	...	...	...	1 year
Case IV	...	...	...	8 years

Group II (Barroway Drove)—

Case V	...	...	...	2½ years
Case VI	...	...	...	3 years
Case VII	...	...	...	1½ years
Case VIII	...	...	...	1½ years

Group III (Crimplesham)—

Case IX	...	...	...	2½ years
Case X	...	...	...	1½ years

Mortality.

Of the above 10 cases 2 died, the second child dying
some two months after the acute stage of broncho-
pneumonia.

Previous to this outbreak another child had died of
pneumonia, in the neighbourhood of Southery, with sus-
picious symptoms of acute anterior poliomyelitis, but
which I was unable to verify.

Transmission of the Disease.

I think the following facts will aid in showing the
infectivity of the disease. Although there seems to have
been no connexion between the three groups, there is
distinct evidence of contact in several of the cases.

GROUP I.—All the cases lived in that part of the district
which is in close proximity to a part of Littleport parish
where there had been cases of a similar nature. In Case I
the father had attended Littleport market. The mother
of Case II had been helping to nurse Case III for ten days
before her own child was taken ill. Case IV attended the
school in Littleport parish, and had visited on more than
one occasion a house in this parish where there had been
3 cases with 1 death.

GROUP II.—These cases lived in Barroway Drove, which
is some eight or nine miles from Southery. There is little
or no intercommunication between the parishes. All the
cases attended the same school. Cases VI and VII were in
the same house, and Case VII was taken ill a week after
Case VI.

GROUP III.—These cases lived at Crimplesham. This
village is in a different direction about nine miles from
Southery and five from Barroway Drove. Case IX lived
next door to Case X. The parents were relatives and
were constantly in each other's houses.

Cerebro-spinal Fluid.

From Case V I took some cerebro-spinal fluid, and the
result of the bacteriological examination was as follows:

The fluid is fairly alkaline and fairly albuminous; it gives a
normal reduction with Fehling's solution. The centrifugalized
deposit is small in amount and contains a few blood discs and
leucocytes, the latter practically all of the lymphocyte type, and
but little in excess of the normal. Neither tubercle bacilli nor
pyogenic micrococci are detected in the centrifugalized deposit.
The media inoculated with this specimen only show a few
baccillary organisms which do not conform to any of the known
pathogenic types, and are most probably of extraneous origin.

The following are the notes of the cases, kindly supplied
me by the doctor in attendance:

GROUP I.

CASE I (Dr. E. G. Wales).

J. J., aged 6, Hilgay Fen. Had been ailing three days with a
"cold." On October 7th was noticed to be helpless in arms
and legs. This was complete on October 8th. When I saw him
the child was lying on his back in bed quite helpless, colour
natural, breathing natural, no pain, no vomiting, no retraction
of head. He could not raise his head from the pillow or move a
limb. Knee-jerks abolished, pupils contract moderately, no
movement on tickling soles of feet, no anaesthesia, no *tache*
cérébrale or Kernig's sign, no rash. He could swallow liquids.
Temperature 100°, pulse 86, regular. Back tender all along spine.
On October 9th he began to breathe with difficulty, turned
cyanosed and died at 10 a.m.

There were several other children in the house, but none were
affected. No connexion with Littleport except that the father
had been going to Littleport Town at intervals. A week previous
a child on Ten Mile Bank died from acute pneumonia which
might have been secondary to acute poliomyelitis. He was
helpless and moribund when seen, with both lungs intensely
congested. Father a travelling baker who visits all the houses
in the neighbourhood.

CASE II (Dr. Glenfield Martin).

V. L., aged 11 months, Southery Ferry. For a fortnight pre-
vious the mother had frequently visited the house of Case III and
had nursed the child. Taken ill on September 18th with fever
and bronchitis. The mother noticed some paralysis and that the
stools were very offensive. Seen by Dr. Martin on October 2nd.
Signs typical of acute anterior poliomyelitis. Paralysis of left
leg. The offensive stools could not be affected by medicine.

Later in December he reports: "Tibial muscles of left ankle
are paralysed and wasted. If foot were not kept in a splint it
would become deformed (nature of valgus). Three other
children in house all well. Prognosis is fair but hardly as good
as two months ago."

CASE III (Dr. Hutt).

A. L., aged 12 months, Southery Ferry. I first attended the
child on September 12th. The mother thought the right
shoulder had been damaged by undressing the child. Tempera-
ture 102°, neck somewhat swollen, and did not seem to be able
to keep the head in any position long. Deltoid (right) paralysed
and perhaps other muscles. September 14th, temperature
normal, paralysis right upper arm and both legs, adductors
most marked. Present condition, January, 1912: Right leg much
improved and arm very fair movement, left leg some slight
shortening and foot in marked talipes and equinus position.
Adductors (thigh) and peronei still useless.

CASE IV (Dr. Martin).

B. J., aged 8 years, near Southery Ferry. Previous to being
taken ill had visited a house in Black Horse Drove, where there
had been three cases of illness (one fatal). Was first taken ill
on October 5th, with fever and bronchitis. On October 11th,
when first seen, had pyrexia and headache. On October 15th
slight paralysis, well marked the following day. Tenderness of
affected limbs (both legs), muscles of trunk also affected, tender-
ness and pain at back of neck. Stools highly offensive. At first
apparently no progress, but lately the progress has been good.
Weakness of the flexor muscles of both thighs, no deformity
although some rigidity. This case was subsequently notified
for tuberculosis.

GROUP II.

CASE V (Dr. Wales).

F. B., aged 2½ years, Barroway Drove. Brought to surgery
with symptoms of meningitis on October 22nd. Had a convul-
sion on night of 20th. No vomiting, fretful, sleepless. Tempera-
ture 104°, pulse 130, regular. Pupils equal and react. No
paralysis of limbs. Slight retraction of head, cannot sit up
cries when touched. Rapid improvement under treatment.
Temperature normal on 24th, when slight internal strabismus
of left eye was noticed. By 30th was much better, taking good
food, sleeping well. Internal strabismus (paralysis left external
rectus—third cranial nerve) more marked. Several other
children in house all well.

CASES VI AND VII (Dr. Wales).

A. W., aged 3 years, Hooten's Row, Barroway Drove. Seen
on October 24th. Had been ailing two or three days, became
paralysed in all limbs and back. Temperature normal,
pulse 120, breathing normal, no pain, can move feet if soles are
tickled, but not legs. November 1st, Dr. Cross visited with me

and made a lumbar puncture. In January this child much improved, but cannot use right leg.

A. B., aged 1½ years, occupying the same room, became ill with loss of power in back, cannot sit up and cries if "bent." Temperature normal, limbs unaffected.

Case VI died in January from bronchopneumonia.

CASE VIII (Dr. Cross).

A. J. S., aged 1 year 8 months, Barroway Drove, near Case V. Mother noticed for a week previous that he fell about, and attributed this to some new boots.

On October 31st he fell down and could not get up again. On November 1st visited and found child lying on sofa. Temperature 101.4°, restless and fretful, breathing natural. Any attempt to straighten limbs caused child to cry out. Rigidity of muscles of neck, and spine slightly arched. Bowels constipated. Two days afterwards temperature was normal and child appeared better. Left leg appeared useless. The child gradually got better, and after four weeks was able to sit up and crawl about.

Present condition: Can stand with assistance, but cannot control the left leg, which is slightly wasted. No shortening. Health otherwise good.

GROUP III.

CASE IX (Dr. Wales).

S. S., aged 2½ years, Crimbleham. Taken ill November 5th. On 6th temperature 103°, no definite symptoms, was thought to be a stomach attack. Seen on the 8th. Temperature 101°. Crying about aching in arms, difficulty in arm movements, moved legs freely. Left arm gradually became quite paralysed and remained so. The acute illness lasted five days.

CASE X.

A. R., aged 1 year 3 months, Crimbleham (living next door to Case IX). Mrs. R. had been backwards and forwards. Became ill about November 15th. A very indefinite illness, fretful and refuses food. Declines to be "sat up," and keeps back rigid, lies on back passively and cries if moved. On November 28th all symptoms have disappeared except for weakness and rigidity in back.

REMARKS.

There had been no abnormal amount of sickness in the district, and very few cases of infantile diarrhoea.

Four of the cases lived in good farmhouses with well-to-do parents, three in good cottages, and there were only two houses in which the sanitary conditions were unsatisfactory.

Eight of the cases were in the fen, some being a good distance from the high road, and were therefore not subject to any unusual amount of dust.

There were no known cases of paralysis in animals or fowls on the various premises.

Flies and other insects were abundant, and in three of the houses there were undoubtedly a number of fleas.

With the exception of Cases III and IV there were other children in all the houses, but in only one house was there more than one case.

SERUM AND VACCINE THERAPY IN CONNEXION WITH DISEASES OF THE EYE.

THE MIDDLEMORE PRIZE ESSAY, BRITISH MEDICAL
ASSOCIATION, 1911.

By C. W. G. BRYAN, F.R.C.S. ENG.,

LATE ASSISTANT IN THE DEPARTMENT FOR THERAPEUTIC INOCULATION,
ST. MARY'S HOSPITAL; AND HOUSE-SURGEON, OXFORD EYE HOSPITAL.

(Concluded from page 665.)

III.

THE SERUM TREATMENT OF DISEASES OF THE EYE.

SERUMS, antitoxic and antibacterial, have been used very extensively in treating cases of eye disease, and may be divided into two classes—those which have a specific action and those which are non-specific (the so-called "paraspecific serums").

Serum has usually been injected either into the subcutaneous tissues, the most frequently used and safest method; into muscular tissues; or into the blood stream, the method by which the most rapid effect is obtained. It has also been given by mouth and by rectum. Darier⁸ concludes that serums given by mouth immunize but exert no bactericidal effect.

Specific Serums.

1. *Diphtheria Antitoxic Serum*.—As in diphtheria elsewhere, so in diphtheria of the eye extremely good results have been obtained by the use of this method of treatment. Sydney Stephenson¹ has reported cases successfully

treated in this way. The dose should be 2,000 to 10,000 units, and at least two doses are usually necessary. In membranous conjunctivitis due to the Klebs-Loeffler bacillus the Roux serum is valuable; Axenfeld² has analysed the results obtained and reports favourably. In corneal diphtheria the serum is less effectual, and this is due to the fact that, though the serum does overcome the toxic effects of the *Bacillus diphtheriae*, the severity of the corneal complications depends on the mixed nature of the infection from the presence of the ordinary organisms of suppuration, staphylococci, streptococci, pneumococci, etc.; the serum has only a non-specific effect on these organisms. In such mixed infections Römer's antipneumococcus serum and polyvalent anti-streptococcus serum are of use. Post-diphtherial paralysis of eye muscles: Aubineau³ reports a case of paralysis of accommodation cured by injection of anti-diphtheria serum. Three relapses subsequently occurred, each of which yielded to the treatment. Four doses (10 c.cm., 20 c.cm., 30 c.cm., 60 c.cm.) were given in this case.

2. *Tetanus Antitoxic Serum*.—This serum, of the same nature scientifically as antidiphtheria serum, is of value in treating tetanus having its primary point of infection in the eye. Such cases are rare, but Ellis has had two recoveries of three cases treated. Lewis also has reported the recovery of a case. Diphtheria and tetanus are the two diseases due to organisms possessing an extracellular toxin, and for this reason are especially suitable for the use of specific antitoxic serums. Specific serums have been prepared and used in infections by many other organisms with varying results. Possibly the benefit obtained in some cases does not depend on the specific nature of the serum as it does in diphtheria and tetanus. As I shall mention later, in infections due to the ordinary pyogenic cocci excellent results have been obtained with non-specific serums.

3. *Specific Antipneumococcus Serum*.—Many methods of preparing antipneumococcus serum have been employed. Kornel Scholtz⁴ finds that the serum produced by infecting an animal with one culture of pneumococcus usually has no effect in agglutinating other cultures of that organism; these experiments demonstrate the existence of different strains of pneumococci. It is impracticable in clinical medicine to prepare a serum by immunizing animals to cultures obtained from individual patients, and therefore the employment of a polyvalent serum, prepared from many strains, becomes necessary. The best known polyvalent antipneumococcus serum was that of Römer. Römer⁵ has lately modified the method of preparation of his serum, and has introduced a new serum on the lines of Bail's antiaggressin-containing serum; using this new serum, he reports success in the treatment of pneumococcal ulcers of the cornea.

4. *Antirheumatic Serum*.—Rosenthal claims good results in rheumatic iritis by the use of a specific antibacterial serum prepared by immunizing animals against the *Streptococcus rheumaticus*. The results have not been corroborated, and Darier⁸ says better results are obtained in such cases by using Roux's serum.

5. *Antistreptococcus Serum*.—Polyvalent antistreptococcus serum has given occasional good results in streptococcal infections of the eye, but the results are not constant owing to the enormous number of strains of streptococcus which have been demonstrated. Recent methods of analysing the strain infecting any particular case by means of the sugar reactions may make possible the use of antistreptococcus serum prepared with a corresponding strain. Snell⁶ reports a case greatly benefited by antistreptococcus serum. The patient had suffered from suppuration of the lids of the right eye for seven days, the temperature was 103°, delirium was marked, and the general condition grave. Streptococci were found in pure culture. Three injections, each of 10 c.cm. of polyvalent antistreptococcus serum, were given over a period of forty-eight hours. On the following days the temperature was subnormal, delirium abated, and eventually a cure resulted. On the other hand, Lawson⁷ found antistreptococcus serum of no use in a case of acute streptococcus infection of the cornea.

Antigonococcus Serum.—An antigenococcus serum has been prepared by Rogers and Torry by injecting living gonococci into the peritoneum of sheep. Knapp⁷ reports results of the use of this serum. He has found it of no