Presidential Addresses

TO THE

BRANCHES OF THE BRITISH MEDICAL ASSOCIATION.

LANCASHIRE AND CHESHIRE BRANCH.

SURGICAL MILESTONES.

By ROBERT C. BROWN, M.A.CANTAB., M.B.,

F.R.C.P.LOND., F.R.C.S.ENG.,

Consulting Medical Officer to the Preston and County of Lancashire Royal Infirmary.

"Did you see *Milestones*?" is a question very frequently put by the friends of those who have been spending a few days in London; and the reply is almost invariably, "Yes; I was delighted with it." And the play suggested to me a title for the remarks which I now propose to make to you on "some items in the progress of surgery and its auxiliaries between 1836 and 1913."

Milestones presents to our notice several families; animated discussions in reference to the relative value of wood, iron, and steel in the construction of ships; changes in the styles of ladies' dress at three periods—1860, 1885, and 1912; the pictorial illustrations which accompany the programmes showing the crinoline or inflated period, the dress improver with local prominence period, and the non-crinoline or collapsed period; daughters who resented the dictation of their parents as to whom they should marry, and a young lady who, in a moment of revolt, broke off her engagement in 1860, and repented ever afterwards having done so, and whom we recognize in 1912 as a charming old maid. The piece has been running since March, 1912, and is likely to continue to attract for some time to come.

The subject of my address is a much more serious and important one, and my milestones are: 1836, 1847, 1867, 1882, and 1895. The progress of surgery and its auxiliaries between 1836 and 1913 has conferred upon mankind benefits and blessings which cannot be too highly appreciated, although their great value is apt to be overlooked on account of our familiarity with them.

If any one had ventured to prophesy in 1836 that such operations as now come under the category of the surgery of the abdomen would be performed in 1913 he would have been regarded as a lunatic. I heard the late Sir W. Mitchell Banks tell his audience, at the opening of the operating theatre at the Liverpool Royal Infirmary by Sir F. Treves a few years ago, that if in the earlier part of his career he had attempted to perform operations which he was then undertaking as a portion of his daily work he would have landed himself in the hands of the police. The mention to him of such operations as gastro-enterostomy, or the removal of the uterus or of the prostate gland, would have made him shrug his shoulders and use some of the impressive language with which his friends were familiar.

First Milestone, 1836.

The modern system of nursing had its commencement in 1836. The credit of inaugurating the new order of things belongs to Germany, for it was at Kaiserwerth that the institute for training deaconesses was founded, and it was at this institute that Florence Nightingale acquired the practical knowledge which enabled her afterwards to turn her remarkable gift of organization to such brilliant account. During the nineteenth century the qualifications of nurses were greatly raised by the influence of Florence Nightingale, Mrs. Fry, and others, amongst the most important of whom may be numbered Queen Victoria, Queen Alexandra, and our present Queen. In 1877 the St. John Ambulance Association was founded, and in association with it lectures to women on home nursing and hygiene are now given regularly in all parts of the country. About ten days ago Queen Alexandra, accompanied by her sister the Dowager Empress of Russia, inspected at Devonshire House over eleven hundred of the nurses associated with the Queen Victoria Jubilee Institute. They came from all parts of England, Wales, Scotland, and Ireland. Every one of these

women has had three years of hospital training and an additional special preparation for work in ill-equipped homes. They spend their days in the homes of the poor, going from case to case, dressing, advising, instructing, or contriving some rough-and-ready device for a back-country case where necessity must invent. The nurses in the early part of the century were of a type so graphically described by Dickens in *Martin Chuzzlewit*, *Sketches by Boz*, and other papers. Most of you will remember one of them, who was called Mrs. Gamp, and who Dickens tells us was a fat old woman with a husky voice; her face, the nose in particular, was red and swollen, and it was difficult to enjoy her society without being conscious of a smell of spirits. Her usual accompaniments were a species of gig umbrella, a pair of pattens, and a snuff-box. Mrs. Gamp lodged at a bird fancier's in Kingsgate Street, High Holborn, next door but one to the celebrated mutton-pie shop, and directly opposite the celebrated cats' meat warehouse. Her bedroom was in the first floor front, and the window was easily assailable at night by pebbles, walking sticks, fragments of tobacco pipes, or a cabdriver's whip; these were all much more efficacious than the street-door knocker, which was so constructed as to wake the street with ease, and even spread alarms of fire in Holborn without making the smallest impression on the premises to which it was addressed. When thus vigorously applied every window in the street became alive with female heads. Every surgeon knows what an important auxiliary a good nurse is both before, during, and after an operation. It is unnecessary for me to offer an abstract notion of such a treasure, as all are familiar with concrete examples, and you also know how the nurses' home of to-day differs from that of Mrs. Gamp.

Milestone 1847.

When I became a student at King's College Hospital in 1855 anaesthetics had not been very long in general use, for it was only in 1846 that ether was employed, whilst chloroform was used for the first time in a surgical operation in the Edinburgh Royal Infirmary on November 15th, 1847. Sir James Simpson made the first experiment with chloroform in his own house on November 4th, 1847. Before chloroform was suggested to him by Waldie, a Liverpool chemist, Simpson personally experimented with several chemical liquids in the hope of finding something less disagreeable and persistent in smell than ether. The dream of Simpson's life had been to soothe the pain of his patients, and, before he realized his dream, he had been almost driven by the painful scenes which he witnessed in the operating theatre to abandon medicine for the law. Sir William Fergusson was in the height of his fame when I went to King's College Hospital. He had been appointed one of its surgeons in 1840, and as chloroform was not used until 1847, he had the experience of performing operations before the introduction of anaesthetics, and comparing it with that of performing operations under anaesthesia. He could tell us how patients narcotized with opium, or intoxicated with alcohol, and uttering piercing screams and sometimes startling oaths, were forcibly held down by strong assistants. He could contrast the agony which, before the introduction of chloroform, people had to undergo both in anticipation of and during the performance of surgical operations, with the comparative ease of mind with which such proceedings could after its introduction be contemplated.

Milestone 1867.

In 1867, the antiseptic or aseptic method of treatment (as it is now called) was introduced by Lord Lister, and as a result of it the number of operations performed for the cure of disease or deformity increased enormously, and rates of mortality which in some cases were scarcely less than 50 per cent. were reduced to as low a point as 1 per cent. Many diseases which were previously beyond the reach of treatment are now easily cured by an operation, and therefore an inestimable benefit has been conferred on mankind. The use of chloroform and other anaesthetics, by removing the fears and dangers attendant upon pain, led to a great increase in the number and complexity of operations, but the mortality following some of them was so terrible that they had to be abandoned. Ovariectomy was one of these, but since the introduction

of the antiseptic or aseptic method of treatment, it has been undertaken again, and performed with such remarkably successful results as to make it clear that the aseptic method was the one thing necessary to complete the usefulness of chloroform and other anaesthetics. The number of females who have not only been rescued from certain death by the operation of ovariectomy, but have been restored in the course of a few weeks to perfect health, is one of the most startling facts in the history of operative surgery. It is very interesting to read some remarks which Sir Frederick Treves made in 1897 and compare them with the state of affairs at the present time. He said: "Sixty years ago the operating room was the dirtiest room in the hospital; the surgeon operated in the dirtiest coat in his possession—a coat stiff with blood and animal filth; he was as proud of this blood-stained rag as a peer of ancient lineage may be of his faded ceremonial robes." An up-to-date operating theatre is an ideal of the most perfect cleanliness. The floor, walls, and roof are covered with glazed tiles, which can be washed with the greatest ease; there are no angles in which dust can accumulate, and the most complete arrangements are provided for washing the hands and the instruments; there is a separate room for the sterilization of the dressings, the basins, and other utensils, and for boiling the water; there is a room in which the patient is anaesthetized by a doctor in the presence of one nurse before being wheeled into the operating theatre; there is a room in which the surgeons hold their consultations, and the equipment of the lavatories is all that could be desired. On a dull day, or in the night, the theatre is brilliantly illuminated by electricity. The operating table is a model of ingenuity. The operating surgeon and his assistants wear a complete suit of white material, such as drill; they have caps on their heads and india-rubber gloves on their hands, and their faces are covered as far as the eyes with a mask of gauze, so that the eyes are the only portion of the body which is visible.

Milestone 1882.

In 1882 very valuable information was communicated by Koch in reference to the tubercle bacillus, although its existence had been discovered by Baumgarten a few years previously. The revolution which has been caused by the discovery of the tubercle bacillus is something like a romance. Attention has lately been called to the circumstance that the common household fly, which is such an annoyance to housekeepers, and with whose unpleasant tickling we are all so familiar, is a great friend of the tubercle bacillus and is a factor in the dissemination of tuberculosis, as well as of typhoid fever and other diseases.

Milestone 1895.

The x rays, one of the most important auxiliaries to the surgeon of the present day, were discovered in 1895. The wonderful revelations made by the x rays have excited profound interest and placed the treatment of fractures of the bones and dislocation of the joints on a more satisfactory basis than it ever was before. It is now possible to examine them as if the flesh was transparent, and not only to examine them, but also to take photographs of them. The position of bullets, needles, pieces of steel, and many other foreign bodies can be ascertained and photographed without resorting to the painful process, so long in vogue, of probing for them. Some of the diseases of internal organs can also be elucidated by the x rays, and by their agency several diseases of the skin can be cured.

The importance of the telephone to the surgeon can be best appreciated by supposing the privilege of using it to be withdrawn. The first successful exhibition of the telephone was made in London on July 14th, 1877. With the telephone the surgeon makes many of his appointments, both at the hospital and in his private practice; he gives instructions to his house-surgeons and others; he sends for a taxicab; when detained at an operation and unable to return home at the time appointed for his consultation hour, he is able to relieve the anxiety of the patients who are waiting to see him by explaining to his servant the cause of his detention. There is another use of the telephone which I can best illustrate by giving you a case which was recorded in some of the newspapers. The child of a young married lady was taken ill in the night with symptoms which she thought were those of

croup, a disease of which she had had no previous experience. She at once telephoned to her medical man and asked him what she ought to do. His reply was, "Hold Johnny up to the telephone and let him cough." When this had been done the doctor told the lady that Johnny was not suffering from croup, that she might put him in bed again, and he would call and see him in the morning.

We did not hear much about appendicitis until 1901, when our dearly beloved peace-making King Edward VII narrowly escaped being taken from us by that disease. I remember that at the commencement of his illness the newspapers informed us that he was suffering from a "chill on the liver," a complaint which has no place in the recognized nomenclature of diseases. Sir Samuel Wilks has called attention to the fact that the disease existed and was recognized long before the term "appendicitis" was applied to it, and he has shown that Dr. Addison described "coecitis" or "typhilitis" in his regular course of lectures on medicine at least as early as 1836. Very little has been added since then except that which relates to bacteria and surgical treatment. Moreover, the distinct share of the appendix in the causation of the disease was recognized, although it was not credited with quite so much responsibility as we are now inclined to ascribe to it. In the index to Sir Thomas Watson's lectures on the *Principles and Practice of Physic*, which was the standard work when I was a medical student, neither in the 1857 edition nor in that of 1871 does the word "caecum" or "appendix" occur, but in the chapter which enumerates the causes of peritonitis there is the following passage: "It is a curious fact that the vermiform appendix of the caecum is not infrequently the seat of a penetrating ulcer."

I have traced little groups of glands in that slender tube, and I have known perforation to happen from the specific ulceration of typhoid fever, and from the accidental ulceration caused by a cherry-stone lodged there in one instance, and by a pellet of hard faecal matter in another. The last mentioned cause is of not infrequent occurrence. I and others can recall—and it is a painful reflection—many patients who were undoubtedly suffering from appendicitis, although we were not aware of the fact, and whose lives would probably have been saved by the operation which is now so frequently and successfully performed in all parts of the world and at all hours of the day and night. An American surgeon who is in favour of an early operation says: "The affected appendix is a cap which sometimes snaps, sometimes flashes, and sometimes causes an explosion, and none of us can tell in advance what is going to happen."

Few operations are more beneficent in their design or more successful in their result than those which have been undertaken for the removal of gall stones.

The lives of those who are liable to recurrent attacks of biliary colic, which cannot be prevented by medical treatment, are rendered miserable by the constant fear of a return of their trouble. Sir Thomas Watson says: "The pain that attends the passage of a gall stone is often dreadful. Perhaps there is no pain to which the body is subject that is more severe. You will not wonder at this when you consider that through a tube of which the natural size scarcely exceeds that of a goosequill there sometimes passes a stone as big as a walnut." Even the most simple and uncomplicated form of the operation—which consists in incising the abdominal wall, opening the gall bladder, removing the gall stones, and stitching the edges of the gall bladder to the abdominal wall, leaving a fistula which heals within a few weeks—would not be undertaken with safety before the adoption of the aseptic method of treatment.

The operation of tapping the chest, which is now performed so very frequently, must have been a somewhat unusual occurrence ten years before the introduction of Lord Lister's principles. Writing in 1857, Sir Thomas Watson, one of the most eminent physicians of that day and a colleague of Sir William Fergusson, said: "On six occasions I have myself witnessed the evacuation, by puncture, from the human pleura of a clear transparent liquid. Three of the patients recovered, three died. Again, I have twice seen pus let out by the primary puncture of the chest; one of these recovered and the other died. This constitutes the amount, or nearly so, of my

personal experience of the operation of paracentesis thoracis." In reference to healing up the orifice or keeping it open, Sir Thomas remarks: "On this point practical men differ. I should say, if pus comes out, by all means make the aperture large and keep it open; and inasmuch as detention of the pus would be injurious and the depending point is difficult to hit, and the orifice is apt to clog, I would do more than leave it open; I would draw the puriform fluid off twice a day by a siphon; if serum be let out, by all means close and heal the wound." He then mentions the circumstances under which he would reopen the wound. When I see fluid let out of the chest with that beautiful instrument, Dieulafoy's aspirator, I recall the only occasion on which I saw paracentesis thoracis performed at King's College Hospital. It was in 1858, and the fluid was conveyed, not through an india-rubber tube into a basin, but through a piece of chicken's intestine, the distal end of which rested under the surface of water which had been poured into a bucket.

When I went to King's College Hospital in October, 1855, Sir William Fergusson was in the height of his fame, and the brilliancy with which he performed an amputation or a lateral lithotomy, the delicate manipulation with which he operated for cleft palate, attracted many visitors to his theatre. Amputations, however, were becoming less and less frequent as the conservative method of excising the diseased joint, instead of removing the limb, was becoming more generally adopted. Indeed, Sir William Fergusson and Mr. Syme of Edinburgh were two of the chief pioneers in this branch of operative surgery. This was, of course, twelve or fifteen years before the introduction of antiseptics, but since this great revolution in treatment excisions have been more frequently undertaken and with more satisfactory results. But just as amputation has been superseded in suitable cases by excision, so excision is nowadays superseded by arthroectomy. In this operation, instead of sawing off layers of bone, diseased foci only are gouged out, and the resulting cavities are disinfected by carbolic acid and packed with sterilized iodoform. Influenced by Sir William Fergusson's great reputation, and attributing to him a power which he did not possess, many patients came to see him under the impression that he could restore to its normal flexibility and usefulness, in the course of a few days, an arm which had been fractured and from which the splints had only just been removed. He always treated these people with great kindness and sympathy, and informed them that in due course the limb would be as good as it was before the accident. He also instructed them to come and see him again in three months if by that time the limb had not recovered its normal flexibility and usefulness. Knowing, however, as he did that most of these patients would not be satisfied unless they had "something to take," he almost invariably told his house-surgeon to order them 5 grains of compound galbanum pill three times a day. The composition of this pill, I remember, was galbanum, myrrh, sagapenum, asafoetida, soft soap, and treacle.

In these days, when so many people are knocked down and run over by motor omnibuses, motor cars, and motor cycles, serious internal injuries are often sustained, which render it necessary to make an incision through the soft parts between the ribs and the hips, or, in other words, to do a laparotomy, for the purpose of exploration. Before the days of aseptic surgery this procedure would have been regarded as unjustifiable. Since its adoption, however, tears in the stomach, intestine, bladder, and other organs have been stitched up so as to prevent any further escape of their contents into the peritoneal cavity, and in this way many lives have been saved which would otherwise have been lost. Death from internal haemorrhage has also been prevented by discovering the bleeding vessel and ligaturing it. It is a very wonderful indication of the advance which has been made in operative surgery since the aseptic method of treatment was adopted, that cases are on record in which the entire stomach has been successfully removed, and that it is now no uncommon occurrence to remove the kidney, spleen, gall bladder, prostate gland, and uterus; to suture a perforating ulcer of the stomach, duodenum, and other portions of the intestine; to excise a diseased part of the stomach or intestine, and to unite the proximal and distal ends by suture. It has now even been undertaken to open the

skull and remove a tumour from the brain, and to open the spinal canal for the removal of tumours from the spinal cord.

An American surgeon is said to have succeeded in transplanting the limb of one animal to some other animal, and in the transference of vital organs, such as the kidney, from one animal to another. We live in an age of wonders, and it will not be surprising if what has already been done for certain of the lower animals may be accomplished for human beings.

When we reflect on what operative surgery has accomplished between 1836 and 1913, we are disposed to wonder what its record will be at the end of the century, and perhaps to regret that we were not born seventy years later, in order that we might see the still further progress which will be made, not only in surgery, but in art, literature, and all other branches of knowledge. We ought to be very thankful that we have been allowed to live during a portion of the century which has been more remarkable than any of its predecessors for the discovery and invention of appliances which have added so much to the interest and enjoyment of life, and it is our duty to do our best to continue the good work which was carried on by those who have gone before us.

YORKSHIRE BRANCH.

THE DEFECTS OF THE INSURANCE ACT AND THEIR REMEDY.

By WILLIAM STANGER, F.R.C.S.ENG.,

President of the Yorkshire Branch.

THE Insurance Bill threatened to place us in an entirely new relation to the State. It imposed upon those who accepted it a statutory obligation to attend insured persons. Our responsibility would therefore be doubled. Before it was to the patient alone, now it would be to the patient and the Insurance Commissioners. It is true many of us had been accustomed to take clubs, but this was partly as a charity, and the tie which bound us to clubs was one of mutual good will. We had not to keep records, file prescriptions; there was no question of medical inspection or of a court of complaints.

When the bill became an Act of Parliament, we were in this position: Those of us who had clubs found that almost all former club patients were now insured persons, and, worse still, many of our former private patients were now entitled to medical benefit in return for their weekly subscription of threepence or fourpence. We were therefore threatened with a great loss of income unless we accepted the Act.

In the meantime, owing to the action of the British Medical Association, and our unanimous refusal to work the Act as at first passed, the Chancellor was forced to make certain important concessions. The inadequate sum of 6s. was raised to 8s. 6d., or rather 9s., including 6d. for tuberculosis. The records were to be of the simplest kind. The medical inspection was declared to be only for statistical purposes, and the court of complaints was transformed into the Medical Service Subcommittee. The profession was very reluctant to give up its old freedom, which it valued far more than money. At the same time, it was felt that refusal to work the Act would mean to many men working in industrial districts absolute financial ruin, and this feeling became so strong that, when the time came to decide, a large majority of medical men, driven by economic necessity, resolved to accept the Act. At the present time, 13,600 doctors are treating 14 millions of insured persons.

On the whole, the Act has proved financially beneficial to those practitioners who work in industrial districts. In some cases their incomes have been doubled. On the other hand, doctors who work amongst the middle class have in many cases suffered a great loss. Patients who were well able to pay a moderate bill have to be treated now for 7s. a year. Doctors who practise amongst the upper class, and consultants and specialists, are not affected by the Act.

As we have accepted the Act, though with reluctance, it behoves us to try and improve it either by amendments to the amending bill which has been introduced, or by