season, it was still in a rather lethargic state, thus permitting a human being to get close to it, otherwise it would have been off at the first approach.

AN ENORMOUS PAROTID TUMOUR IN A CHINESE WOMAN.

By HAROLD BALME, F.R.C.S.ENG.,

SCHOFIELD MEMORIAL MISSION HOSPITAL, TAI YUAN FU, NORTH CHINA.

THE accompanying photograph is that of a Chinese woman, aged 53 years, who was first seen by me at a mission station in Southern Shansi, North China, in February, 1910. According to her history, she had first noticed a lump some eleven or twelve years previously, which she described as a small, round, hard substance, near the right ear. The tumour had grown quite slowly for the first few years, but during the last eighteen months it had increased much more rapidly, until its dimensions and weight had become so considerable that she could not bear it unsupported. For the last few months she had been obliged to sit on the Chinese kang all day long supporting the tumour with her hands. She was advised to make her way to the Baptist Mission Hospital in Tai Yuan Fu, the capital of Shansi, where she arrived some days later.

Condition on Admission.

The patient was very *sallow* and *emaciated* on admission; this was attributed to the fact that she was an opium smoker. On the right side of the neck, extending from the malar bone to the level of the breast, and from the middle of the chin to some distance beyond the posterior border of the sterno-mastoid muscle, was an enormous tumour firmly attached to the structures in the parotid region but quite separate from the lower portion of the neck, which it overhung. The tumour was firm and elastic and definitely lobulated; its total circumference, so far as it could be measured, was considerably larger than that of the patient's own head. The skin was freely movable over the growth, on the surface of which several large vessels ramified; the tumour itself was apparently encapsulated.

Before any surgical interference was possible, the patient underwent the ordinary course of treatment for the breaking off of the opium habit, and it was not until she was almost free from opium, and in a more favourable condition of health, that an operation was attempted.

Operation.

With the assistance of Dr. Paula Maier, an enormous elliptical incision was made from the malar bone to the

sternal end of the clavicle, and back to the lobe of the ear. Little difficulty was experienced in shelling out the tumour, though a great number of large vessels were met with, and in the enfeebled state of the patient these all needed to be isolated and clamped before section. There was a large hollow between the angle of the jaw and the mastoid process, forming the bed of the tumour, the surrounding bone having to a considerable extent atrophied from pressure. A large portion of skin, which was extremely redundant, was removed with the tumour, and the wound was closed in the ordinary way.

After-Treatment.

It was necessary to give two or three pints of saline intravenously before she left the operating table, and for the first twenty-four hours the patient suffered severely from shock. She needed close attention for some considerable time after this, and for the first two or three days Dr. Paula Maier remained almost continuously with her. Her subsequent recovery proved uneventful. When leaving the hospital she had a remarkable hollow behind the angle of the jaw, but I have since heard that this has to a large extent filled up.

Description of Tumour.

The tumour, which weighed over 20 oz., has been presented to the Museum of the Royal College of Surgeons, and the following report upon it has been received from Mr. Shattock:

Tumour from Neck of Chinese Woman.—This is a myxo-endothelioma. No cartilage found in the sections examined. Some parts of the stroma are very cellular, indicating rapid growth, but the zone of fibrous tissue limiting the growth excludes any sarcomatous change having occurred at the periphery.

PERFORATED ENTERIC ULCER: OPERATION: RECOVERY.

By WILLIAM WASHBURN,

SURGEON TO THE GLOUCESTERSHIRE ROYAL INFIRMARY AND EYE INSTITUTION.

J. M., male, aged 10 years, was admitted to the Gloucestershire Royal Infirmary on February 3rd, under Dr. Grosvenor, with enteric fever, characterized by dry tongue, constipation, much evacuation, and a temperature ranging up to 102°. There was a positive Widal reaction. On March 1st, during the fifth week of his illness, he suddenly became collapsed, and complained of pain in the right side of the abdomen. His temperature fell to 97° and the pulse rose from 110 to 156. The abdomen was retracted and hard, and there was considerable impairment of movement on respiration.

The house-physician, Mr. H. Hopps, recognizing that in all probability perforation had occurred, immediately sent for me, and within three-quarters of an hour of the onset of the collapse I opened the abdomen in the middle line below the umbilicus, and found a perforation of the ileum $\frac{1}{2}$ in. in diameter, some 3 in. from the ileo-caecal valve. The intestines were bathed in clear serous fluid, and there were about 3 oz. of a similar fluid in the pelvis. There was no other evidence of commencing peritonitis. I closed the perforation with a sero-muscular purse-string suture of fine silk, mopped the fluid from the pelvis, and inserted a small rubber drain. The wound was closed with through-and-through silkworm gut sutures.

The patient was back in bed within half an hour of his leaving the ward. The tube was removed twenty-four hours after the operation. Its insertion was probably an unnecessary precaution.

Recovery was uneventful, and the patient is now (May 9th) in excellent health.

At the request of the Bishop of London, we draw attention to the seaside camps for London working boys. The first of them was opened some twenty-four years ago, and year by year the work has extended, until last year about 3,700 lads were provided with accommodation for a week or two at the seaside. Many readers have probably seen these camps, and none can have any doubt as to the physical and moral advantages likely to be derived from their upkeep. Each boy contributes to the expense on an age scale, but, of course, the total outlay is not covered thereby. Contributions will be gladly received by the secretary of the movement, Mr. A. Abel Bloxam, 22, Northumberland Avenue, W.C.