the more common becomes this disease in the female sex; whereas general paralysis is common to all grades of society in the male sex, it is very rare in the better and respectable classes of females in all grades of society. Probably these statistics give a rough idea of the relative prevalence of syphilis in the two sexes.

We may conclude, therefore, that syphilis is comparatively commoner in the West End among males and comparatively in the poor East End among females. Seeing that a very considerable number of paralytics in the West End do not come into the county asylums, it may be assumed that the ratio of six males to one female is well under the mark.

BIBLIOGRAPHY.

- ¹ Bruce, Sir David: Sleeping Sickness Commission, vol. iv.
- ² Candler, J. P.: Results of Wassermann Reaction: Discussion on Syphilis, *Proc. Roy. Soc. Med.*, vol. v, Supplement, 1912.
- ³ Ehrlich: The Harben Lectures on Experimental Researches on Specific Therapeutics, Lecture I, *Lancet*, June 15th, 1907, p. 1634, and Lecture III, *Lancet*, August 10th, 1907, p. 351.
- ⁴ Forster and Tomaszewski: *Deut. med. Woch.*, 1913, Band xxxix, p. 1237.
- ⁵ Fournier: *Les Affections Parasyphilitiques*, 1894.
- ⁶ Goldmann, E.: *Vital Färbung am Zentralnervensystem*. Verlag der Königl. Akademie der Wissenschaften, Berlin, 1913.
- ⁷ Levaditi et Yamanouchi: L'Orientation de la Syphilis, *Comptes Rendus de la Société de Biologie*, vol. lxxiv, 1908. *Ibid.*, *Annales de l'Institut Pasteur*, vol. xxii, No. 10, October, 1908.
- ⁸ Marie, Levaditi, et Bauchowski: Présence Constante du Tréponème dans le Cerveau des Paralytiques Généraux Morts en Ictus, *Comptes Rendus de la Société de Biologie*, vol. lxxiv, No. 17, 1913.
- ⁹ Marinesco et Minea: Absence de *Spirochaeta pallida* dans le Système Nerveux Central des Paralytiques, *Société de Neurologie*, April 5th, 1906. *Ibid.*, Présence du *Treponema pallidum* dans la Paralyse Générale, *Comptes Rendus Hebdomadaires*, vol. lxxiv, No. 16, 1913. *Ibid.*, Association de Ménigite Syphilitique et de Paralyse Générale. Présence de Tréponèmes dans les Méninges, *Réunion Biologique de Bucarest*, March 6th, 1913.
- ¹⁰ Mott, F. W.: *Proc. Roy. Soc. Med.*, Special Syphilis Number, 1912. *Ibid.*, *System of Syphilis*, vol. iv, Diseases of the Nervous System. *Ibid.*, Histological Observations on the Changes in the Nervous System in Trypanosome Infections, especially Sleeping Sickness and Dourine, and their Relations to Syphilitic Lesions of the Nervous System, *Archives of Neurology*, vol. iii. *Ibid.*, Tabes in Hospital and Asylum Practice, *Archives of Neurology*, vol. ii. *Ibid.*, The Changes in the Central Nervous System of Two Cases of Negro Lethargy, *BRITISH MEDICAL JOURNAL*, 1899, p. 1666.
- ¹¹ Neisser, Albert: *Beiträge zur Pathologie und Therapie der Syphilis*.
- ¹² Noguchi: Studien über den Nachweis der *Spirochaeta pallida* im Zentralnervensystem bei der progressiven Paralyse und bei Tabes dorsalis, *Muench. med. Woch.*, April 8th, 1913. *Ibid.*, *La Presse Méd.*, No. 81, 1913.
- ¹³ Noguchi and Moore: A Demonstration of *Spirochaeta pallida* in the Brain in Cases of General Paralysis, *Journ. Exper. Med.*, February, 1913.
- ¹⁴ O'Farrell and Balfour: Granule-shedding in *Treponema pallidum* and Associated Spirochaete, *Journal of the Royal Army Medical Corps*, vol. xvii, No. 3, September, 1911.
- ¹⁵ Robertson, Geo.: General Paralysis of the Insane, Morrison Lectures, *Journal of Mental Science*, April, 1913.
- ¹⁶ Swift: *Studies from the Rockefeller Institute*, vol. xvii, 1913.

THE RELATION OF THE INTERNAL SECRETIONS TO THE FEMALE CHARACTERISTICS AND FUNCTIONS IN HEALTH AND DISEASE.

A PAPER INTRODUCTORY TO A DISCUSSION ON THE SUBJECT
AT THE OBSTETRICAL AND GYNAECOLOGICAL SECTION
OF THE ROYAL SOCIETY OF MEDICINE.

By W. BLAIR BELL, M.D.,

GYNAECOLOGICAL SURGEON TO OUT-PATIENTS, ROYAL INFIRMARY,
LIVERPOOL; EXAMINER IN GYNAECOLOGY AND OBSTETRICS,
ROYAL COLLEGE OF SURGEONS, ENGLAND, AND TO
DURHAM UNIVERSITY.

[Abstract.]

In dealing with the problems under discussion it will be best, I think, first to outline the more purely morphological and physiological aspects. I shall do this briefly, for I have already this year, in the Arris and Gale Lectures,[†] gone somewhat fully into the details of this part of the subject. Afterwards I shall indicate what I believe to be the associated pathological conditions to be recognized, and in some cases the methods whereby they may be treated.

PART I.

MORPHOLOGICAL AND PHYSIOLOGICAL CONSIDERATIONS.

At the commencement it is essential that we consider how far the internal secretions are concerned in producing

the feminine characteristics and ensuring the full development of the genital organs. Afterwards I shall support with experimental evidence my statement that the internal secretions of the ovary alone are not the sole factors in the preservation of the integrity of the genital functions once these have been fully established.

PRODUCTION OF THE FEMALE CHARACTERISTICS AND FUNCTIONS.

What is it that makes a woman a woman? With our present knowledge it appears probable that, although the potentiality to produce femininity exists in the earliest stages of segmentation of the ovum, this potentiality is directed towards the future development and correlations of the endocrinous glands which are to control the sexual evolution of the individual.[‡]

It may be stated most emphatically that the ovaries alone are not responsible for the female characteristics. This is not mere conjecture, for it was first well shown by Bulloch and Sequeira¹⁰ that the suprarenal cortex is largely responsible for the secondary sexual characteristics. Most of the evidence, however, in regard to the influence of the suprarenals has been obtained from pathological phenomena which will be discussed later.

We know less of the actual influence over the secondary characteristics of the other organs of internal secretion, but there is a large body of evidence gradually accumulating which tends to show that the pure female and the pure male are produced by all the internal secretions, probably acting in association with that of the gonads.

We must next consider the influence the endocrinous glands exert on the secondary development of the female genital organs and their functions in the individual perfectly formed in regard to the genitalia and secondary characteristics. This part of the subject is one which in its pathological aspect constantly confronts us in our clinical work.

Although the genitalia may be normal morphologically at birth (primary development), they only become functionally active at puberty (secondary development) if the whole endocrinous system is in perfect harmony and acting efficiently. Thus, thyroid or pituitary insufficiency may cause the genital organs to remain infantile; and, as we shall see later, disease of these structures may cause retrogression in the genitalia even after they have functionated normally. Further, it is known that the gonads and uterus remain underdeveloped until the thymus atrophies at the time of puberty. At this period of life, owing to the withdrawal of the thymus secretion, as some authorities think, the genital organs begin to develop. Other investigators, however, believe that it is the development of the gonads which causes retrogression in the thymus. The probability is that both views are correct; for experimentally it is found that removal of the ovaries leads to hypertrophy of the thymus, even after atrophy has occurred in the normal course of events. Likewise removal of the thymus in male guinea-pigs is said to be followed by a rapid development of the genital glands (Paton).¹⁷

MAINTENANCE OF THE FEMALE CHARACTERISTICS AND FUNCTIONS.

It follows logically that the factors which give rise to femininity and the normal development of genital activity should also be chiefly concerned in the maintenance of the female characteristics and functions. I shall adduce further evidence to show that this is so when we consider the pathological aspect of the subject under discussion. Here the normal processes at work may be described.

That the ovary alone is not solely responsible for the changes which occur or for the integrity of the genitalia is shown by many facts, both experimental and clinical. It is well known, however, that the removal of the ovaries in young animals tends to produce sometimes adiposity and almost invariably overgrowth of the skeleton; the animal is never stunted, nor do masculine characteristics develop. The uterus, however, remains infantile and does not undergo secondary development. So, also, it is certain that if the female genital glands be removed during adult

* The paper will be published in full in the *Proceedings* of the Royal Society of Medicine.

† These lectures may be read in conjunction with this paper, which is in part an attempt to apply the experimental findings to clinical practice.

‡ The word "developmental" may be used in connexion with two periods or stages in the development of the genital organs. The first stage is during fetal life, when the organs are originally formed; and the second is at puberty, when full anatomical and physiological development takes place.

life the uterus retrogresses and becomes atrophied. Some years ago Hick and I¹ showed that the primary change is in the musculature, and not in the endometrium. This change is the obvious evidence of the ovarian function in regard to the uterus. Atrophy of the uterus can, however, be prevented for a time if an ovarian graft—especially an autogenous one—be made when oöphorectomy is performed. It is found experimentally that the follicles in grafts always degenerate, but that the interstitial cells remain unchanged for many months. The success of grafts depends on the technique employed. Thin grafts buried in a vascular site do much better than thick grafts in avascular areas. When finally the graft disappears the uterus atrophies. It is obvious, therefore, that the integrity of the uterus depends, so far as the ovary is concerned, on the interstitial cells. But Crowe, Cushing and Homans,² Aschner,³ Biedl⁴ and others have shown that the same degree of uterine atrophy—and also, it is said, ovarian atrophy—may supervene after partial removal of the pituitary body. Which portion of the pituitary is supposed to be concerned is not clear from the contradictory statements given. It is possible that the divergent opinions result from the belief that the pituitary body consists of two separate organs with different functions. I believe, however, that the pituitary body is *one organ*. In spite of the histological differences seen in the various parts we must remember that all the epithelial elements are derived from the same source—Rathke's pouch.

Further, I have shown that removal of the thyroid alone produces an intense degree of atrophy in the uterus. The ovaries, however, do not retrogress; on the contrary, there appears to be increased activity, especially in the follicles.⁴ In pregnancy, also, we know that the ovaries, the thyroid, the pituitary, and the suprarenals are all concerned in the changes that take place locally and generally. The corpus luteum in the ovary appears to assist in the implantation of the ovum, especially in rodents. This effect apparently is less pronounced in the human subject. The thyroid is especially active in early pregnancy, and the suprarenals and pituitary in the later stages. The changes in the pituitary are shown by very remarkable modifications in the staining reactions of the cells of the anterior lobe. These structural alterations are doubtless essential to the changes in the maternal metabolism, but it is unnecessary here to discuss the chemical details.

In addition to the facts mentioned concerning the local effects of the removal of the ovaries, pituitary, and thyroid on the genital organs and their functions, we have very definite and precise information as to the general effects produced. And since the removal of the ovaries is an experiment some gynaecologists perform on the human subject with regularity and without a qualm—indeed they defend it by saying when they perform oöphorectomy the bad results are not worth considering—I am sure we should give this aspect of the subject special consideration, since we are not all of the same mind as to the needless removal of the genital glands.

In the first place, I would call attention to the well-known fact that women as well as animals vary very considerably in the reactions that occur after oöphorectomy. Age, individual variations in ovarian activity and in the correlations with other endocrinous glands, habits of life, health, and so on, all play an important part in the result.

We might also include psychical effects, because they are related to the functions of the internal secretory system as a whole; but these effects are difficult to estimate in animals, so I shall not discuss them until we come to the consideration of the pathological disorders of the internal secretions in women. Since, then, there are these individual variations, it is somewhat difficult to create a quantitative standard, and this is, I think, the reason of so many divergent opinions in regard to the value of the ovaries. We can, however, recognize the qualitative changes with no small degree of precision.

In animals oöphorectomy produces the following alterations in the general metabolism.* There is a marked retention of calcium, which is, of course, a phenomenon

normally associated with senility. The phosphorus excretion is much increased, as is the nitrogen. These observations I shall refer to again in discussing ovarian insufficiency in the human subject.

When we come to examine the other endocrinous glands after oöphorectomy in animals we find somewhat different changes according to the order to which the animal employed for experimentation belongs. I have elsewhere alluded to the normal morphological differences found in the organs of internal secretion in different animals.⁴ The cat is the most convenient of the commoner animals used in laboratories, in that its endocrinous glands more nearly resemble the human than do those of the rabbit and other rodents. In this animal we find marked changes in the anterior portion of the pituitary and in the thyroid after oöphorectomy. In the thyroid there is a great increase in the colloid collections. In the anterior lobe of the pituitary there is a marked eosinophilia which I have discussed elsewhere, and have described as representing an increase in secretory activity.⁴

We must not consider, however, that our experimental results are final in regard to the human subject until they have been verified by *post-mortem* evidence in women, or by the evidence of disease. Rapid progress to this end would be assured if pathologists would make a routine practice of removing all the organs of internal secretion during *post-mortem* examinations, and of examining histologically the material obtained. Meanwhile laboratory experiments have this effect at least: they give us indications for clinical observation which so often leads to a complete vindication of the laboratory results, and so to settled knowledge.

PART II.

PATHOLOGICAL AND THERAPEUTICAL CONSIDERATIONS.

In considering the relation of pathological conditions of the internal secretions to the genital organs and their functions I shall follow as far as possible the arrangement adopted in describing the physiological processes—that is to say, I shall first discuss the derangements in the normal development of the genital organs and their functions, and subsequently the derangements associated with the maintenance of these functions.

DERANGEMENTS IN THE DEVELOPMENT OF THE GENITAL ORGANS AND THEIR FUNCTIONS.

1. *Imperfect and Irregular Primary Development of the Genitalia.*

Primary maldevelopment of the genital organs may result in the following conditions:

- (a) Under-development of the ovaries and uterus, which may otherwise be perfectly formed.
- (b) Under-development or imperfect development of the uterus with normal development of the ovaries.

The first of these conditions is usually associated with what is known as asexual ateleiosis, the cause of which is due, no doubt, to some general disorder in the endocrinous system. At the present time we have no definite knowledge on this matter, but it is hardly possible that genital under-development with ateleiosis can be due to primary ovarian insufficiency alone in view of our experimental evidence.

The second class of case, in which the ovaries are normal while the rest of the genitalia are imperfectly developed or even assume male characteristics, is interesting if only because some of the conditions met with are liable to be confused with those in the first class.

It is probable, however, that in the presence of normal ovaries imperfect development of the uterus and incomplete union of the Müllerian ducts, with the various associated abnormalities which may be found, are due to some local structural irregularities which are, as yet, not fully understood.⁵ But where the genitalia, apart from the ovaries, assume the male type, and, in fact, give rise to the condition known as hermaphroditismus femininus, we have some definite evidence of the influence of an internal secretion not produced by the genital glands on this condition.

2. *Irregular Secondary Development of the Female Genitalia and Characteristics.*

By secondary development of the genital organs and female characteristics I refer to the changes which

* In his recent work on the organs of internal secretion, under the title of *The Regulators of Metabolism*, Paton calls attention to the fact that only observations conducted over a length of time can have any value, and assumes that none have been made. But, as is evident in various parts of his book, he has not made a first-hand study of the literature, for my experiments, of which he makes no mention, were conducted over far longer periods than he suggests as being necessary.

normally occur at puberty. The onset of the functional activity of the ovaries and the associated secondary sexual characteristics which become evident at this time change the girl into a woman.* Since the advent of puberty itself is dependent on the general metabolism, which is influenced by the internal secretions, we can easily understand how disorders of the endocrinous system influence the phenomena associated with this period of life.

(a) *Precocious Puberty*.—This is an extremely interesting phenomenon, wholly dependent, as far as we know, on disturbances in one or other of the endocrinous glands. It is probable, of course, that a lesion in one organ affects the others, but with this we are not at the moment concerned. In the first place, it is important to remember that the results produced by tumours and hyperplasias in the organs of internal secretion other than the gonads are not the same in the different sexes. So that while suprarenal hypernephromata in male children are practically always associated with precocity of the sexual organs and secondary characteristics, in female children with suprarenal hypernephromata the sexual organs are not precociously developed, for the tendency is rather to produce in them the characteristics of the male—a deep voice, enlargement of the clitoris, hirsuties, and so on. It is true that cases of precocity in females with these suprarenal lesions have been described, but it appears to me that pubic hirsuties with over-development of the clitoris—a masculine transformation—may have been wrongly interpreted into excessive development of the female external genitalia, as was certainly done in Orth's case.¹⁰ Bulloch and Sequeira,⁷ however, report the case of a girl, aged 11, whom they describe as precocious, and in whom menstruation existed. But I do not think the evidence submitted is sufficient to justify the statement as to female precocity, especially in view of the facts that she only commenced to menstruate at the age of 10, and that subsequently she had much hair on the lips and chin.

It is probable, therefore, that the view I have stated is the correct one; indeed, I fail to understand how it can be otherwise when we consider the evidence at our disposal as to the masculine metamorphosis that occurs with these same lesions in the suprarenals in young adult women. Most of the cases of suprarenal hypernephromata observed in children have occurred in girls. In the case of a boy, who died at the age of 5, recorded by Linser,¹³ there was precocious development of the genital organs with sexual activity. Likewise it is interesting to note that tumours of the pineal in boys always lead to sexual precocity, but do not do so in girls.

It is difficult to interpret this curious sex selection in regard to these lesions in different organs, but certainly it appears at first sight as if both the pineal and suprarenal cortex influenced the production of male characteristics. We must remember, however, that we have no reliable evidence as to whether tumours of the pineal give rise to excessive or decreased secretion, but the above suggestion does not necessarily depend on an increased secretion from the pineal in the circumstances mentioned.

Although the foregoing facts concerning the suprarenal cortex and the pineal are negative in regard to sexual precocity in girls, I have dwelt on them as it is necessary to clear the ground in this matter, for most clinicians seem to be under a misapprehension, since they conclude that corresponding lesions produce sexual precocity in either sex. This, as I have endeavoured to show, is not so except in regard to lesions in the gonads. Indeed, to come to the point, it appears probable that nothing in regard to lesions in the endocrinous glands except certain tumours and hyperplasia in the ovaries will give rise to true sexual precocity in the female. Many such cases are on record. A typical example is that related by Lucas:¹⁴ A girl, aged 7, had markedly precocious development with menstruation. A solid ovarian tumour of doubtful nature was removed, and the signs of puberty thereupon gradually disappeared. It follows, therefore, that unlike sexual precocity in boys, this phenomenon in girls is only associated with tumours or hyperplasia of the gonads.

(b) *Delayed Puberty*.—Delayed puberty may be due to a variety of causes, such as bad feeding, unsuitable occupation, chlorosis, and so on, in which it is difficult to trace a

primary cause in the internal secretions; at any rate, one need take no account of them in the treatment of such cases. But there is a large number of instances in which the organs of internal secretion are primarily at fault. As already stated, the internal secretions of the ovary mainly produce their effects in correlation with other internal secretions; and the full development of the genitalia, therefore, is dependent on the general metabolic conditions prevailing under the influence of all the endocrinous glands. So that, strictly speaking, one cannot consider under-development of the ovary as a primary factor in delayed puberty, although, undoubtedly, it may be found to be under-developed as a correlated condition. Practically, it appears that the thyroid and pituitary, in association with the ovaries, are the factors most concerned in the final development of the female genital organs. It will easily be understood how important a matter it is that all cases of delayed puberty—both in major degree where the patient remains childlike altogether, and in the minor degree where she is well developed in every particular except in regard to the genital functions—should be carefully taken in hand at the earliest possible moment. These patients are frequently obese, consequently this condition has been considered to be a primary cause of delayed puberty. This is not so; both conditions are dependent on the same cause.

Many cases can be treated successfully with thyroid extract, and some with pituitary extract in large doses. Extract of the whole gland should be prescribed.

There appears to be little chance of effecting full development of the uterus after the age of 20 years. This, at first sight, appears to be a somewhat curious fact, for one might think that if a uterus can be, and normally is, roused into activity after fifteen years of primary inactivity, ten more years of quiescence would make no difference. It must be remembered, however, that the body metabolism alters rapidly after the eighteenth year of life, when the development of the skeleton is practically completed: so that, unless genital activity be aroused during the period of change, it is impossible sufficiently to control the metabolism, which has become rearranged, in order to produce the effect desired.

DERANGEMENTS OF THE FULLY ESTABLISHED FEMALE CHARACTERISTICS AND FUNCTIONS.

Derangements of the Sexual Characteristics.

The secretions of the endocrinous glands which have produced the normal female continue subsequently to act in the same way—passively, no doubt—and it is only some extensive alteration in the structure and function of certain units of the internal secretory system which can bring about an alteration in the female sexual characteristics once they have been determined, and the genital organs have become functionally active. So far as we know, the withdrawal of any one secretion will produce no change. There must be some overgrowth in those parts of the endocrinous system, apart from the genital glands, which normally produce masculinity—a potentiality which appears to be concentrated in the suprarenal cortex, the pituitary body, and probably in the pineal. Apert¹ has asserted that masculine characteristics may be produced in women by simple ovarian tumours. But, as Glynn¹¹ points out, there is no evidence in the cases reported that the suprarenals were examined, or, I might add, the pituitaries and pineals.

In adult women, before the menopause, the changes towards masculinity associated with suprarenal hyperplasia and neoplasia may be most marked. Menstruation ceases, the breasts shrink, hair grows on the face, the voice deepens; and last, and not least, profound psychological changes have been known to occur, a previously gentle woman becoming rough and aggressive. A number of such cases have recently been collected by Glynn.^{11 11a}

With acromegaly, too, I have seen a somewhat similar change. In one case menstruation ceased, the skin became coarse, the voice deep, the breasts shrunken, and last, and most interesting of all, the clitoris hypertrophied until it resembled a small penis. We know that in the male acromegaly (hyperpituitarism) at first gives rise to increased sexuality, so that the reverse obtains in the female in whom there is a tendency towards masculinity. In the case of the internal secretions of the pituitary, therefore, we see evidence of a difference of sex stimula-

* It has already been mentioned that the uterus may be malformed owing to local pelvic conditions, with full development of the ovaries. In such circumstances puberty occurs, although the uterus may be functionless.

tion in the male and the female, just as we saw it in connexion with the suprarenals and pineal.*

These, then, are the main facts in connexion with the maintenance and alteration of sex characteristics in the adult female.

Derangements of the Genital Functions.

In considering derangements of the fully established functions I shall take in order each of the chief organs of internal secretion and endeavour to show how they are individually related to the various disorders met with. In some cases we have to take into account certain correlations; especially is this so where the ovary is concerned.

Primary disorders of the endocrinous organs may be considered as giving rise to insufficiency or excess of the normal secretions.

Ovarian Insufficiency.

Ovarian insufficiency occurring during the active sexual period is rare as a primary disorder, except when produced by operative interference. As we have already seen, when the ovaries are removed experimentally definite changes occur in the general metabolism, in the uterus and in the structure of the other organs of internal secretion; and these changes appear to be more prominent in the higher than in the lower mammals.

From observations made on the human subject there is no doubt that a similar train of events follows complete oöphorectomy. In addition, however, there are frequently to be noticed psychical phenomena, which are doubtless dependent on the general disturbance of the metabolism; but marked individual variations exist in the symptoms produced just as normally there are considerable variations in the femininity and sexuality of different women. Unfortunately, those who have asserted that oöphorectomy is of small moment to the patient have failed to make good their scientific position in the matter by the exercise of a discretion that appears obvious in view of what has been said regarding individual variability. If one were to gauge, and in many cases it would not be difficult, the degrees of femininity and ovarian activity in the patient, it might be possible to assert that in those cases in which these were obviously in abeyance little disturbance of the metabolism would result; and, consequently, it might be justifiable in such cases to remove the ovaries with a fibromyomatous uterus or with infected tubes. But unless such a scientific distinction, which undoubtedly does exist and accounts for the individual variations after oöphorectomy, be made, it does not seem right blindly and systematically to remove the genital glands.

It is more or less generally supposed that the natural menopause occurs in women as the result of an ovarian insufficiency. Undoubtedly there is ovarian insufficiency sooner or later, but it has always seemed to me that the symptoms produced depend for their severity on the correlations that exist between all the internal secretions, on the uterine changes which prevent the monthly excretions, and, lastly, on individual variability as to sexuality and to the stability of and capacity for readjustment in the endocrinous system. That the phenomena associated with the natural menopause partly depend on ovarian insufficiency is proved by the onset of similar manifestations in most women after oöphorectomy. That they are related to the coincidental uterine atrophy is proved by the fact that women may suffer from these same symptoms after removal of the uterus alone. Also, I have seen monthly menses which reminded one strongly of some of the symptoms of the menopause in women in whom there was congenital absence of the uterus or malformation sufficient to prevent menstruation, together with functional ovaries. Finally, the natural menopause varies in its severity in different women, just as does the artificial; but probably chiefly according to the extent of the femininity, which is accentuated by ovarian activity.

The main types of menopausal phenomena to be observed clinically may be classified as follows:

1. Psychical.
2. Vasomotor.
3. General metabolic.
4. Gross changes in the endocrinous glands.

* The thymus appears to take no part in the regulation of sex characteristics, in spite of its supposed influence over the formation of the skeleton. Further investigations may, however, prove that it is so concerned.

Thus we may observe:

1. Mental irritability or instability.
2. Flushing with sweating, cardiac distress, and intestinal movements.
3. The deposition of fat and the retention of calcium salts in the tissues.
4. Excessive changes in the endocrinous glands, especially the thyroid and pituitary body. One of my patients died of exophthalmic goitre one year after the removal of both ovaries for malignant disease. Myxoedema, which recent views associate with previous excess of secretory activity in the thyroid, also usually occurs after the menopause. The changes in the pituitary are less noticeable, but the deposition of fat may be the result of what is known as carbohydrate tolerance, due to hypopituitarism. This condition is always seen in the disease known as dystrophia adiposo-genitalis.

It must be remembered, of course, that the phenomena mentioned above are classified clinically. They are all primarily due to disturbances in the endocrinous organs.

The information required before we can speak with certain knowledge about the changes at the menopause is considerable. Work must be done on the normal age changes in the endocrinous glands, and especially the changes that occur at the menopause; and the metabolism before, during, and after the menopause must be worked out in a number of cases and compared. But however little we know about the metabolism of, and the structural changes in, the glands of internal secretion at the natural menopause, we recognize the fact that the more gradual the onset and completion the less the general disturbance. Many women, of course, have prolonged and distressing menopauses, but in some, at least, I am inclined to think that the menopausal retrogression is spasmodic rather than continuous. The cases of the natural menopause in which the disturbance is great are undoubtedly those which occur suddenly, and for this reason the artificial menopause is liable to be more serious than the natural.

There seems, too, to be some relation between the severity of symptoms and the age of the patient. In this respect, I believe, the natural often differs from the artificial menopause. When it occurs in young women (35 to 40 years of age) naturally, or pathologically as in superinvolution of the uterus, the symptoms are usually slight. Possibly this is because the genital functions in such women have never been predominant. When the natural menopause occurs in elderly women (50 to 55) the symptoms are often severe. On the contrary, the artificial menopause usually produces far worse effects in young women than in those over 45 years of age, the degrees of femininity being equal.

To complete the picture of ovarian insufficiency it is, necessary that we should turn for a moment to the question of the ovarian secretions in pregnancy. It was first shown by Fränkel¹⁰ that destruction of the corpus luteum in the early stages of pregnancy inevitably caused abortion in animals. Hick and I⁵ and others have confirmed the importance of the ovarian secretions in the implantation of the ovum in rabbits. But this does not hold good to the same extent in the human subject, for both ovaries have been removed as early as the sixth week of pregnancy without producing abortion. In such cases the question of the entire removal of the ovary is easily settled by the subsequent history of the patient. At the same time it is almost certain that the initial implantation of the ovum could not occur in the absence of the secretion of the corpus luteum. And, indeed, such insufficiency may be the cause of many early abortions.

It has been suggested that during pregnancy ovulation does not occur, and that the internal secretions of the ovary, apart from that of the corpus luteum, are in abeyance. I do not think there is any real evidence in support of these statements. Nevertheless, if such were the case, it might indicate that ovarian insufficiency during pregnancy threw a considerable strain on the other organs of internal secretions such as the thyroid and anterior lobe of the pituitary, and, in fact, might be the original cause of some hyperplasias in these structures.

When we consider the treatment of ovarian insufficiency, the question of ovarian transplantation naturally first crosses our minds. Experimental and operative evidence goes to show that autogenous grafts alone are of service. In rabbits, grafts from one animal to another have been

shown to be effective, but only for a short time. Professor Tuffier of Paris has found that, in the human subject, only autogenous grafts are of any use at all. Ovarian transplantation, therefore, can only be employed in mitigation of the artificial menopause. That is to say, autogenous ovarian grafts may be of value in women under 40 years of age in whom both ovaries have been removed; but they can be employed only in those cases in which the lesion is not malignant. Thus there is a very restricted field for this method of treatment, especially when we remember how rarely it is necessary entirely to sacrifice the whole of both ovaries in other pathological conditions. In the circumstances indicated, however, I believe it to be a most valuable and even imperative procedure. When a graft is made, it is better, I think, to use a thin, flat, or wedge-shaped piece of ovary without the cortex, in order that the extensive raw surfaces may quickly obtain attachment and an efficient blood supply be established, and that by the removal of the cortex cyst formation in the follicles, which is otherwise inevitable, may be avoided. Recently I saw Professor Tuffier of Paris implant an ovary. He simply buried the whole ovary in the anterior abdominal wall. This appears to me a very bad technique, for the reasons indicated above. In any case, as far as we know, the menopause is only postponed and thereby mitigated by ovarian implantation.

Excessive Ovarian Secretion.

The clinical effects produced by excessive ovarian secretion may be discussed under two headings:

1. Effects on the sexual functions, characteristics, and psychology.
2. Effects on the general metabolism.

It must not be forgotten that the effects mentioned in the first group are chiefly dependent on alterations in the general metabolism.

1. An excessive ovarian secretion leads to an increase in sexual activity both locally and generally. The local effects may be seen in an increased and prolonged menstruation. This condition occurs chiefly in girls about and just after puberty who indulge in masturbation or are sexually insane, in unmarried women over 30 years of age, and in women whose married life is a failure from a sexual point of view.

Before long it is probable that we shall find that injections of suprarenal and pituitary extract may be employed to control the effects of excessive ovarian secretion, and in these circumstances operation will not be called for.

2. An excessive ovarian secretion affects the metabolism in the opposite way to oöphorectomy, and one of the most important metabolic disturbances is the abnormally large excretion of calcium salts. In the non-pregnant woman this is a matter of no importance so far as the skeleton is concerned, but it possibly accounts for some of the nervous and muscular irritability, which are constant features of these cases. In pregnancy and lactation, however, there is a large demand for calcium over and above the amount necessary for the normal maternal metabolism, and, as a result, softening of the bones may occur. This condition is known as osteomalacia. Such cases are not common in some localities, and personally I have not seen a case for many years. This disease used to be treated by removal of the ovaries, and many patients were cured. A few years ago Bossi, of Genoa, suggested the injection of suprarenal extract as an alternative to oöphorectomy. Apparently many cures have been effected by this treatment, and also by the injection of pituitary extract. So that it may be argued that there is insufficiency in these organs, as well as an excess of ovarian secretion; and this, indeed, is quite possible. Be this as it may, substitution of other extracts is a great advance on removal of the ovaries in the treatment of osteomalacia; and the success thus obtained indicates the line of attack that will probably be followed in the future in regard to all those diseases of the endocrinous glands in which there is an excessive secretion. That is to say, metabolically antagonistic extracts will take the place of removal of portions of the diseased organs.

By way of further illustration I may here mention that, as I suggested some time ago, the injection of pituitary extract is an excellent method of treatment in hyper-

thyroidism. Suprarenal and pituitary extracts counter-balance the action of the ovaries and thyroid and lead to calcium retention, which, as we have already seen, is a normal function of the suprarenals and pituitary. It is probable that both portions of each of these structures are concerned in the result.

Thyroid Insufficiency.

Thyroid insufficiency always causes a decrease in or the complete cessation of the function of menstruation, according to the degree of insufficiency. I have already alluded to the complete atrophy of the uterus which follows thyroidectomy in animals; and it will be remembered that I pointed out that in my experiments the ovaries were not found to undergo atrophy, but rather to show increased activity. Apparently in the human subject atrophy of the uterus is never complete with pathological insufficiency of the thyroid, not even with myxoedema; for I have known a patient, who had had this condition for some years, not only to menstruate, but to conceive after the administration of thyroid extract.

It is, however, the disturbances arising from minor degrees of thyroid insufficiency which chiefly come under our notice as gynaecologists. Delayed menstruation has already been discussed. Secondary amenorrhoea and scanty menstruation, the latter often associated with dysmenorrhoea, are very frequently seen with thyroid insufficiency. The patients are obese, lack energy, and have deficient sexuality. There is another menstrual disturbance—namely, mastodynia—which sometimes occurs regularly each month before the onset of menstruation, in which I have found thyroid extract of great service; and in consequence of this I believe that the condition may be indirectly connected with thyroid insufficiency. If thyroid extract be given from about the twenty-fourth day of the cycle till menstruation is established this function may come on earlier than usual, with relief of the pain in the breasts. There is a natural aversion from prescribing thyroid indiscriminately, but in these minor degrees of insufficiency thyroid extract is the one preparation which will cure the patient. I have seen quite remarkable results in a number of cases. I always order the preparation—usually 3 grains of the dried gland—to be taken only on going to bed at night. If given in this way and not during the day the patient never suffers from ill effects.

With regard to pregnancy, it is well known that normally the thyroid gland enlarges in the early stages, but I need not here go into the causes of this. It shows that increased thyroid activity, which, I believe, takes the form of a storage of colloid, is required. A marked degree of insufficiency, such as is seen in myxoedema, almost invariably leads to sterility. Cases have been recorded in which pregnancy has occurred in minor or variable states of this disease; and in such circumstances a notable improvement may be produced in the general condition, the thyroid becoming more or less active.

There is one more point in regard to thyroid insufficiency and pregnancy which requires some comment. Nicholson¹⁵ promulgated the view, which has been taken up by many others, that eclampsia was due to thyroid insufficiency in pregnancy. I conducted a series of experiments—reported elsewhere¹⁶—to find out how far this was true; and I came to the conclusion that neither metabolically nor actually was there any real evidence in support of Nicholson's view. Indeed, it appeared to me that the animals in the later stages of pregnancy were considerably less affected than those which were not pregnant. I have accounted for this curious fact by assuming that in the later stages of the development of the fetal thyroid its secretion may be conveyed to the mother. The thyroid, of course, has a most important part to play during pregnancy, but I do not think it has anything to do primarily with the causation of eclampsia.

Excess of Thyroid Secretion.

It is a common experience to meet with an excess of thyroid secretion in connexion with pelvic lesions, and we must be careful to distinguish between those cases which are the result of genital affections and those which are the cause of them. It is certain that after the menopause the thyroid normally undergoes some retrogression, after, perhaps, a period of minor hyperthyroidism; and it is probable that removal of the ovaries leads, eventually a

all events, to the same condition in the majority of cases. But, as I have mentioned, acute exophthalmic goitre may intervene. Experimentally it is seen that a considerable increase in the colloid formation follows oophorectomy, but it is probable that this is a different condition from Graves's disease. At the same time we really know far too little about exophthalmic goitre—especially in its relation to simple goitre and to myxoedema, with which it not infrequently alternates—to speak definitely on this question. Excessive thyroid secretion may act on the genital functions in two ways. First, it may stimulate them to unusual activity, or, secondly, the general metabolism may be so upset that the genital functions cease. I am quite certain that there is a condition of hyperthyroidism which produces excessive menstruation, and which is totally unassociated with any of the symptoms of exophthalmic goitre. This condition is usually seen in girls about puberty usually just before and during menstruation. It may be that the excessive activity is due to the incomplete functional development of the ovaries; but, whatever the cause, there is not infrequently menorrhagia. I have, however, seen this same enlargement of the thyroid with amenorrhoea. This paradoxical phenomenon is easily explained on the supposition of ovarian insufficiency, and an inability of the thyroid in some cases, in spite of its enlargement, to meet the deficiency. When there is menorrhagia I have found that it can always be controlled by calcium lactate.

The relation of excessive thyroid secretion to conception and pregnancy is very interesting, and much has been written on the subject. I think there can be no doubt that excess of thyroid secretion, except that seen in connexion with exophthalmic goitre, does not interfere with fertility. It would be strange if it did, since we know there is an increase of thyroid activity during the early months of pregnancy, and possibly throughout. But it is no less a fact that women with marked exophthalmic goitre rarely become pregnant, and if they do the disease is aggravated. I remember one patient who had suffered from exophthalmos for some years, yet she had had several children, but there had always been haemorrhage *post partum*—an occurrence which has frequently been noted in these cases. She consulted my colleague, Mr. Thelwall Thomas, and he removed one half of the thyroid gland, with the result that there was marked amelioration in her symptoms. Shortly afterwards she became pregnant again, and Mr. Thomas referred her to me for observation and treatment. She became somewhat worse than she had been since her operation, so I gave large doses of calcium lactate and ordered rest. She went comfortably to full term, and there was no haemorrhage *post partum*. I advised her not to nurse the child, which was normal. She improved very considerably after parturition, and is now, so far as I know, in good health. I think it is most important that these patients should be given large doses of calcium salts if they become pregnant, as there is such an excessive excretion of lime salts in this disease, and so much danger of haemorrhage *post partum* and even *ante partum*. Infundibulin should be given immediately after labour, to prevent bleeding.

Pituitary Insufficiency.

We have seen that, like the thyroid, the pituitary is actively concerned in the proper development and the maintenance of the genital functions. Consequently, as we would expect, insufficiency of secretion is associated with amenorrhoea or scanty menstruation. I have already mentioned the fact that various investigators claim for different portions of the pituitary body different functions, and I have stated my belief in the unity of the whole gland. Patients with pituitary insufficiency are always obese and dull, and it is often a difficult matter to say whether the thyroid or the pituitary, or both, be at fault. Sometimes it is quite impossible to make sure, and in such cases one can only try first one and then the other extract. I have occasionally given both together, for we know from experiments that a very intimate connexion exists between these two organs. If it be true that there is atrophy of the ovary with pituitary insufficiency, as described by Cushing, it would probably be useful to give both pituitary and ovarian extract together. I have shown that ovarian atrophy does not occur with thyroid insufficiency in adults, and therefore I am somewhat sceptical as to the findings of Crowe and his co-workers,

already alluded to, in regard to the ovaries with pituitary insufficiency. Sometimes the condition of the skin may help one in forming an opinion as to which gland is at fault. If it be dry and rough one would suspect the thyroid, if fine and smooth it is more likely that the pituitary is at fault. I have come across no case recorded in which a patient suffering from a major degree of pituitary insufficiency has become pregnant. As already mentioned, pituitary extract appears to act as well as suprarenal extract in osteomalacia, so that it is possible that there is a minor degree of insufficiency of the pituitary body in this disease.

Excess of Pituitary Secretion.

We have seen that an excess of pituitary secretion, such as occurs in acromegaly, produces masculinity, and while in the male this may give rise to excessive sexuality, in the female it causes amenorrhoea and loss of desire, as is only natural if masculinity be produced. The amenorrhoea may be spasmodic, for acromegaly tends to improve for a time and then again to retrogress. Cushing⁹ has shown that eventually acromegaly (hyperpituitarism) changes into dystrophia adiposo-genitalis—a condition of pituitary insufficiency. The effect of excessive pituitary secretion must, therefore, necessarily tend to produce sterility. A case, however, has recently been recorded¹² in which a woman suffering from acromegaly conceived after having been treated, before becoming pregnant, with ovarian extract, which was believed to have improved her general condition and to have caused the appearance of menstruation. I have myself seen an improvement with the reappearance of the menses take place in acromegaly under ordinary circumstances. So that it is difficult to say with certainty that pregnancy resulted in the above case from the treatment with ovarian extract.

Suprarenal Insufficiency.

This condition has only received clinical recognition in Addison's disease, in which there is an extensive derangement of the metabolism. The amenorrhoea that is found in this disease is probably due partly to the general disturbances and partly to uterine atrophy. I know of no record of a proper *post-mortem* examination of the genital organs in this disease. Experimentally I have found a moderate grade of uterine atrophy with suprarenal insufficiency.

We have already seen that the administration of suprarenin has been successfully employed for osteomalacia. This, as I have pointed out, may be not only because suprarenin is antagonistic to the ovarian secretion, but also because in this disease there may be suprarenal insufficiency, a supposition which is supported by the fact that in one case in which I produced experimentally suprarenal insufficiency in a rabbit the bones of the fore-legs became bowed. Further, in most of the animals operated on there was an enormous increase in the urinary excretion of calcium. It appears, therefore, that the secretions of the suprarenals are of great importance during pregnancy in assisting the absorption and retention of lime.

Excess of Suprarenal Secretion.

There is little to add to what has already been said concerning the sex changes that occur in adult women with suprarenal cortical hyperplasia. Amenorrhoea is a constant phenomenon, and pregnancy never occurs in well-defined cases. At present we have no remedy for suprarenal hyperplasia, but in the future some treatment may be forthcoming for the minor degrees of this condition, and this will probably be of the nature of an antagonistic combination of secretions. That there are many cases of slight hyperplasia which go unobserved seems probable, for it is a question whether the shame of a moustache or pride in masculinity has the greater weight with some women. I have such a case under observation at the present time. The patient is aged 19, her menses have ceased, and she has a masculine voice and appearance. The growth of hair on the face alone is required to complete the picture.

Thymus Insufficiency.

This, so far as we know, is natural after puberty. It is possible that in cases of early retrogression of the

thymus sexual precocity occurs. It is difficult, however, to know which is the cause and which the effect. Further clinical investigations should throw light on this question.

Excess of Thymus Secretion.

In the condition known as status lymphaticus there is abnormal persistence of the thymus, and since the secondary or pubescent development of the sexual organs is associated with retrogression of the thymus there is under-development of the genital organs with amenorrhoea if the thymus persist. Individuals with a persistent thymus usually die early, and I do not think the condition has as yet been considered at all seriously from the genital point of view. Retrogression of the thymus might be assisted, if the view be true that development of the genitals causes retrogression of the thymus—a question which has already been discussed.

REFERENCES.

- ¹ Apert, *Bull. méd.*, Décembre 21st, 1910. ² Aschner, B., *Wien. klin. Woch.*, 1909, xii, p. 1730. ³ Bell, W. Blair, *Proc. Roy. Soc. Med.*, July, 1909. ⁴ Idem, Arris and Gale Lectures, *Lancet*, March 22nd and April 5th, 1913. ⁵ Bell, W. Blair, and Hick, P., *BRITISH MEDICAL JOURNAL*, March 13th, 1909, p. 656. ⁶ Biedl, A., *Innere Sekretion*, Berl., 1910. ⁷ Bulloch, W., and Sequeira, J. H., *Trans. Path. Soc.*, 1905, lvi, p. 189. ⁸ Crowe, S. J., Cushing, H., and Homans, J., *Bull. Johns Hopkins Hosp.*, 1910, xxi, p. 127. ⁹ Cushing, Harvey, *The Pituitary Body and its Disorders*, 1912. ¹⁰ Fränkel, L., *Arch. f. Gyn.*, 1905, lxxv, p. 224. ¹¹ Glynn, E. E., *Quart. Journ. Med.*, 1912, v, No. 18, p. 156. ¹² Glynn, E. E., and Hewelson, J. T., *Journ. Path.*, 1913, xviii, p. 81. ¹³ Kelladay, L., *Zentralbl. f. Gyn.*, 1913, xviii, p. 1. ¹⁴ Linser, P., *Beitr. z. klin. Chir.*, 1903, xxxvii, p. 282. ¹⁵ Lucas, R. C., *Trans. Clin. Soc.*, 1888, xxi, p. 224. ¹⁶ Nicholson, H. O., *Scot. Med. Surg. Journ.*, June, 1901. ¹⁷ Orth, *Arbeiten aus den Path. Institut zu Göttingen*, Berl., 1893 (quoted by Bulloch and Sequeira). ¹⁸ Paton, Noel, *Journ. Physiol.*, 1904, xxxii, p. 59.

EPIDEMIC CERVICAL ADENITIS.

BY

J. GORDON SHARP, M.D. EDIN.,

LEEDS.

DURING the spring we had in Leeds a long-continued epidemic of influenza of a moderately severe type. Side by side with this there had been in some districts numerous cases of mumps, diphtheria, and measles, but amongst the cases of illness there were some which could not be fitted with a diagnostic label. The obscurity attaching to them is now removed to some extent by the descriptions given in the article which appeared in the *JOURNAL* of June 14th, 1913, p. 1282, entitled "An Epidemic characterized by Adenitis, Tonsillitis, and Cardiac Complications."

I cannot say how many cases I have seen—perhaps twenty, thirty, or more in all—for at first they fell into that long list of undescribed conditions which we all meet with; and it was only when some half-dozen cases of more than ordinary severity were seen that one took special notes with a view to arriving at an exact diagnosis. These were as follows:

CASE I.—*Cervical Adenitis and Tonsillitis.*

The first case that arrested my attention was a girl of 12, who in April, 1913, had a swelling at the angle of both jaws and in the triangles of both sides, with a red rash, limited to the neck, which completely disappeared after half a day. There was slight swelling of both tonsils, an oedematous uvula, a white-coated tongue, and a temperature which never was higher than 101°, a pulse of 110, stiffness of the neck and shoulder muscles, malaise, nausea, complaint of palpitation and pain at the pit of the stomach. One is almost ashamed to write that the diagnosis varied between rheumatism, influenza, German measles, mumps, and diphtheria. Rheumatism because a brother and sister suffered from heart disease, the result of this affection, and because of the stiffness from which the girl herself suffered; influenza because it was prevalent; German measles because of the rash; diphtheria because it, too, was then in evidence, but a swab cleared this up; mumps because three cases had occurred in the near neighbourhood. All were excluded and the diagnosis was left blank. The most prominent signs and symptoms were the cervical adenitis, which persisted for more than a week after the acute symptoms; the rapid heart action, which persisted for three weeks; and the later debility.

CASE II.—*Cervical Adenitis, Measly Rash, and Tonsillitis in an Adult.*

This patient, seen in April, 1913, was a lady of 35, who had a large swelling at the angle of the left lower jaw, and the same on both sides of the sterno-mastoid muscle, accompanied by a measly rash on the neck, chest, and hands, which appeared suddenly and disappeared suddenly within two days. The tongue was coated white and the tonsil on the left side was large and purple red. There was great stiffness of the muscles

of the neck. The pulse was 120, and palpitation was complained of. The temperature rose to 103° on two days. There was nothing suggesting quinsy. One morning she said she was sure something had burst in her mouth, and she complained of a salty taste for two or three days. After this the tonsil decreased in size and the swelling on the jaw also disappeared. The stiffness remained long after all acute symptoms had gone. The swelling of the glands along the sterno-mastoid muscle did not go till four weeks after its first appearance, and convalescence was not established till this time. The exhibition of the hypophosphites and Fowler's solution seemed to hasten the disappearance of the enlargement.

CASE III.—*Mild Cervical Adenitis.*

A boy of 7 was suddenly seized in March, 1913, with a temperature of 101°, headache, slight sore throat, and enlarged glands on both sides and in both triangles. He got rapidly well on 5-gram doses of sodium salicylate, but the glandular enlargement remained for over fourteen days, much to the anxiety of the parents, who feared acute tuberculosis. The boy ultimately got quite well.

CASE IV.—*Adenitis with Moderately Severe Tonsillitis.*

A girl of 11 had stiff neck, enlarged glands at the angle on both sides, swelling in both anterior triangles, enlarged tonsils, especially on the left side, and on this side there was a patch of yellow which it was thought might be diphtheria. A swab gave a negative result, and the child soon got well as far as the acute symptoms were concerned. The temperature was never higher than 102°. The glands in the anterior triangles did not go away till the girl was up and moving about.

CASE V.—*Very marked Glandular Enlargement with Severe Constitutional Symptoms.*

A girl of 13 was taken ill with headache, stiffness of neck and shoulders, temperature of 103°, sickness, vomiting, large swelling at the angles, in the triangles, tender tonsils, and oedematous uvula. Under time and treatment the mouth symptoms, swellings at the angles, and the sickness and vomiting cleared away, but the swelling of the glands remained in the triangles till the child was able to be out—some three weeks after the commencement of the illness. Earlier in the year four brothers (she was the only girl in the family and slept alone) and the father and mother were all laid up with influenza.

CASE VI.—*Enlarged Cervical Glands, Tonsillar Abscess, and Palpitation.*

A woman of 25 was seized in May, 1913, suddenly with pain in the throat, stiff neck, enlarged glands at the angles and in both triangles, epigastric pain, suffocating feeling, palpitation, and a temperature of 103° in the armpit. On examining the mouth the tonsils were swollen and red, but more particularly on the left side (by a strange coincidence the left side is nearly always the more severely affected), which was bulging. On the third day this swelling burst and discharged pus and blood, and left a ragged opening. The heart was 140, and the first sound in the mitral area was not clear; but this was not seriously considered, for the patient is somewhat "nervous." This may have been a mistake on my part, and the heart condition may have been part of the general disturbance. The swelling at the angles of the jaw rapidly subsided, but in the case of the glands along the sterno-mastoids the subsidence only took place gradually and only after a few weeks. The convalescence was prolonged.

In all the cases the tonsils played an important part, and they may have been the seat of entry of infection. I can now recall several cases of a mild nature having occurred in families, so that the disease may be of a nature which is communicable from person to person; and I now see cases in which stiffness along the sterno-mastoids is complained of, and on examination tender glands are discovered near the muscle. Are they late cases of the kind now under review? They disappear on being treated with organically combined iodine and massage.

THE will of the late Mr. Walter Whitehead, consulting surgeon to the Manchester Royal Infirmary, has been proved at £176,892 net. Among the bequests are £250 to the North Wales University College, and an annuity of £20 to the Manchester Royal Infirmary for the Bradley scholarship.

WE learn from the *Australasian Medical Gazette* that in order to promote research in cancer Mr. John Grice, a member of the Melbourne University Council, has offered a sum of £1,000, to be spent at the rate of £125 a year, to establish a scholarship for work at the University and the Melbourne Hospital. The State Government makes an annual grant of £1,000 to the University for research purposes, and the Minister of Education recently conferred with Dr. MacFarland (the vice-chancellor) and Dr. J. W. Barrett, and agreed to raise the amount to £1,125 for three years, so that the extra £125 could be devoted to cancer research during that period.