was not detected in the routine examination of the urine, and the diagnosis was only made *post mortem*.

High arterial tension and consequent changes in heart and arteries have usually been present in these cases, and have been regarded by some as constant. They were not recognizable in a case recorded recently,³ and in Cases I and II above they were not present to any appreciable degree clinically, in spite of the fact that they were carefully sought for. There were some signs of retinal changes in Case I, and microscopically the walls of the arterioles of the kidneys were distinctly thickened; but considering the extreme degree of fibrosis of the kidneys and their minute size, in conjunction with the comparatively long duration of the symptoms, any changes present would seem unaccountably slight. Dr. Sawyer⁴ suggests one year as an average duration of symptoms in similar cases, and notes three years in the exceptionally longer ones; but this boy had obvious symptoms, which hardly varied at all over a period of seven and a half years.

Pathologically the condition appears to be similar to other forms of interstitial nephritis, although the capsule of the kidney was not adherent in either fatal case.

In diagnosis, when the symptoms are obvious, one would suppose that cystic disease of the kidneys would be the only condition to exclude.

That so rare an affection should be met with in two children of the same parents is of extreme interest from an etiological point of view. It would seem to confirm the view that such cases may be due to some intrauterine infection, or that heredity plays some part. In these particular cases neither the results of the Wassermann tests nor any other findings would seem to indicate syphilis. In Dr. Sawyer's cases the affection was more frequent in girls. I do not know whether the suggestion has been made that a *Bacillus coli* infection might produce this disease. The right kidney in Case I was more affected than the left, which has been noticed in *coli* infections. There was no dilatation of the pelvis or ureters, which, present in some cases, has suggested gravel or calculi as a disposing cause.

I am indebted to Mr. Luce for leave to publish Case III, which I only saw *post mortem*.

REFERENCES.

¹ Allbutt and Rolleston's *System of Medicine*. ² *Proc. Roy. Soc. Med.*, Dr. Miller, December, 1911. ³ *Ibid.* ⁴ *Chronic Interstitial Nephritis in Children*, *Birmingham Review*, 1903, Dr. J. E. H. Sawyer; also *St. Thomas's Hosp. Rep.*, 1906.

YEAST INFECTION IN CHILDREN.

BY

ROBERT CRAIK, M.D. GLASG

EALING, W.

AN infant, 4 months old, was said to be always crying. It was a big child, but thin, pale, and peevish-looking, and sucking a comforter. It had been ailing for about two months. There was a history of frequent motions, which were seldom watery, but often jelly-like. The mother's milk had been scanty from the beginning, and after the first fortnight breast milk was given at night only. During the day a mixture of barley water and cow's milk was used. Examination revealed little beyond an inflammation of the perineal region. The mouth showed no sign of stomatitis.

A swab from the tongue and a loopful of freshly-passed motion were smeared on malt extract agar. After twenty-four hours, at 37° C., the latter showed confluent colonies of the thrush fungus along its whole length. Only a few discrete colonies appeared on the smear from the mouth. Appropriate treatment soon gave a satisfactory result.

Stomatitis due to the thrush fungus is familiar to every one, and even a marked attack may not greatly upset an infant. The source of infection is the mouth of the mother or nurse, and a comforter is often the carrier. Along with stomatitis or following it, or, as in this case, without definite stomatitis, the fungus may be associated with troublesome intestinal catarrh and be present in the motion to the number of one million cells per gram.

On culture media the organism often appears in its

yeast conidial form only, and may be mistaken for a saccharomyces. Thus, on isolating it from inflamed gums, Mr. K. Goadby named it *Sac. neoformans*. On meeting with it in a case of sore throat, Mr. A. G. R. Foulerton named it *Sac. tumefaciens albus*, and described it at length.¹ Grown in litmus milk or beer wort at 20° C. to 22° C. for a week, it appears in chains of elongated cells, with yeast conidia round the septa. The common form in London ferments maltose, but neither saccharose nor dextrin. Less common is one which ferments the three. Both forms liquefy glucose gelatin within one month, but they never ferment galactose, lactose, or raffinose. In textbooks it is still sometimes stated that the nature of this fungus is not understood. The work of Reess, Hansen, and Zopf set the matter at rest twenty-five years ago, and the latter named the organism *Monilia albicans*. Since then a French mycologist has referred it to a higher group of fungi, but it is certain that the organism with which he worked was not the common thrush fungus at all. Moreover, his work has not been confirmed.

REFERENCE.

¹ *Experimental Blastomycosis*, *Journ. Path. and Bact.*, 1893.

SALICYLIC IONIZATION IN ACUTE PERICARDITIS.

By J. S. MACKINTOSH, M.D., M.R.C.S., L.R.C.P.,

HAMPSTEAD.

THE value of salicylic ionization in some forms of pleurisy has already met with some appreciation by the profession. The following case apparently shows that it is also useful in acute pericarditis.

A thin, delicate girl, aged 18, developed acute rheumatism after a quinsy, and in the course of the attack acute pericarditis appeared. The pain was severe, the patient spent the whole of one night in a position of complete orthopnoea and was quite sleepless. The next day salicylic ionization was tried, the active electrode in the form of a pad soaked in a 2 per cent. solution of sodium salicylate being placed over the precordium and the indifferent electrode on the back. A current of 40 milliamperes was used for about half an hour, and had not run many minutes before the patient declared herself eased. At the end of the session the patient lay back on her pillow and slept soundly. The ionization was repeated on subsequent days, and the symptoms of pericarditis gradually subsided; aceto-salicylic acid was also being taken by the mouth. It is difficult to believe that the great and rapid relief afforded in this case was due merely to tissue-ionization by the galvanic current or to its moral effect. The patient was a favourable subject for the treatment, the thinness of her chest-wall offering a reasonable expectation that the salicylic ions could be made to penetrate to the requisite depth.

The benefit of salicylic ionization in joints the subject of acute rheumatism or sprain is well established, and the resemblance between the pathology of the pericardium and that of the synovial sac of a joint is an axiom of pathology.

FRACTURE of the clavicle, especially when incomplete, is not always easy to detect in the newborn. Sometimes it is not betrayed until the callus around the seat of fracture causes pressure effects. Dr. Azema, of Toulouse (*Bulletin de la Soc. d'Obstét. et de Gynéc. de Paris*, etc., May, 1913, p. 530), has noted two instances of this condition, though in one there was strong evidence that the injury occurred a few days before delivery. The possibility of intrauterine fracture has always to be taken into account. Dr. Azema likewise reports a fatal case of osteomyelitis developing in the fractured bone through infection from an inflamed umbilicus. *Post-mortem* examination revealed an umbilical abscess not communicating with the peritoneal cavity and general peritonitis. In this case an obstetrical fracture of the clavicle became infected from a focus elsewhere, a common seat of infection, which was noted at birth and treated appropriately by the medical officer and nurses of the maternity clinic in Toulouse where the child was born.