heart disease complicated with hysteria, but must not further refer to them. I will only say that in 13 out of the 18 sleeplessness due to restlessness or pain was a prominent symptom, and tended to aggravate the condition of the patients; 4 of the 18 were males between the ages of 18 and 30. The remainder were females from 16 to 35, with one exception, a woman of 45.

The alliance between functional and organic diseases of the nervous system claims much closer attention than I can give. As Charcot affirmed, neurasthenia is often found united with hysteria, particularly in men as a result of traumas and in women at the menopause. As regards the former class, he considered that neurasthenia was the first to evolve under the influence of the sudden violent shock, and that later the physical and mental depression prepared the way for hysteria.

My experience in the examination of men who have met with accidents leads me entirely to agree with this statement, for not a few of these subjects in a month or two after an accident, become worse rather than better, and on examination prove to be hysterical. It is difficult sometimes to estimate how far symptoms belong to hysteria or neurasthenia. In some traumatic cases, however, the neurasthenic symptoms are pretty definite—namely, psychical asthenia, characterized by diminished power of attention, want of energy, muscular weakness, constant headache or pressure in the head, and backache.

The frequent association of hysteria with the mental stigmata of degeneracy has led to the misconception that the hysterical were necessarily deceitful, that they lied for the sake of lying, that they were simulators and charlatans. As Jane¹ points out, though their mental condition tends to be modified in certain directions (as memory, will, power of attention), and their disposition becomes mobile and contradictory, their character, as the character of every one, depends upon their primary intelligence and their early surroundings and education. If time permitted I might refer to cases illustrating the occasional association of hysteria and obsessions.

I must say but little of hysteria as a complication of organic nervous diseases, and shall only refer to it as an occasional adjunct to structural diseases of the spinal cord. Dr. Buzzard directed attention to the relation of hysteria and insular sclerosis, with special reference to diagnosis, to the tendency for "functional symptoms to be mixed up with those belonging to organic change," and to the temporary intervals of improvement. Hysteria is probably more frequently associated with disseminated sclerosis than with any other affection of the cord, and this to some extent is accounted for, since this disease is frequently met with in early life and in women. For a similar reason hysteria is likely to occur in hereditary ataxia. I do not think I have met with more than half a dozen cases of this affection, but two of these—boys between the ages of 14 and 18—were distinctly hysterical. Young people with vertebral caries, too, are at times hysterical. A girl under my care had caries in the dorsal region of the spine with spastic paraplegia and an hysterical monoplegia of an arm. On the other hand, in that very common disease, locomotor ataxy, occurring as it does in adults, hysteria is more rarely found as a complication. I have only notes of four cases where it was present, and two of these were aged 30 and 34.

The recognition of hysteria in these cases accounted for certain symptoms not directly due to the organic disease, and in a few instances there was evidence that temporary improvement followed and depended on mitigation of the hysterical condition. I will only refer to one of several cases which warrants this assumption. A boy of 17 came under my care with such symptoms of disseminated sclerosis as nystagmus, intentional tremor, and also great weakness of the legs, especially the right; he was unable to walk, and he staggered and dragged the right foot. He was a degenerate with badly-shaped head, jug-handle ears, and supernumerary nipples. He had been out of work some time and was in a miserable physical condition. He suffered from occasional attacks of faintness and vomiting. There was anaesthesia of the left forearm and right leg and restriction of the fields of vision. He was treated by massage, faradic current, and cold sponging. He was soon able to walk well, and in two months left the hospital apparently in good health. At this time the only evidence of hysteria was some restriction of the visual field. He

afterwards enlisted, and at the end of two years was discharged owing to some return of his old trouble. At the age of 20 he again came under my care with the former symptoms of disseminated sclerosis. His gait was spastic, but he walked fairly well. There was now complete anaesthesia of the left side of the face and left leg. He again improved under treatment, and left the hospital to seek light employment.

And now, Mr. President and gentlemen, I must bring this lecture to a close. None can be more sensible of its defects than myself. If it should fail in arousing a little more interest in the study of hysteria as it appears in association with other diseases, the want of success in this direction will be due to my incapacity rather than to the nature of my subject.

REFERENCES.

¹ *Hystérie*, par le Dr. Voronoff, p. 65. ² Gilles de la Tourette, *Traité de la hystérie*, vol. i, p. 541. ³ Oppenheim, *Textbook of Nervous Diseases*, translated by Alexander Bruce, vol. ii, p. 1061. ⁴ *Leçons du Mardi*, 1888-9, p. 457.

THE RATIONAL TREATMENT OF CHRONIC BACILLARY DYSENTERY;

AND THE ADVANTAGES OF ENEMATA OF SILVER GELATOSE.

By LEONARD ROGERS, M.D., F.R.C.P., I.M.S.,

PROFESSOR OF PATHOLOGY, CALCUTTA.

ACUTE bacillary dysentery is essentially an inflammation of the mucous membrane of the large bowel, and the causative organisms are limited to the intestinal tract, except in a very few acute instances in which they invade the blood stream. In early cases the serum and saline treatments are usually effective in dealing with the toxæmia and congestion of the mucous membrane, and commonly subdue the disease before much lasting damage has been done.

Cases of chronic bacillary dysentery in which extensive ulceration has already been produced before the patient comes under skilled treatment are much more difficult to deal with. I have frequently found the saline treatment do more harm than good, while in one large Calcutta hospital, where several hundred cases are treated yearly, the mortality from this distressing disease is about 40 per cent. In fact, I know of no common tropical disease in India at the present time which is more difficult to deal with successfully, with the single exception of the terrible kala-azar. Contrary to the experience of some other writers, I have not had good results with stock Shiga dysentery bacillus vaccines, as recommended by Forster, and, owing to the great toxicity of Shiga's bacillus, they require to be used with caution. Sensitized dead dysentery bacillus vaccines, I think, promise better, but I have not yet been able to try them. Occasionally I have obtained good results with an autogenous dysentery vaccine, but in other cases even this failed, while the method is obviously not of general applicability, as, apart from the time and expense required to carry it out, there are few laboratories in the tropics.

In short, it may be said that the treatment of chronic bacillary dysentery is in a very unsatisfactory state at the present time, especially as compared with the formerly intractable amoebic variety of the disease.

THE CHARACTER AND DISTRIBUTION OF THE BOWEL LESIONS AND THEIR BEARING ON TREATMENT.

An analysis of considerably over one hundred *post-mortem* examinations made by me in Calcutta during the last twelve years has brought out a fact which appears to be of great practical importance in connexion with the problem under consideration. I refer to the very marked contrast in the distribution, as well as in the character, of the bowel lesions in acute and chronic fatal bacillary dysentery respectively. These features of the disease are fully described and illustrated in my recently issued small work on dysenteries, and I only need here refer to the points of importance in practice.

Briefly it may be said that in fatal acute bacillary dysentery the whole length of the large intestine and the lowest part of the ileum are uniformly involved in

a fibrinous inflammation of the mucous membrane, with only superficial ulceration, if any. Death results early from the absorption of a lethal dose of toxin, aided by the shock induced by the acute affection of such a large extent of the mucous membrane of the bowel.

On the other hand, in examinations after death from chronic bacillary dysentery—including in that category all those in which the disease has lasted from one month up to a year or more—the lesions are, in my experience, nearly always limited to the lower portions of the large intestine, and in the exceptional cases in which they extend above the descending colon they are comparatively slight in the upper part, although they involve the whole of the mucous membrane of the lower half of the large gut. Moreover, the character of the lesions is quite different from that of cases in which death has taken place in the early acute stages of the disease. In the chronic cases extensive depressed, and often serpiginous, ulcers are found on a generally thickened bowel wall, with small islets of remaining mucous membrane between them, which as a rule no longer shows the acute fibrinous deposits of the early stages. In this class of case the strength is steadily sapped by the frequent discharges of blood and mucus and the loss of absorptive powers, and although the early toxic and febrile stage has been survived—probably largely on account of the lesser extent of the bowel involved—the tendency of the disease is to run on for months, unchecked by the numerous drugs poured into the stomachs of the unfortunate patients.

The very frequent limitation of the lesions of chronic bacillary dysentery to the lower half of the large gut, as opposed to the general involvement in acute fatal cases, was new to me; but the important bearing of this fact on the treatment of the chronic disease is evident. Any drugs given by the mouth have to traverse the whole length of the gastro-intestinal canal down to the lower part of the large gut before they reach the seat of the disease. When they do arrive in the descending colon and sigmoid the irritability of the ulcerated mucous membrane is not likely to allow them to remain long in contact with the diseased parts before being expelled in one of the numerous stools. In short, but little can be expected from medicines given by the mouth, except perhaps from large doses of bismuth, some of which might help to coat and protect the ulcers.

INDICATIONS FOR THE USE OF MEDICATED ENEMATA IN CHRONIC BACILLARY DYSENTERY.

The very distribution of the lesions which is so inimical to the action of remedies administered orally is, on the contrary, of the greatest assistance to the action of enemata containing active drugs; for the lower half of the large intestine is much more readily and effectually reached in this way than the upper portion. The question of treatment, therefore, resolves itself mainly into one of ascertaining which are the best remedies for rectal administration.

One of the most frequently used and effective of these is silver nitrate, which was advocated in chronic bacillary dysentery by Stephen Mackenzie in 1882, and since used by many others, especially in the United States, although not so much in the tropics, probably on account of its failure in the numerous chronic amoebic dysenteries of warm climates. Silver nitrate, however, has two serious disadvantages in practice. The most important of these is that it is precipitated by both chlorides and albuminous substances, so that its action is very uncertain. Secondly, when used in sufficient strength to be likely to exert marked antiseptic and astringent properties, in the presence of salts and albumin, it is apt to be very painful.

EXPERIMENTAL INVESTIGATION OF THE ACTION OF VARIOUS SILVER COMPOUNDS AND OTHER ANTISEPTICS ON THE DYSENTERY BACILLUS IN THE PRESENCE OF SALTS AND ALBUMIN.

The foregoing considerations led me to undertake an investigation with a view to ascertaining the substances most likely to be useful in the treatment of chronic bacillary dysentery by means of medicated enemata, especial attention being directed to the numerous organic combinations of silver which have been so largely used in the treatment of inflammatory conditions of the urethral and conjunctival mucous membranes. The details of these

experiments will appear in the second number of the *Journal of Indian Medical Research*, so only the results need be mentioned here in so far as they bear on the practical treatment of the disease. Each substance was tested against Shiga's bacillus in two sets of dilutions—one dissolved in water, and a second in which half the diluting fluid was sterile broth containing both chlorides and albumins. The importance of the second set will be evident from the fact that whereas silver nitrate killed the dysentery bacillus within five minutes (the time adopted in all the tests) up to a dilution of 1 in 10,000, yet in the presence of a little broth it failed to kill in a dilution of 1 in 100 in one experiment, and 1 in 500 in another, being at once precipitated by the broth.

On the other hand, I found that those organic silver compounds which were not precipitated by broth had a much more marked action against the dysentery bacillus in the broth series of tubes than had silver nitrate, although less active than the latter in water. Of the organic silver substances silver gelatose (albagin) gave the best results in broth, killing the Shiga bacillus in five minutes up to dilutions of 1 in 500 and 1 in 1,000 respectively in two trials. Nargol came next, acting up to 1 in 500. Protargol and mercuriol (mercury nucleate) were nearly as active, but it would not appear to be advisable to inject large quantities of a mercury compound into the bowel in emaciated dysenteric subjects. On the other hand, argentamin had little action, while collargol, ichthargan, and argyrol were practically inert in the presence of broth, as were copper sulphate and cuprol (copper nucleate).

A second series of experiments was carried out in a similar manner with certain antiseptics, with the following results: Cyllin was active in broth up to 1 in 2,500. Iodine, dissolved with the aid of iodide of potassium, and izal both proved effective up to 1 in 1,000, but lysol was less active in the one test made. Permanganates, with or without the addition of a little dilute sulphuric acid, quinine bisulphate, and boracic acid, all proved to be nearly inert in the presence of broth.

It would thus appear that under the conditions noted albagin was the best of the silver compounds tested, being closely followed by nargol and protargol, while cyllin, iodine, and izal were the most active of the other antiseptics tested.

THE PRACTICAL APPLICATION OF THE ABOVE INVESTIGATION.

In making use of the foregoing data in the treatment of chronic bacillary dysentery we are first confronted with the differential diagnosis of this class from more especially the amoebic form. The clinical differences are dealt with in my book, but they are not sufficiently characteristic and constant to enable them to be completely relied on for diagnostic purposes. Of greater value is the absence of pathogenic amoebae from the stools after repeated examination, as they may not be found at a single attempt in chronic amoebic disease. The isolation of the dysentery bacillus is a much more difficult procedure, even when a well equipped laboratory is available, and often fails in experienced hands in chronic bacillary cases, as subsequently verified *post mortem*. Fortunately we have a simple clinical test for excluding the amoebic variety in the use of emetine, for if 1-grain doses of this drug given hypodermically on three or four successive days has no effect on the disease, amoebic infection may safely be excluded and the case treated as one of chronic bacillary dysentery in the absence of signs or microscopical evidence of other rarer forms of lower bowel disease.

During the last few months I have treated a number of non-amoebic cases in the light of the above experiments, with sufficiently promising results to deserve record so as to allow of others giving them a trial in intractable chronic bacillary dysentery.

DYSENTERY CASES TREATED WITH ENEMATA OF ALBAGIN OR SILVER GELATOSE.

My experiments having shown albagin to be the most effective of the organic silver compounds against the dysentery bacillus in the presence of salts and organic matter, I proceeded to try it in dysentery cases other than the amoebic variety. The results have been distinctly promising, while the solution has the great advantage of being

painless, and usually easily retained for a sufficient length of time to have a good chance of exerting its disinfectant and astringent properties. The following brief notes will serve to illustrate the effects of albargin in different types of cases.

CASE I.—Acute Bacillary Dysentery Resisting Purgative Treatment, but Cured by Three Albargin Bowel Washes.

An Indian Christian male, aged 38, admitted for suspected cholera, but proved to be a case of acute bacillary dysentery. From eleven to seventeen stools daily, containing much blood and mucus. No improvement after eight days' treatment with castor oil and opium, followed by salines. One pint of albargin of the strength of 1 grain in 1 oz. (approximately 1 in 500 solution), given as an enema, and increased to 1½ pints on each of the following two days. The evening temperature, which had been rising to 99° and a little over, fell to 97°; the blood disappeared at once, and the mucus finally after six days, by which time the stools had become reduced to one or two healthy motions daily, and he was on solid food. He was discharged four days later, and came to see me several times during the next three months, and has remained well. The rapid improvement after the failure of other treatment in a severe case, which was threatening to become chronic, was striking.

CASE II.—Chronic Bacillary Dysentery of Eight Months' Duration Rapidly Cured by Albargin.

European female, aged 17, admitted for chronic dysentery for eight months without intermission. No amoebae found on repeated examination of the stools, which contained blood and mucus and numbered three to four daily, being passed with gripping pains. History of fever at first, but none now. Treated with castor oil emulsion and bismuth salicylate for five days without improvement. Then one and a half pints of one grain to the ounce albargin solution given as an enema on three successive days, and retained for from fifteen to thirty minutes without pain. The blood and mucus disappeared after the second injection, and one healthy stool was passed daily during the following week, although solid diet was given two days after the last bowel wash. No relapse occurred, and she has been seen repeatedly during the two months following her discharge from hospital and has remained in good health. Such a rapid recovery after eight months' suffering is very encouraging.

CASE III.—Severe Bacillary Dysentery in a Young Child Cured by Albargin.

A European child, aged 2½, had been treated for severe dysentery, with up to twenty stools a day, for twelve days with castor oil, emetine, and saline enemata. No material improvement had resulted and the condition of the child was becoming serious. Captain Green-Armitage, I.M.S., now asked me to see the case, and I advised that 8 oz. enemata of albargin of the strength of 1 grain to 1 oz. should be given. Improvement began after the first injection, and shortly after a second, four days later, no further blood or mucus was passed, but a third was given as a precautionary measure. The child rapidly picked up and was discharged cured two weeks after the commencement of the albargin treatment. The absence of pain after the injections in so young a child is of interest, as well as the immediate improvement after the failure of other treatment.

The foregoing cases will suffice to illustrate the value of the new treatment in certain cases of bacillary dysentery, while the following is a brief summary of the results obtained in a consecutive series of all the cases so far treated in my ward. They number twelve, including those already recorded above. Four acute cases of less than two weeks' duration on admission were all rapidly cured by from three to five albargin enemata, although two of them had suffered from other attacks within the previous few months. Of the 8 chronic cases 5 were of from one to four months' duration, 1 of eight months and 2 of sixteen months. Of these 8 cases, one patient left hospital two days after the treatment was begun, so should be excluded. One of sixteen months' duration improved considerably, but the drug failed to cure him, and he is still in hospital. The remaining 6 were cured, from three to five injections being given in 5 of them, while a larger number were required in the sixteen-months' case, who had previously been treated unsuccessfully with nargol enemata. As all these cases had resisted the usual treatment with castor-oil and salines for an average of seven days (mild cases clearing up under purgative treatment not being given bowel washes), these results are distinctly encouraging, and are certainly better than I have previously obtained in similar cases by older methods.

Lastly, it should be mentioned that albargin bowel washes failed to do good in two patients, who were

subsequently found to have pathogenic amoebae in their stools, although on their admission amoebae had not been found, so that the silver gelatose appears to be useless in amoebic cases, although of great value in bacillary dysentery.

I have not yet been able to test the value of the other antiseptics which my experimental investigation show to be worthy of trial in non-amoebic dysentery. The main object of this communication is to draw attention to the principle that medicated enemata are clearly indicated in the treatment of chronic bacillary dysentery, owing to the lesions being usually limited to the lower half of the large intestine, and consequently easily reached by bowel washes.

GASTRIC TETANY IN THE ADULT.

BY

W. K. IRWIN, M.D. ABERD.,
LONDON.

In 1869 Kussmaul drew attention to the occurrence of tetany in cases of chronic dilatation of the stomach. Since that date about sixty cases of a similar nature have been recorded in adults. In the majority the dilatation of the stomach was due to the cicatrization of a chronic ulcer in or near the pylorus, but in a few tetany has been observed to follow dilatation of the stomach due to other causes than ulceration. Thus, in a case recorded by Trevelyan, it occurred as a complication in a carcinomatous stricture of the duodenum, and in a case recorded by Fleiner it was associated with a sarcomatous growth of the stomach, one attack being immediately fatal. Also, tetany has ensued from dilatation of the stomach due to twisting of the duodenum, pressure of an enlarged gall bladder, and pressure on the duodenum by a pancreatic cyst. The most prominent feature of this disease is the occurrence of a tonic spasm of the muscles of the upper and lower limbs, but sometimes intermittent spasm of the jaw, neck, and back muscles is present; these attacks last from a few minutes to half an hour, and disappear as suddenly as they came. During their continuance the patient is unable to open the mouth or to swallow, and in several instances opisthotonos was a well-marked symptom.

In other cases the tetany is followed by convulsions that closely resemble epilepsy, the fits are usually repeated in rapid succession. In typical cases the wrists and elbows are partially flexed, and the forearms strongly pronated; the fingers are bent over the thumbs, flexion being most marked at the carpo-phalangeal joints; the palms are hollowed by the approximation of the thenar and hypothenar eminences, and therefore the hand becomes cone-shaped (the accoucheur's hand). In the lower extremities the legs are rigidly extended, the heels drawn up, the soles of the feet turned inwards, and the toes are bent downwards and adducted. The patient often complains of pain while the spasm lasts, and the affected parts are blue and cold. The deep reflexes are much increased, the superficial reflexes may also be exaggerated, and the muscles react very readily to the interrupted current. The patient generally remains quite conscious during an attack. The breathing is quickened, the pulse full and regular. The temperature is at first often depressed, but in fatal cases it often rises to 108° F. before death. The pupils are often contracted, but react both to light and accommodation. Cutaneous sensibility rarely undergoes any decided alteration, but in two or three of the recorded cases transient hyperaesthesia was observed. Retention of urine usually occurs, and the urine often contains a small quantity of albumin. Occasionally a trace of sugar or acetone is present. Delirium, rise of temperature, excess of albumin in the urine, coma with dilatation of the pupils, and the presence of tetanic or epileptiform seizures are all evil signs. In almost every case the first attack is soon followed by several others. The duration of the spasm may be from four or five minutes to several hours, but it may remain almost constant for two or three days. It often follows an attack of vomiting or diarrhoea, and a spasm may sometimes be induced by percussing the epigastrium, by passage of a stomach tube, or by pressure on the main artery of a limb. Sometimes patients have