Published in final edited form as:

Tetrahedron Lett. 2007 February 12; 48(7): 1291–1294.

Coupling of activated esters to gramines in the presence of ethyl propiolate under mild conditions

David T. Jones, Gerald D. Artman III, and Robert M. Williams*

Department of Chemistry, Colorado State University, Fort Collins, CO 80523, USA University of Colorado Cancer Center, Aurora, CO 80045

Abstract

The coupling of activated esters to gramine derivatives is described using ethyl propiolate. A series of substrates have been prepared using these mild conditions to provide a scope and limitations for this methodology.

Keywords

indoles; tryptophans; gramines; malonates; indolines

During the course of the total syntheses of several natural products and potential biomimetic precursor substrates, we have needed to couple functionalized activated esters to gramine derivatives of various indole ring systems. Three literature methods can be used to form the desired carbon-carbon bond. First, heating of the gramine **1** to temperatures greater than $>120^{\circ}\text{C}$ leads to spontaneous formation of the 3-methylene-3*H*-indole (**2**) via loss of the dimethyl amine (Scheme 1).¹ Trapping of **2** with nucleophiles such as a malonate enolate or cyanide generates the desired 3-substituted indole **3**. This procedure can result in good yields of the coupled product, but with the high temperatures employed may not be suitable for sensitive substrates. Alternatively, Somei² and Kametani³ independently reported that the temperature can be reduced to $\sim 90^{\circ}\text{C}$ by the addition of 0.4 equivalents of *tri-n*-butylphosphine. The precise role of the phosphine catalyst is still unclear at this time. However, one mechanistic hypothesis is that it acts as nucleophilic catalyst forming a highly reactive phosphonium salt, which is more susceptible to elimination and formation of **2**. As a milder alternative to the thermal conditions, the indole nitrogen of the gramine **1** can be protected as the corresponding *N*-silyl gramine **4**.⁴ Exposure of **4** to MeI or Me₂SO₄ generates a quaternary ammonium salt, which is readily eliminated in the presence of TBAF to generate **2**, which is trapped as before to yield **3**. These reaction conditions are quite general for a variety of substrates, but the use of a fluoride source to generate intermediate **2** makes it incompatible with substrates that require silyl protecting groups elsewhere in the molecule.

Previously, our laboratory has successfully employed the conditions developed by Somei and Kametani for the preparation of substituted indole derivatives utilized in the synthesis of the brevianamides and paraherquamides.⁵

* Corresponding author. Tel.: (970)-491-6747; fax: (970)-491-3944; e-mail: rmw@chem.colostate.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Unfortunately, as our malonate and indole substrates have become more complex the yields for this condensation reaction can be quite variable and range from good (~70%) to poor (~30%). As result, we have examined alternative conditions for this key bond-forming reaction and were drawn to a report by Sainsbury, who demonstrated the facile coupling of gramine and an activated ester at room temperature using activated acetylenes.⁶ Activation of gramine **5** occurs *via* conjugate addition on dimethyl acetylenedicarboxylate (DMAD, **6**) generating the zwitterion **7** (Scheme 2). Loss of the ammonium species *via* elimination leads to formation of the enamine **8** and the reactive 3-methylene-3*H*-indole (**2**). Trapping of the anion of *N*-formyl diethyl aminomalonate (**9**) generates the desired product **10** in 90% yield after only 15 min. With only two examples of this reaction reported by Sainsbury, we have performed a series of experiments to explore the scope and limitations of these conditions in effort to better understand how this mild coupling method can be applied to natural product synthesis.⁷

In the original report by Sainsbury, the only activated acetylene employed was DMAD. As such, we commenced our experiments by screening a panel of commercially available activated acetylenes (Table 1).^{8,9} Due to the speed of the reaction, two important details were quickly ascertained from the coupling of gramine (**5**) to diethyl malonate (**11**) in THF at rt. First, acetylenes possessing a single carboxylate group (entries *a*, *b* and *c*) generated **12** in moderate to good yields (47%-78%). Second, once the acetylene was di-substituted with a carboxylate group at one terminus and an alkyl or aryl group at the other, no reaction was observed between the malonate and the gramine (entries *e-h*). It is postulated that these results are observed due to the acetylene becoming: 1) more electron-rich and therefore less susceptible to conjugate addition (entries *d*, *e* and *f*); and 2) more sterically encumbered (entries *g* and *h*). This initial screening of reagents revealed that commercially available ethyl propiolate readily affords the alkylated malonate **12** in good yield (entries *i-l*).

With the reagent screen completed, we began to explore the influence of different solvents on this reaction. Our initial reagent screen was conducted in THF, the same solvent employed by Sainsbury. We have found that performing the coupling in different solvents can significantly influence the yield of this reaction. Switching from THF to CH₃CN (entry *i*) or CH₂Cl₂ (entry *j*) increased the yield from 78% to 85% and 86%, respectively. Interestingly, a protic solvent like MeOH (entry *k*) also generates the desired product, albeit in a depressed 60% yield. However, Et₂O (entry *l*) proved to be the best solvent for this reaction resulting in 90% yield of the alkylated product **12** after 15 minutes at room temperature.

Based on these preliminary results, we alkylated a series of malonate derivatives (**13**) with gramine (**5**) in the presence of ethyl propiolate in less than 20 min (Table 2).¹⁰ As a result, several trends were observed. First, the activated esters (**13**) needed to have a pK_a lower than 16 in DMSO for good to excellent yields of the alkylated product (entries *a-d*). Second, alkylations of activated esters with low pK_a's like ethyl cyanomalonate (entry *a*) and ethyl nitroacetate (entry *d*) were often accompanied by small amounts of dialkylated products (<20%) thereby depressing the yield of the monoalkylated product **14**. Malonates that contain bulky substituents like *N*-Boc diethyl aminomalonate (entry *b*) also gave lower yields (65%) presumably due to increased steric interactions. Interestingly, coupling of the benzophenone imine of glycine (entry *e*) did afford product, but in a disappointing 41% yield. We believe this example represents the upper limit of the pK_a range required for this methodology (pK_a = 19.5). This is further confirmed by the failed alkylation of *N*-Boc ethyl glycine (entry *f*) and ethyl azidoacetate (entry *g*), both of which have pK_a's greater than 20.

In addition to the scope of the activated ester, we explored the influence of indole ring system of the gramine and its ability to couple to the activated ester in the presence of ethyl propiolate. Commercially available indoles were converted to their respective gramines **15** using known literature procedures (CH₂O, Me₂NH, AcOH, rt) in good to excellent yields. Exposure of the

gramine **15** to *N*-Boc diethyl aminomalonate (**13b**) in the presence of ethyl propiolate in Et₂O readily generated the desired tryptophan derivatives **16** (Table 3). Both electron-poor and electron-rich indole ring systems couple under these conditions in good yields. In addition, sterically demanding C2-substituted gramines (Table 3, entry *g*) readily afford the tryptophan derivative **16g** in 62% yield. The coupled products **16** can readily be converted to the racemic tryptophan derivatives using known literature methods.¹¹

In conclusion, we have studied the scope and limitations on the mild coupling of activated esters to gramines using the inexpensive reagent ethyl propiolate. In comparison to similar reactions, this carbon-carbon bond-forming event takes place at room temperature in moderate to good yields in a several common organic solvents using many different activated esters and gramines. Finally, we are attempting to develop an asymmetric variant of this coupling reaction for the enantioselective synthesis of tryptophans analogues readily derived from various gramines.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

The authors acknowledge financial support from the National Institutes of Health (CA70375) and the NSRA Postdoctoral Fellowship for GDA (GM72296). Mass spectra were obtained on instruments supported by the NIH Shared Instrumentation Grant GM49631.

References

1. For the generation of 3-methylene-3*H*-indoles under thermal conditions, see: (a) Diker K, Maindreville D, Lévy J. *Tetrahedron Lett* 1999;40:7459–7462. (b) Quick J, Saha B, Driedger PE. *Tetrahedron Lett* 1994;35:8549–8552. (c) Kozikowski AP, Okita M, Kobayashi M, Floss HG. *J Org Chem* 1988;53:863–869.
2. Somei M, Karasawa Y, Kaneko C. *Heterocycles* 1981;16:941–949.
3. Kametani T, Kanaya N, Ihara M. *J Chem Soc Perkin Trans* 1981;1:959–963.
4. For recent examples of the Somei-Kametani coupling, See: (a) Novikov AV, Sabahi A, Nyong AM, Rainier JD. *Tetrahedron:Asymmetry* 2003;14:911–915. (b) Freed JD, Hart DJ, Magomedov NA. *J Org Chem* 2001;66:839–852. [PubMed: 11430103] (c) Boyles DA, Nichols DE. *J Org Chem* 1988;53:5128–5130.
5. For representative examples using *N*-silyl indoles to generate 3-methylene-3*H*-indoles, see: (a) Crawley SL, Funk RL. *Org Lett* 2006;8:3995–3998. [PubMed: 16928057] (b) Pérez-Serrano L, Casarrubios L, Domínguez G, Freire G, Pérez-Castells J. *Tetrahedron* 2002;58:5407–5415. (c) Meseguer B, Alonso-Díaz D, Griebenow N, Herget T, Waldmann H. *Chem Eur J* 2000;6:3943–3957. (d) Iwao M, Ishibashi F. *Tetrahedron* 1997;53:51–58.
6. For representative examples from our laboratory, see: (a) Williams RM, Cao J, Tsujishima H, Cox RJ. *J Am Chem Soc* 2003;125:12172–12178. [PubMed: 14519003] (b) Cushing TD, Sanz-Cervera JF, Williams RM. *J Am Chem Soc* 1996;118:557–578. (c) Williams RM, Glinka T, Kwast E, Coffman H, Stille JK. *J Am Chem Soc* 1990;112:808–821.
7. (a) Hogan I, Jenkins P, Sainsbury M. *Tetrahedron Lett* 1988;29:6505. (b) Hogan I, Jenkins P, Sainsbury M. *Tetrahedron* 1990;46:2943.
8. For an alternative application of activated actelyenes and gramines, see: Voskressensky LG, Borisova TN, Kulikova LN, Varlamov AV, Catto M, Altomare C, Carotti A. *Eur J Org Chem* 2004:3128–3135.
9. All reagents were used without further purification from commercial sources. For detailed experimental procedures, see Supporting Material section below.
10. **GENERAL PROCEDURE:** The gramine (1.1 equiv.) is suspended in Et₂O (0.2M). The activated ester (1.0 equiv.) is added in a single portion followed by ethyl propiolate (1.1 equiv.) via syringe. The reaction mixture is stirred until no more starting material is visible by TLC (~20 min). The

reaction mixture is poured into a separatory funnel containing EtOAc. The organic solution is washed with 1N HCl, water and Brine. The organic layer is dried over Na₂SO₄ and the volatile organics removed under reduced pressure. The crude product is purified by flash silica gel chromatography.

11. (a) Marcq V, Mirand C, Decarme M, Emonard H, Hornebeck W. *Bioorg Med Chem Lett* 2003;13:2843–2846. [PubMed: 14611841] (b) Endo Y, Shudo K, Itai A, Hasegawa M, Sakai S. *Tetrahedron* 1986;42:5905–5924.

Scheme 1.

Two methods for the coupling of gramines to activated esters.

Scheme 2.

DMAD mediated coupling of gramine to *N*-formyl diethylaminomalonate.

Table 1

Alkylation of gramine (**5**) with diethyl malonate (**11**) in the presence of commercially available activated acetylenes.

Entry	R ₁	R ₂	Solvent	Yield of 12
<i>a</i>	H	Et	THF	78%
<i>b</i>	H	Me	THF	77%
<i>c</i>	H	<i>tert</i> -Bu	THF	47%
<i>d</i>	<i>n</i> -propyl	Me	THF	0%
<i>e</i>	Me	Et	THF	0%
<i>f</i>	Et	Et	THF	0%
<i>g</i>	Ph	Et	THF	0%
<i>h</i>	Ph	Me	THF	0%
<i>i</i>	H	Et	CH ₃ CN	85%
<i>j</i>	H	Et	CH ₂ Cl ₂	86%
<i>k</i>	H	Et	MeOH	60%
<i>l</i>	H	Et	Et ₂ O	90%

Table 2Coupling of gramine (**5**) with different activated esters **13** in the presence of ethyl propiolate.

Entry	Ester Derivatives	Yield of 14
<i>a</i>	$\text{R}_1 = \text{CN}$, $\text{R}_2 = \text{H}$	67%
<i>b</i>	$\text{R}_1 = \text{CO}_2\text{Et}$, $\text{R}_2 = \text{NHBoc}$	65%
<i>c</i>	$\text{R}_1 = \text{Ac}$, $\text{R}_2 = \text{H}$	83%
<i>d</i>	$\text{R}_1 = \text{NO}_2$, $\text{R}_2 = \text{H}$	65% ^{<i>l</i>}
<i>e</i>	$\text{R}_1 = \text{N}=\text{CPh}_2$, $\text{R}_2 = \text{H}$	41%
<i>f</i>	$\text{R}_1 = \text{NHBoc}$, $\text{R}_2 = \text{H}$	0%
<i>g</i>	$\text{R}_1 = \text{N}_3$, $\text{R}_2 = \text{H}$	0%

^{*l*} Due to solubility problems, this coupling was conducted in CH_2Cl_2

Table 3

Coupling of *N*-Boc diethyl aminomalonate (**13b**) with various indole derivatives of gramine **15** in the presence of ethyl propiolate.

Entry	Gramine	Yield of 16
<i>a</i>		54%
<i>b</i>		44%
<i>c</i>		65%
<i>d</i>		42%
<i>e</i>		53%
<i>f</i>		61%
<i>g</i>		62%